

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 March 2003 (20.03.2003)

PCT

(10) International Publication Number
WO 03/022219 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US02/28731
- (22) International Filing Date:
9 September 2002 (09.09.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/950,543 11 September 2001 (11.09.2001) US
- (71) Applicant: **ALBANY MOLECULAR RESEARCH, INC.** [US/US]; 21 Corporate Circle, Albany, NY 12212-5098 (US).
- (72) Inventor: **TROVA, Michael, Peter**; 3002 Morgan Court, Schenectady, NY 12306 (US).
- (74) Agents: **GOLDMAN, Michael, L.** et al.; Nixon Peabody LLP, P.O. Box 31051, Clinton Square, Rochester, NY 14603-1051 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 03/022219 A2

(54) Title: NITROGEN SUBSTITUTED BIARYL PURINE DERIVATIVES AS POTENT ANTIPROLIFERATIVE AGENTS

(57) Abstract: The compounds of the present invention are 2,6,9-trisubstituted purine derivatives which are inhibitors of cyclin/cdk complexes. The compounds of the current invention also are potent inhibitors of human cellular proliferation. As such, the compounds of the present invention constitute pharmaceutical compositions with a pharmaceutically acceptable carrier. Such compounds are useful in treating a disorder mediated by elevated levels of cell proliferation in a mammal compared to a healthy mammal by administering to such mammal an effective amount of the compound. Examples of the compounds of the present invention are represented by the following chemical structures; with X, Y, D, Q, V, A, R₁, R₂, R₃, R₄, and n₁ defined herein.

NITROGEN SUBSTITUTED BIARYL PURINE DERIVATIVES AS POTENT ANTIPROLIFERATIVE AGENTS

FIELD OF THE INVENTION

5

The present invention relates to compounds that are shown to be potent cyclin/cyclin dependent kinase (cdk) inhibitors. Compounds with these properties are shown to be potent inhibitors of cell growth and proliferation. Such compounds can be used to treat the following conditions in mammals: rheumatoid arthritis, lupus, type 1 diabetes, multiple sclerosis, cancer, restenosis, gout and other proliferative diseases involving elevated levels of cell proliferation compared to healthy mammals. Compounds of the present invention which are biaryl substituted purine derivatives are shown to be potent antiproliferative agents against a number of human transformed cell lines, and also inhibitors of human cyclin/cdk kinase complexes.

15

BACKGROUND OF THE INVENTION

Cellular Proliferation and Cancer.

The disruption of external or internal regulation of cellular growth can lead to uncontrolled proliferation and in cancer, tumor formation. This loss of control can occur at many levels and, indeed, does occur at multiple levels in most tumors. Further, although tumor cells can no longer control their own proliferation, they still must use the same basic cellular machinery employed by normal cells to drive their growth and replication.

25

Cyclin Dependent Kinases and Cell Cycle Regulation.

Progression of the normal cell cycle from the G1 to S phase, and from the G2 phase to M phase is dependent on cdks (Sherr, C.J., Science 274:1672-1677 (1996)). Like other kinases, cdks regulate molecular events in the cell by facilitating the transfer of the terminal phosphate of adenosine triphosphate (ATP) to a substrate protein. Isolated cdks require association with a second subunit, called cyclins (Desai et al., Mol. Cell. Biol., 15:345-350 (1995)). Cyclins cause conformational changes at the cdk active site, allowing ATP access and interaction with the substrate protein.

- 2 -

The balance between its rates of synthesis and degradation controls the level of each cyclin at any point in the cycle (Elledge, S.J., et al., Biochim. Biophys. Acta 1377:M61-M70 (1998)). The influences of cyclin/cdk activity on the cell cycle and cellular transformation are summarized in Table 1.

5

Abnormal Cyclin/cdk Activity in Cancer.

In a normal cell, interlocking pathways respond to the cell's external environment and internal checkpoints monitor conditions within the cell to control the activity of cyclin/cdk complexes. A reasonable hypothesis is that the disruption of normal control of cyclin/cdk activity may result in uncontrolled proliferation. This hypothesis appears to hold in a number of tumor types in which cyclins are expressed at elevated levels (Table 1). Mutations in the genes encoding negative regulators (proteins) of cyclin/cdk activity are also found in tumors (Larsen, C.-J., Prog. Cell Cycle Res., 3:109-124 (1997)); (Kamb, A., Trends in Genetics, 11:136-140 (1995)).

Members of the Cip family of cdk inhibitors form a ternary complex with the cyclin/cdk and require binding to cyclinA, cyclinE, or cyclinD (Hall, M., et al., Oncogene, 11:1581-1588 (1995)). In contrast, Ink family members form a binary complex with cdk4 or cdk6 and prevent binding to cyclinD (Parry, D.; et al., EMBO J., 14:503-511 (1995)).

20

Table 1. Associations Among Cyclins and Cancers

Cyclin	Cell Cycle Role	Associated cdk	Cancer
A	S, G2 to M	cdk1, cdk2	hepatocellular carcinoma (Wang, J.; et al., <u>Oncogene</u> , 8:1653-1656 (1992))
B1/B2	G2 to M	cdk1	none yet defined
D1	G1	cdk4, cdk6	parathyroid adenoma (Motokura, T., et al., <u>Nature</u> , 350:512-515 (1991)) centrocytic B cell lymphoma (Withers, D.A., et al., <u>Mol. Cell. Biol.</u> , 11:4846-4853 (1991)) esophageal carcinoma (Jiang, W., et al., <u>Cancer Res.</u> , 52:2980-2983 (1992)) breast cancer (Dickson, C., et al., <u>Cancer Lett.</u> , 90:43-50 (1995)) squamous cell carcinoma (Bartkova, J., et al., <u>Cancer Res.</u> , 55:949-956 (1995)) hepatocellular carcinoma (Nishida, N., et al., <u>Cancer Res.</u> , 54:3107-3110 (1994))
D2	G1	cdk4, cdk6	colorectal carcinoma (Leach, F.S., et al., <u>Cancer Res.</u> , 53:1986-1989 (1993))
E	G1 to S	cdk2	breast cancer (Keytomarsi, K., et al., <u>Cancer Res.</u> , 54:380-385 (1994)) gastric carcinoma (Akama, Y.; et al., <u>Jap. J. Cancer Res.</u> , 86:617-621 (1995)) colorectal carcinoma (Kitihara, K.; et al., <u>Int. J. Cancer</u> , 62:25-28 (1995))

5 Inhibitors of Cyclin/cdk Complexes as Potential Anticancer Agents.

Tumors with elevated cyclin/cdk activity, whether from the over expression of cyclins or the loss of an endogenous cdk inhibitor, are prime targets for potential therapies based on small molecule cyclin/cdk inhibitors. In fact, several small molecule inhibitors of cyclin/cdks are reported (Meijer, L., et al., "Progress in
10 Cell Cycle Research," Plenum Press: New York, 351-363 (1995)) and appear to bind at the ATP site of the kinase. Some information is known about small molecule inhibitors of other kinases, such as PKC (serine kinase) (Murray, K.J. et al., "Ann. Rep. Med. Chem.," J. Bristol, Ed., Academic Press, Inc.: New York, Chapter 26 (1994)) and tyrosine kinases (Fantl, W.J., et al., Ann. Rev. Biochem., 62:453 (1993);
15 Burke, T.R., Drugs of the Future, 17:119-1131 (1992); Dobrusin, E.M. et al., "Ann.

Rep. Med. Chem," J. Bristol, Ed., Academic Press, Inc.: New York, Chapter 18 (1992); Spence, P., Curr. Opin. Ther. Patents, 3:3 (1993)). A number of known inhibitors were obtained from commercial sources or were synthesized by literature procedures.

5

Purine Compounds as Cyclin/cdk Inhibitors.

There are several reports of 2,6-diamino substituted purine derivatives as cyclin/cdk inhibitors and as inhibitors of cellular proliferation. Among those are reports by U.S. Patent No. 5,583,137 to Coe, et al., olomoucine (Vesely, J., et al., Eur. J. Biochem., 224:771-786 (1994)), roscovitine (Meijer, L., Eur. J. Biochem., 243:527-536 (1997)), WO 97/16452 to Zimmerman, Imbach, P., et al., Bioorg. Med. Chem. Lett., 9:91-96 (1999), Norman, T.C., et al., J. Amer. Chem. Soc., 118:7430-7431 (1996), Gray, N.S., et al., Tetrahedron Lett., 38:1161-1164 (1997), Gray, N.S., et al., Science, 281:533-538 (1998), WO 98/05335 to Lum, et al., Schow, S.R., et al., Bioorg. Med. Chem. Lett., 7:2697-2702 (1997), US Patent No., 5,886,702 to Mackman, et al., Nugiel, D.A., et al., J. Org. Chem., 62:201-203 (1997), and Fiorini, M.T. et al., Tetrahedron Lett., 39:1827-1830 (1998). Many of these reported compounds are shown to inhibit cyclin/cdk complexes and have modest cellular proliferation inhibition properties.

20

The compounds of the present invention are shown to have far superior biological activities as cyclin/cdk complex inhibitors as well as inhibitors of cellular proliferation compared to those previously reported. In fact, the art (e.g., Fiorini, M.T. et al., Tetrahedron Lett., 39:1827-1830 (1998)) teaches away from compounds of this invention, claiming lack of cellular proliferation inhibition.

25

SUMMARY OF THE INVENTION


The compounds of the present invention are 2,6,9-trisubstituted purine derivatives which are inhibitors of cyclin/cdk complexes. The compounds of the current invention also are potent inhibitors of human cellular proliferation. As such, the compounds of the present invention constitute pharmaceutical compositions with a pharmaceutically acceptable carrier. Such compounds are useful in treating

30

- 5 -

conditions in a mammal mediated by elevated levels of cell proliferation compared to a healthy mammal by administering to such mammal an effective amount of the compound.

In one embodiment, the compounds of the present invention are
 5 represented by the chemical structure found in Formula I



Formula I

10 wherein:

R₁ are the same or different and independently selected from the group consisting of:

- H;
- C₁-C₆-straight chain alkyl;
- 15 C₂-C₆-straight alkenyl chain;
- C₃-C₆-branched alkyl chain;
- C₃-C₆-branched alkenyl chain;
- C₃-C₇-cycloalkyl;
- CH₂-(C₃-C₇-cycloalkyl);
- 20 CH₂CF₃;
- CH₂CH₂CF₃; and
- CH(CF₃)₂;

the combination of X, D, and Q are either:

- 25 D=Q=N, and X=CH; or
- D=X=N, and Q=CH; or
- Q=X=N, and D=CH; or
- Q=N, and D=X=CH;

- 30 V= NH;
- O;
- S; or
- CH₂;

- 35 R₂= phenyl;

- 6 -

substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F , Cl , Br , CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;

5 1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;

3-pyridyl;

10 4-pyridyl;

2-pyrimidyl;

4-pyrimidyl;

5-pyrimidyl;

thiophene-2-yl;

15 thiophene-3-yl;

2-furanyl;

3-furanyl;

oxazol-2-yl;

oxazol-4-yl;

20 oxazol-5-yl;

thiazol-2-yl;

thiazol-4-yl;

thiazol-5-yl;

imidazol-2-yl;

25 imidazol-4-yl;

pyrazol-3-yl;

pyrazol-4-yl;

isoxazol-3-yl;

isoxazol-4-yl;

30 isoxazol-5-yl;

isothiazol-3-yl;

isothiazol-4-yl;

isothiazol-5-yl;

1,3,4-thiadiazol-2-yl;

35 benzo[b]furan-2-yl;

benzo[b]thiophene-2-yl;

2-pyrrolyl;

3-pyrrolyl;

1,3,5-triazin-2-yl;

40 pyrazin-2-yl;

pyridazin-3-yl;

pyridazin-4-yl;

2-quinolinyl;

3-quinolinyl;

45 4-quinolinyl;

1-isoquinolinyl;

3-isoquinolinyl; and

4-isoquinolinyl; or

- 7 -

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

5 R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 10 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

R₄= H;
 15 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

20 n₁= 0-3;

n= 0-3;

25 A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;


30 Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 35 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;

40 R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 45 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention is directed to a compound of the following formula:



Formula III

wherein:

10 R_1 are the same or different and independently selected from the group consisting of:

H;
 C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 15 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 20 CH₂CH₂CF₃; and
 CH(CF₃)₂;

the combination of X, D, and Q are either:

25 D=Q=N, and X=CH; or
 D=X=N, and Q=CH; or
 Q=X=N, and D=CH; or
 Q=N, and D=X=CH;

30 V= NH;
 O;
 S; or
 CH₂;

R₂= phenyl;

- 9 -

substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

5 1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;

3-pyridyl;

10 4-pyridyl;

2-pyrimidyl;

4-pyrimidyl;

5-pyrimidyl;

thiophene-2-yl;

15 thiophene-3-yl;

2-furanyl;

3-furanyl;

oxazol-2-yl;

oxazol-4-yl;

20 oxazol-5-yl;

thiazol-2-yl;

thiazol-4-yl;

thiazol-5-yl;

imidazol-2-yl;

25 imidazol-4-yl;

pyrazol-3-yl;

pyrazol-4-yl;

isoxazol-3-yl;

isoxazol-4-yl;

30 isoxazol-5-yl;

isothiazol-3-yl;

isothiazol-4-yl;

isothiazol-5-yl;

1,3,4-thiadiazol-2-yl;

35 benzo[b]furan-2-yl;

benzo[b]thiophene-2-yl;

2-pyrrolyl;

3-pyrrolyl;

1,3,5-triazin-2-yl;

40 pyrazin-2-yl;

pyridazin-3-yl;

pyridazin-4-yl;

2-quinolinyl;

3-quinolinyl;

45 4-quinolinyl;

1-isoquinolinyl;

3-isoquinolinyl; and

4-isoquinolinyl; or

- 10 -

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

5 n= 0-3;

 A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
10 OCH₂CH₂; or
 CHCH₃;

 Y= H;
 OR₁;
15 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
20 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;

R₃ are the same or different and independently selected from the group consisting of:

25 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
30 above in R₂;


 R₅= C₃-C₇-cycloalkyl;

 R₆ = C₁-C₄-straight chain alkyl;
35 C₃-C₄-branched chain alkyl; or
 C₂-C₄-alkenyl chain;

or a pharmaceutically acceptable salt thereof.

40 The present invention is also directed to a process for preparation of a purine derivative compound of the formula:

- 11 -



Formula I

wherein:

R_1 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_6 -straight chain alkyl;
 C_2 - C_6 -straight alkenyl chain;
 C_3 - C_6 -branched alkyl chain;
 C_3 - C_6 -branched alkenyl chain;
 C_3 - C_7 -cycloalkyl;
 CH_2 -(C_3 - C_7 -cycloalkyl);
 CH_2CF_3 ;
 $CH_2CH_2CF_3$; and
 $CH(CF_3)_2$;

the combination of X, D, and Q are either:

$D=Q=N$, and $X=CH$; or
 $D=X=N$, and $Q=CH$; or
 $Q=X=N$, and $D=CH$; or
 $Q=N$, and $D=X=CH$;

$V=$ NH;
 O;
 S; or
 CH_2 ;

$R_2=$ phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F, Cl, Br, CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;
 1-naphthyl;
 2-naphthyl;
 heterocycles selected from the group consisting of:
 2-pyridyl;
 3-pyridyl;

- 12 -

- 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 5 thiophene-2-yl;
 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 10 oxazol-4-yl;
 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 15 imidazol-2-yl;
 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 20 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 25 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 30 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 35 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 40 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

- R₃ are the same or different and independently selected from the group consisting of:
 45 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;

- 13 -

$(\text{CH}_2)_n\text{Ph}$; and
 $(\text{CH}_2)_n$ -substituted phenyl, wherein the phenyl substituents are as defined
 above in R_2 ;

5 $\text{R}_4 =$ H;
 C_1 - C_4 -straight chain alkyl; or
 C_3 - C_4 -branched chain alkyl;

R_3 and R_4 can be linked together by a carbon chain to form with intervening atoms a
 10 5-8-membered saturated or unsaturated ring;

$n_1 =$ 0-3;

$n =$ 0-3;

15 $\text{A} =$ CH_2 ;
 $(\text{CH}_2)_2$;
 $(\text{CH}_2)_3$;
 OCH_2CH_2 ; or
 20 CHCH_3 ;

$\text{Y} =$ H;
 OR_1 ;
 $\text{N}(\text{R}_1)_2$;
 25 $\text{N}(\text{R}_1)\text{C}(\text{O})\text{R}_3$;
 $\text{N}(\text{R}_1)\text{C}(\text{O})\text{R}_5$;
 $\text{N}(\text{R}_1)\text{C}(\text{O})\text{CH}(\text{R}_6)\text{NH}_2$;
 $\text{N}(\text{R}_1)\text{SO}_2\text{R}_3$;
 $\text{N}(\text{R}_1)\text{C}(\text{O})\text{NHR}_3$; or
 30 $\text{N}(\text{R}_1)\text{C}(\text{O})\text{OR}_6$;


$\text{R}_5 =$ C_3 - C_7 -cycloalkyl;

$\text{R}_6 =$ C_1 - C_4 -straight chain alkyl;
 35 C_3 - C_4 -branched chain alkyl;
 C_2 - C_4 -alkenyl chain;
 $(\text{CH}_2)_n\text{Ph}$; or
 $(\text{CH}_2)_n$ -substituted phenyl, wherein the phenyl substituents are as defined
 above in R_2 ;

40 or a pharmaceutically acceptable salt thereof, said process comprising:

reacting a first intermediate compound of the formula:

- 14 -

**Formula IX**


where Z= Br or I

5

with a compound of the formula: $R_2-B(OH)_2$, $R_2-Sn(n-Bu)_3$, or $R_2-Sn(Me)_3$, or mixtures thereof, under conditions effective to form the purine derivative compound.

Another aspect of the present invention is directed to a process for preparation of a purine derivative compound of the formula:

10

**Formula I**

15 wherein:

R_1 are the same or different and independently selected from the group consisting of:

H;

C_1 - C_6 -straight chain alkyl;

20 C_2 - C_6 -straight alkenyl chain;

C_3 - C_6 -branched alkyl chain;

C_3 - C_6 -branched alkenyl chain;

C_3 - C_7 -cycloalkyl;

CH_2 -(C_3 - C_7 -cycloalkyl);

25 CH_2CF_3 ;

$CH_2CH_2CF_3$; and

- 16 -

benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 5 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 10 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 15 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:
 20 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 25 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 30 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a
 5-8-membered saturated or unsaturated ring;

35 n₁= 0-3;

n= 0-3;

A= CH₂;
 40 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

45 Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;

- 17 -

5 $N(R_1)C(O)R_5$;
 $N(R_1)C(O)CH(R_6)NH_2$;
 $N(R_1)SO_2R_3$;
 $N(R_1)C(O)NHR_3$; or
 $N(R_1)C(O)OR_6$;


$R_5 =$ C₃-C₇-cycloalkyl;

10 $R_6 =$ C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

15

or a pharmaceutically acceptable salt thereof, said process comprising:

reacting a first intermediate compound of the formula:




20

Formula XV

under reductive or hydrogenation conditions effective to form the purine derivative compound.

25 Another aspect of the present invention is directed to a process for
 preparation of a purine derivative compound of the formula:

- 18 -



Formula I

wherein:

5

R_1 are the same or different and independently selected from the group consisting of:

- H;
 C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 10 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 15 CH₂CH₂CF₃; and
 CH(CF₃)₂;

the combination of X, D, and Q are either:

- D=Q=N, and X=CH; or
 20 D=X=N, and Q=CH; or
 Q=X=N, and D=CH; or
 Q=N, and D=X=CH;

- V= NH;
 25 O;
 S; or
 CH₂;

- 30 R_2 = phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of R_1 ,
 OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F,
 Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;
 1-naphthyl;
 35 2-naphthyl;
 heterocycles selected from the group consisting of:
 2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 40 2-pyrimidyl;

- 19 -

4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;
 5 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 oxazol-5-yl;
 10 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;
 imidazol-4-yl;
 15 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 20 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 25 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 30 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 35 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 40 Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 45 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and

- 20 -

$(\text{CH}_2)_n$ -substituted phenyl, wherein the phenyl substituents are as defined above in R_2 ;

5 $R_4 =$ H;
C₁-C₄-straight chain alkyl; or
C₃-C₄-branched chain alkyl;

R_3 and R_4 can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

10 $n_1 =$ 0-3;

$n =$ 0-3;

15 $A =$ CH₂;
(CH₂)₂;
(CH₂)₃;
OCH₂CH₂; or
CHCH₃;


20 $Y =$ H;
OR₁;
N(R₁)₂;
N(R₁)C(O)R₃;
25 N(R₁)C(O)R₅;
N(R₁)C(O)CH(R₆)NH₂;
N(R₁)SO₂R₃;
N(R₁)C(O)NHR₃; or
N(R₁)C(O)OR₆;

30 $R_5 =$ C₃-C₇-cycloalkyl;

$R_6 =$ C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
35 C₂-C₄-alkenyl chain;
(CH₂)_nPh; or
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R_2 ;


40 or a pharmaceutically acceptable salt thereof, said process comprising:
reacting a first intermediate compound of the formula:

- 21 -

**Formula XIV**

with a compound of the formula:


5

**Formula VIII**

10 where V₁= NH₂;
OH; or
SH;

under conditions effective to form the purine derivative compound.

15 Another aspect of the present invention is directed to a process for preparation of a purine derivative compound of the formula:

**Formula XX**

wherein:

- 22 -

R_1 are the same or different and independently selected from the group consisting of:

- H;
- C_1 - C_6 -straight chain alkyl;
- C_2 - C_6 -straight alkenyl chain;
- 5 C_3 - C_6 -branched alkyl chain;
- C_3 - C_6 -branched alkenyl chain;
- C_3 - C_7 -cycloalkyl;
- CH_2 -(C_3 - C_7 -cycloalkyl);
- CH_2CF_3 ;
- 10 $CH_2CH_2CF_3$; and
- $CH(CF_3)_2$;

the combination of X, D, and Q are either:

- D=Q=N and X=CH; or
- 15 D=X=N and Q=CH; or
- Q=X=N and D=CH; or
- Q=N and D=X=CH;

- V= NH;
- 20 O;
- S; or
- CH_2 ;

- R_2 = phenyl;
- 25 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F, Cl, Br, CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;

- 1-naphthyl;
- 30 2-naphthyl;
- heterocycles selected from the group consisting of:

- 2-pyridyl;
- 3-pyridyl;
- 4-pyridyl;
- 35 2-pyrimidyl;
- 4-pyrimidyl;
- 5-pyrimidyl;
- thiophene-2-yl;
- thiophene-3-yl;
- 40 2-furanyl;
- 3-furanyl;
- oxazol-2-yl;
- oxazol-4-yl;
- oxazol-5-yl;
- 45 thiazol-2-yl;
- thiazol-4-yl;
- thiazol-5-yl;
- imidazol-2-yl;

- 23 -

5 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 10 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 15 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 20 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 25 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:
 30 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 35 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 40 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a
 5-8-membered saturated or unsaturated ring;

45 n₁= 0-3;

 n= 0-3;

- 24 -

- A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 5 CHCH₃;

 Y= NR₁C(O)R₃;
 NR₁C(O)R₅;
 NR₁SO₂R₃;
 10 NR₁C(O)NHR₃; or
 NR₁C(O)OR₆

 R₅= C₃-C₇-cycloalkyl;


 15 R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 20 above in R₂;

or a pharmaceutically acceptable salt thereof;

said process comprising:

- reacting a first intermediate compound having the same formula as the purine
 25 derivative compound except that Y=NHR₁, with R₃COCl or R₅COCl or R₃SO₂Cl or
 R₃NCO or R₆OC(O)Cl under conditions effective to form the purine derivative
 compound.

- Yet another aspect of the present invention is directed to a process for
 30 preparation of a purine derivative compound of the formula:



Formula XIX

wherein:

- 25 -

R_1 are the same or different and independently selected from the group consisting of:

- H;
- C₁-C₆-straight chain alkyl;
- C₂-C₆-straight alkenyl chain;
- 5 C₃-C₆-branched alkyl chain;
- C₃-C₆-branched alkenyl chain;
- C₃-C₇-cycloalkyl;
- CH₂-(C₃-C₇-cycloalkyl);
- CH₂CF₃;
- 10 CH₂CH₂CF₃; and
- CH(CF₃)₂;

the combination of X, D, and Q are either:

- D=Q=N and X=CH; or
- 15 D=X=N and Q=CH; or
- Q=X=N and D=CH; or
- Q=N and D=X=CH;

- V= NH;
- 20 O;
- S; or
- CH₂;

- R₂= phenyl;
- 25 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

- 1-naphthyl;
- 30 2-naphthyl;
- heterocycles selected from the group consisting of:

- 2-pyridyl;
- 3-pyridyl;
- 4-pyridyl;
- 35 2-pyrimidyl;
- 4-pyrimidyl;
- 5-pyrimidyl;
- thiophene-2-yl;
- thiophene-3-yl;
- 40 2-furanyl;
- 3-furanyl;
- oxazol-2-yl;
- oxazol-4-yl;
- oxazol-5-yl;
- 45 thiazol-2-yl;
- thiazol-4-yl;
- thiazol-5-yl;
- imidazol-2-yl;

- 26 -

5 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 10 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 15 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 20 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 25 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:
 30 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 35 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 40 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a
 5-8-membered saturated or unsaturated ring;

45 n₁= 0-3;

n= 0-3;

- 27 -

- 5 A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;
- Y= NHC(O)CH(R₆)NH₂;
- 10 R₅= C₃-C₇-cycloalkyl;
- R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
15 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

or a pharmaceutically acceptable salt thereof;

said process comprising:

- 20 reacting a first intermediate compound having the same formula as the purine
derivative compound except that Y is NH₂, with a compound of the formula:
PNHCH(R₆)CO₂H under conditions effective to form the purine derivative compound
after a suitable deprotection strategy,

wherein

- 25 P= C(O)OtBu;
 C(O)OCH₂Ph;
 Fmoc
 Benzyl; or
 Alloc.

- 30 The compounds of the present invention, as described in Formula I,
show significantly improved growth inhibition of human transformed cell lines and/or
cyclin/cdk inhibition relative to compounds of the prior art. These compounds have
been demonstrated to be potent growth inhibitors in dozens of human transformed cell
lines. Olomoucine, a structurally related purine derivative, is a poor human
35 transformed cell growth inhibition agent with GI₅₀ values in the 20,000-100,000 nM
range over 60-transformed cell lines. By contrast, the compounds of the present
invention demonstrate GI₅₀ values over 60-transformed cell lines in the <10-25,000
nM range, preferably in the <10-100 nM range over 60-transformed cell lines, and,
most preferably, <10 nM across 60-human transformed cell lines. This finding is


- 28 -

unexpected from the prior art, which specifically teaches that compounds of the present invention would not be potent human transformed cell line growth inhibitors.

The R_2 group in Formula I imparts unexpected and significant improvement in growth inhibition in human transformed cell lines, while substitution of various groups at R_3 and R_4 found in Formula I impart important features that contribute to cyclin/cdk inhibition and growth inhibition of human transformed cell lines. Specifically, the combination of the R_2 group and the substitutions within R_3 and R_4 result in compounds with superior biological activity. Compounds which are cyclin/cdk inhibitors and/or human transformed cell line growth inhibitors have utility in treating human proliferative cellular disorders.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention are represented by the chemical structure found in Formula I.




Formula I

wherein:

R_1 are the same or different and independently selected from the group consisting of:

- H;
- C_1 - C_6 -straight chain alkyl;
- C_2 - C_6 -straight alkenyl chain;
- C_3 - C_6 -branched alkyl chain;
- C_3 - C_6 -branched alkenyl chain;
- C_3 - C_7 -cycloalkyl;
- CH_2 -(C_3 - C_7 -cycloalkyl);
- CH_2CF_3 ;



the combination of X, D, and Q are either:

5 D=Q=N, and X=CH; or
 D=X=N, and Q=CH; or
 Q=X=N, and D=CH; or
 Q=N, and D=X=CH;

10 V= NH;
O;
S; or
CH₂;

15 R₂= phenyl;
substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

20 1-naphthyl;
2-naphthyl;
heterocycles selected from the group consisting of:

25 2-pyridyl;
3-pyridyl;
4-pyridyl;
2-pyrimidyl;
4-pyrimidyl;
5-pyrimidyl;
thiophene-2-yl;
thiophene-3-yl;
30 2-furanyl;
3-furanyl;
oxazol-2-yl;
oxazol-4-yl;
35 oxazol-5-yl;
thiazol-2-yl;
thiazol-4-yl;
thiazol-5-yl;
imidazol-2-yl;
40 imidazol-4-yl;
pyrazol-3-yl;
pyrazol-4-yl;
isoxazol-3-yl;
isoxazol-4-yl;
45 isoxazol-5-yl;
isothiazol-3-yl;
isothiazol-4-yl;
isothiazol-5-yl;

- 30 -

1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 5 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 10 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 15 4-isoquinolinyl; or
 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

20 R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 25 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

30 R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a
 5-8-membered saturated or unsaturated ring;

35 n₁= 0-3;

n= 0-3;

40 A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

45 Y= H;
 OR₁;
 N(R₁)₂;

- 31 -

N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 5 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;


R₅= C₃-C₇-cycloalkyl;

10 R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 15 above in R₂;

or a pharmaceutically acceptable salt thereof.

More preferably, the compounds of the current invention are represented by the chemical structure found in Formula III.

20



Formula III

wherein:

25

R₁ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 30 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 35 CH₂CH₂CF₃; and

$\text{CH}(\text{CF}_3)_2$;

the combination of X, D, and Q are either:

- 5 D=Q=N, and X=CH; or
 D=X=N, and Q=CH; or
 Q=X=N, and D=CH; or
 Q=N, and D=X=CH;

- 10 V= NH;
 O;
 S; or
 CH₂;

- 15 R₂= phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any
 position and are independently selected from the group consisting of R₁,
 OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F,
 Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

- 20 1-naphthyl;
 2-naphthyl;
 heterocycles selected from the group consisting of:

- 2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;
 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;

- 33 -

benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 5 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 10 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 15 position and are independently selected from the group consisting of Br,
 Cl, F, R₁, and C(O)CH₃;

n= 0-3;

20 A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

25 Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 30 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;

35 R₃ are the same or different and independently selected from the group consisting of:
 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 40 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

45 R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl; or

C₂-C₄-alkenyl chain;

or a pharmaceutically acceptable salt thereof.

5 In another embodiment, the present invention is directed to a method of treating a mammal with a disorder mediated by elevated levels of cellular proliferation comprising administering a therapeutically effective amount of the compound of the present invention to the mammal under conditions effective to treat the disorder mediated by elevated levels of cell proliferation.

10 The compounds of the present invention can be administered orally, parenterally, for example, subcutaneously, intravenously, intramuscularly, intraperitoneally, by intranasal instillation, or by application to mucous membranes, such as, that of the nose, throat, and bronchial tubes. They may be administered alone or with suitable pharmaceutical carriers, and can be in solid or liquid form such as,
15 tablets, capsules, powders, solutions, suspensions, or emulsions.

 Based on the results obtained in the standard pharmacological test procedures described below, the compounds of the present invention are useful as antineoplastic agents. More particularly, the compounds of the present invention are useful for inhibiting the growth of neoplastic cells, causing cell death of neoplastic
20 cells, and eradicating neoplastic cells. The compounds of the present invention are, therefore, useful for treating solid tumors, including sarcomas and carcinomas, such as astrocytomas, prostate cancer, breast cancer, small cell lung cancer, and ovarian cancer, leukemias, lymphomas, adult T-cell leukemia/lymphoma, and other neoplastic disease states.

25 In addition to the utilities described above, many of the compounds of the present invention are useful in the preparation of other compounds.

 The active compounds of the present invention may be orally administered, for example, with an inert diluent, or with an assimilable edible carrier, or they may be enclosed in hard or soft shell capsules, or they may be compressed
30 into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, these active compounds may be incorporated with excipients and used in the form of tablets, capsules, elixirs, suspensions, syrups, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compound in these compositions may, of course,

be varied and may conveniently be between about 2% to about 60% of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions according to the present invention are prepared so that an oral dosage unit contains between about
5 1 and 250 mg of active compound.

The tablets, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch, or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose,
10 lactose, or saccharin. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar, or both. A syrup may contain, in addition to active ingredient, sucrose as a
15 sweetening agent, methyl and propylparabens as preservatives, a dye, and flavoring such as cherry or orange flavor.

These active compounds may also be administered parenterally. Solutions or suspensions of these active compounds can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be
20 prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Illustrative oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, or mineral oil. In general, water, saline, aqueous dextrose and related sugar solution, and glycols such as, propylene glycol or polyethylene glycol, are preferred liquid carriers, particularly for injectable solutions.
25 Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must
30 be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier

can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

The compounds of the present invention may also be administered
5 directly to the airways in the form of an aerosol. For use as aerosols, the compounds of the present invention in solution or suspension may be packaged in a pressurized aerosol container together with suitable propellants, for example, hydrocarbon propellants like propane, butane, or isobutane with conventional adjuvants. The materials of the present invention also may be administered in a non-pressurized form
10 such as in a nebulizer or atomizer.

General Synthetic Schemes

The compounds of the present invention can be prepared by conventional methods of organic synthesis practiced by those skilled in the art. The
15 general reaction sequences outlined below are general methods useful for preparing the compounds of the present invention and are not meant to be limiting in scope or utility.

Reaction of 2,6-dichloropurine (**Formula IV**) with various amines of **Formula V**, many of which are commercially available or prepared by literature
20 methods or modifications of literature methods, in the presence of a polar solvent, such as ethanol, provides purines of **Formula VI** (General Flowsheet I, *infra*).
Reaction of purines of **Formula VI** with alkyl halides (R_1-Z) in the presence of a base such as potassium carbonate provides N1-alkylated purines of **Formula VII**.
Chloride displacement of N1-alkylated purines of **Formula VII** with amines, thiols or
25 alcohols of structure **Formula VIII**, either in neat solution or in an inert solvent such as ethanol or butanol, with or without a base such as sodium hydride as appropriate, at an appropriate temperature provides purines of **Formula IX** ($V=NH, O, S$).
Transition metal-mediated cross-coupling reaction of purines of **Formula IX** with boronic acid ($R_2-B(OH)_2$) or tin reagents ($R_2-Sn(n-Bu)_3$ or R_2-SnMe_3) provides
30 purines of **Formula X** ($V=NH, O, S$). If in **Formula X** ($Y=NH_2$), then subsequent reaction of **Formula X** ($Y=NH_2$) with acid chloride (R_3COC1), or sulfonyl chloride (R_3SO_2C1), or isocyanate (R_3NCO), or chloroformate ($C1C(O)OR_6$) reagents

- 37 -

provides purines of Formula XI wherein $Y = \text{NHC(O)R}_3$, NHSO_2R_3 , or NHC(O)NHR_3 , or NHC(O)OR_6 , respectively. On the other hand, if in Formula X, Y already is OR_1 or NHC(O)R_3 or NHSO_2R_3 or NHC(O)NHR_3 or NHC(O)OR_6 , as a result of what Y started out as in Formula VIII, then this last step is unnecessary.

5 Reaction of purines of Formula VII, with alkenyl tin reagents of Formula XII, which are prepared by conventional methods described in the literature, in the presence of a transition metal catalyst, such as Pd(0), provides purines of Formula XIII (General Flowsheet II, *infra*). Subsequent reaction of purines of Formula XIII with boronic acid ($\text{R}_2\text{-B(OH)}_2$) or tin reagents ($\text{R}_2\text{-Sn(n-Bu)}_3$ or $\text{R}_2\text{-SnMe}_3$) in the presence of a transition metal catalyst, such as Pd(0),
10 provides purines of Formula XV. Alternatively, by switching the order of reactions dependent on the precise reactivity of the purine of Formula VII, reaction of purines of Formula VII with boronic acid ($\text{R}_2\text{-B(OH)}_2$) or tin reagents ($\text{R}_2\text{-Sn(n-Bu)}_3$ or $\text{R}_2\text{-SnMe}_3$) in the presence of a transition metal catalyst, such as Pd(0), provides purines
15 of Formula XIV. Subsequent reaction of purines of Formula XIV, with alkenyl tin reagents of Formula XII, which are prepared by conventional methods described in the literature, in the presence of a transition metal catalyst, such as Pd(0), provides purines of Formula XV. Finally reduction of the olefin within Formula XV provides purines of Formula X ($\text{V}=\text{CH}_2$).

20

Definitions of the groups include:

Z= Br;
 I;

25 V₁= NH₂;
 OH;
 SH;

R₁ are the same or different and independently selected from:

30 H;
 C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
35 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 CH₂CH₂CF₃;

- 38 -



the combination of X, D, and Q are selected from:

- 5 D=Q=N, and X=CH;
 D=X=N, and Q=CH;
 Q=X=N, and D=CH;
 Q=N, and D=X=CH;

- 10 V= NH;
 O;
 S;
 CH₂;

R₂ can be in any position on the ring and selected from:

- 15 phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any
 position and are independently selected from R₁, OR₁, SR₁, S(O)R₁,
 S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁,
 C(O)NHR₁, phenyl, C(O)NHCHR₁CH₂OH;
 20 1-naphthyl;
 2-naphthyl;
 heterocycles including:
 2-pyridyl;
 3-pyridyl;
 25 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 30 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 35 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;
 40 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 45 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;

- 39 -

1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 5 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 10 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl;
 15 4-isoquinolinyl;
 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from Br, Cl, F, R₁, C(O)CH₃;

R₃ are the same or different and independently selected from:
 20 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh;
 25 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl;
 30 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form a 5-8-membered saturated
 or unsaturated ring;

35 n₁= 0-3;

n= 0-3;

40 A= (CH₂);
 (CH₂)₂;
 (CH₂)₃;
 (OCH₂CH₂);
 (CHCH₃);

45 Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;

- 40 -


5 $N(R_1)C(O)R_5$;
 $N(R_1)C(O)CH(R_6)NH_2$;
 $N(R_1)SO_2R_3$;
 $N(R_1)C(O)NHR_3$;
 $N(R_1)C(O)OR_6$;

$R_5 =$ C₃-C₇-cycloalkyl;


10 $R_6 =$ C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 $(CH_2)_nPh$;
 $(CH_2)_n$ -substituted phenyl, wherein the phenyl substituents are as defined
15 above in R_2 .

- 41 -

General Flowsheet I




General Flowsheet II



Additional, general non-limiting syntheses of compounds of the present invention of

5 **Formula X** and **Formula XI** are shown below in General Flowsheet III.

General Flowsheet III



Reaction of various amines of **Formula V**, many of which are commercially available
 5 or prepared by literature methods or modifications of literature methods, with boronic
 acid ($\text{R}_2-\text{B}(\text{OH})_2$) or tin reagents ($\text{R}_2-\text{Sn}(\text{n-Bu})_3$) or (R_2-SnMe_3) in the presence of a
 transition metal catalyst, such as Pd(0), provides biaryl amines of **Formula XVII**.
 Reaction of 2,6-dichloropurine (**Formula IV**) with various amines of **Formula XVII**,
 in the presence of a polar solvent, such as ethanol, provides purines of **Formula**
 10 **XVIII**. Reaction of purines of **Formula XVIII** with alkyl halides (R_1-Z) in the
 presence of a base such as potassium carbonate provides N1-alkylated purines of

- 44 -

Formula XIV. Chloride displacement of N1-alkylated purines of **Formula XIV** with amines, thiols or alcohols of **Formula VIII**, either in neat solution or in an inert solvent such as ethanol or butanol, with or without a base such as sodium hydride as appropriate, at an appropriate temperature provides purines of **Formula X** ($V=NH$, O, S). If in **Formula X** ($Y=NH_2$), then subsequent reaction of **Formula X** ($Y=NH_2$) with acid chloride (R_3COCl), or sulfonyl chloride (R_3SO_2Cl), or isocyanate (R_3NCO), or chloroformate ($ClC(O)OR_6$) reagents provides purines of **Formula XI** wherein $Y=NHC(O)R_3$, $NHSO_2R_3$, or $NHC(O)NHR_3$, or $NHC(O)OR_6$, respectively. On the other hand, if in **Formula X**, Y already is OR_1 or $NHC(O)R_3$ or $NHSO_2R_3$ or $NHC(O)NHR_3$ or $NHC(O)OR_6$, as a result of what Y started out as in **Formula VIII**, then this last step is unnecessary.

Definitions of the groups include:

15 $Z=$ Br;
I;

$V_1=$ NH_2 ;
OH;
SH;

20 R_1 are the same or different and independently selected from:

H;
C₁-C₆-straight chain alkyl;
C₂-C₆-straight alkenyl chain;
25 C₃-C₆-branched alkyl chain;
C₃-C₆-branched alkenyl chain;
C₃-C₇-cycloalkyl;
CH₂-(C₃-C₇-cycloalkyl);
CH₂CF₃;
30 CH₂CH₂CF₃;
CH(CF₃)₂;

the combination of X, D, and Q are selected from:

35 D=Q=N, and X=CH;
D=X=N, and Q=CH;
Q=X=N, and D=CH;
Q=N, and D=X=CH;

40 $V=$ NH;
O;
S;
CH₂;

R₂ can be in any position on the ring and selected from:

- phenyl;
substituted phenyl, wherein the substituents (1-2 in number) are in any
5 position and are independently selected from R₁, OR₁, SR₁, S(O)R₁,
S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁,
C(O)NHR₁, phenyl, C(O)NHCHR₁CH₂OH;
1-naphthyl;
2-naphthyl;
10 heterocycles including:
2-pyridyl;
3-pyridyl;
4-pyridyl;
2-pyrimidyl;
15 4-pyrimidyl;
5-pyrimidyl;
thiophene-2-yl;
thiophene-3-yl;
2-furanyl;
20 3-furanyl;
oxazol-2-yl;
oxazol-4-yl;
oxazol-5-yl;
thiazol-2-yl;
25 thiazol-4-yl;
thiazol-5-yl;
imidazol-2-yl;
imidazol-4-yl;
pyrazol-3-yl;
30 pyrazol-4-yl;
isoxazol-3-yl;
isoxazol-4-yl;
isoxazol-5-yl;
isothiazol-3-yl;
35 isothiazol-4-yl;
isothiazol-5-yl;
1,3,4-thiadiazol-2-yl;
benzo[b]furan-2-yl;
benzo[b]thiophene-2-yl;
40 2-pyrrolyl;
3-pyrrolyl;
1,3,5-triazin-2-yl;
pyrazin-2-yl;
pyridazin-3-yl;
45 pyridazin-4-yl;
2-quinolinyl;
3-quinolinyl;
4-quinolinyl;

- 46 -

1-isoquinolinyl;
 3-isoquinolinyl;
 4-isoquinolinyl;

5 substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from Br, Cl, F, R₁, C(O)CH₃;

R₃ are the same or different and independently selected from:

H;
 C₁-C₄-straight chain alkyl;
 10 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh;
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

15 R₄= H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;

20 R₃ and R₄ can be linked together by a carbon chain to form a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

25 n= 0-3;

A= (CH₂);
 (CH₂)₂;
 (CH₂)₃;
 30 (OCH₂CH₂);
 (CHCH₃);

Y= H;
 OR₁;
 35 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 40 N(R₁)C(O)NHR₃;
 N(R₁)C(O)OR₆;


R₅= C₃-C₇-cycloalkyl;

45 R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh;

(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂.

Additional, general non-limiting syntheses of compounds of the present invention of
 5 **Formula XVI, Formula XVII and Formula XVIII** are shown below in General Flowsheet IV.

General Flowsheet IV



10 If in **Formula X** (Y=NH₂), then subsequent reaction of **Formula X** (Y=NH₂) with alkyl halide (R₅CH₂Z), an appropriate base, and a solvent; or reaction of **Formula X** (Y=NH₂) with aldehyde (R₅CHO) in the presence of a solvent and a suitable reducing agent provides purines of **Formula XVI** wherein Y=NHR₁, or N(R₁)₂. On the other hand, if in **Formula X**, Y already is NHR₁, or N(R₁)₂, as a result
 15 of what Y started out as in **Formula X**, then this last step is unnecessary. If in **Formula XVI** (Y=NHR₁), then subsequent reaction of **Formula XVI** (Y=NHR₁)

- 48 -

with acid chloride ($R_3\text{COCl}$), or sulfonyl chloride ($R_3\text{SO}_2\text{Cl}$), or isocyanate ($R_3\text{NCO}$), or chloroformate (ClC(O)OR_6) reagents provides purines of **Formula XX** wherein $Y=\text{NR}_1\text{C(O)R}_3$, or $\text{NR}_1\text{C(O)R}_5$, or $\text{NR}_1\text{SO}_2\text{R}_3$, or $\text{NR}_1\text{C(O)NHR}_3$, or $\text{NR}_1\text{C(O)OR}_6$, respectively. On the other hand, if in **Formula XVI**, Y already is $\text{NR}_1\text{C(O)R}_3$, or

5 $\text{NR}_1\text{C(O)R}_5$, or $\text{NR}_1\text{SO}_2\text{R}_3$, or $\text{NR}_1\text{C(O)NHR}_3$, or $\text{NR}_1\text{C(O)OR}_6$, as a result of what started out as in **Formula XVI**, then this last step is unnecessary.

If in **Formula X** ($Y=\text{NH}_2$), then subsequent reaction of **Formula X** ($Y=\text{NH}_2$) with acid ($\text{PNHCH(R}_6\text{)CO}_2\text{H}$), in a suitable solvent in the presence of an appropriate

10 coupling agent provides a purine derivative; which upon suitable deprotection provides purines of **Formula XIX** wherein $Y=\text{NHC(O)CH(R}_6\text{)NH}_2$. On the other hand, if in **Formula X**, Y already is $\text{NHC(O)CH(R}_6\text{)NH}_2$, as a result of what Y started out as in **Formula X**, then this last step is unnecessary.

15 Definitions of the groups include:

Z= Br;
I;

20 P= C(O)OtBu;
C(O)OCH₂Ph;
Fmoc;
Benzyl;
Alloc;

25 R₁ are the same or different and independently selected from:

H;
C₁-C₆-straight chain alkyl;
C₂-C₆-straight alkenyl chain;
C₃-C₆-branched alkyl chain;
30 C₃-C₆-branched alkenyl chain;
C₃-C₇-cycloalkyl;
CH₂-(C₃-C₇-cycloalkyl);
CH₂CF₃;
CH₂CH₂CF₃;
35 CH(CF₃)₂;

the combination of X, D, and Q are selected from:

D=Q=N, and X=CH;
D=X=N, and Q=CH;

- 49 -

 $Q=X=N$, and $D=CH$; $Q=N$, and $D=X=CH$;

5 $V=$ NH ;
 O ;
 S ;
 CH_2 ;

R_2 can be in any position on the ring and selected from:

10 phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any
 position and are independently selected from R_1 , OR_1 , SR_1 , $S(O)R_1$,
 $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F , Cl , Br , CF_3 , $C(O)R_1$,
 $C(O)NHR_1$, phenyl, $C(O)NHCHR_1CH_2OH$;

15 1-naphthyl;
 2-naphthyl;
 heterocycles including:

20 2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;

25 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;

30 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;

35 imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;

40 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;

45 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;

- 50 -

1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 5 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl;
 10 4-isoquinolinyl;
 substituted heterocycle, wherein the substituents (1-2 in number) are in any
 position and are independently selected from Br, Cl, F, R₁, C(O)CH₃;

R₃ are the same or different and independently selected from:

15 H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh;
 20 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl;
 25 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form a 5-8-membered saturated or unsaturated ring;

30 n₁= 0-3;

n= 0-3;

35 A= (CH₂);
 (CH₂)₂;
 (CH₂)₃;
 (OCH₂CH₂);
 (CHCH₃);

40 Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 45 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃;
 N(R₁)C(O)OR₆;

- 51 -

R₅= C₃-C₇-cycloalkyl;


5 R₆= C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh;
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
above in R₂;

10 R₈= C₁-C₅-straight chain alkyl;
C₂-C₅-straight alkenyl chain;
C₃-C₅-branched alkyl chain;
C₃-C₅-branched alkenyl chain;
15 C₃-C₇-cycloalkyl;
CF₃;
CH₂CF₃.

- 52 -


The synthesis of compound 5 is shown below in Scheme I.

Scheme I




The synthesis of compound 11 is shown below in Scheme II.

Scheme II




The syntheses of compounds 12, 13 and 14 are shown below in Scheme III.

Scheme III




The synthesis of compound 17 is shown below in Scheme IV.

Scheme IV




The synthesis of compound 17 is shown below in Scheme V.

Scheme V




The synthesis of compound **25** is shown below in Scheme VI.

Scheme VI




An alternative synthesis of compound 25 is shown below in Scheme VII.

Scheme VII



- 59 -


The synthesis of compound 32 is shown below in Scheme VIII.

Scheme VIII

- 60 -


The syntheses of compounds 33 and 34 are shown below in Scheme IX.

Scheme IX




The syntheses of compounds 36, 38, and 40 are shown below in Scheme X.

Scheme X




The synthesis of compound 43 is shown below in Scheme XI.

Scheme XI




- 63 -

The synthesis of compound 46 is shown below in Scheme XII.

Scheme XII


The syntheses of compound **48** and **50** are shown below in Scheme XIII.

Scheme XIII




The synthesis of compound **53** is shown below in Scheme XIV.

Scheme XIV




The synthesis of compound 54 is shown below in Scheme XV.

Scheme XV




The synthesis of compound 56 is shown below in Scheme XVI.

Scheme XVI




The synthesis of compound **58** is shown below in Scheme XVII.

Scheme XVII




The synthesis of compound **60** is shown below in Scheme XVIII.

Scheme XVIII




The syntheses of compounds **61**, and **62** are shown below in Scheme XIX.

Scheme XIX





The syntheses of compounds 64, and 65 are shown below in Scheme XX.

Scheme XX



The syntheses of compounds 66, and 67 are shown below in Scheme XXI.


Scheme XXI



- 73 -

The synthesis of compound 73 is shown below in Scheme XXII.


Scheme XXII



- 74 -


The syntheses of compounds 74, 75, and 76 are shown below in Scheme XXIII.

Scheme XXIII



The synthesis of compound **77** is shown below in Scheme XXIV.


Scheme XXIV



5

The synthesis of compound **78** is shown below in Scheme XXV.


Scheme XXV



- 76 -


An alternative synthesis of compound 78, and the synthesis of compound 79 are shown below in Scheme XXVI.

Scheme XXVI




The synthesis of compound **80** is shown below in Scheme XXVII.

Scheme XXVII




The syntheses of compounds 86, and 87 are shown below in Scheme XXVIII.

Scheme XXVIII




- 79 -

The synthesis of compound **88** is shown below in Scheme XXIX.

Scheme XXIX


The syntheses of compounds **93**, and **94** are shown below in Scheme XXX.

Scheme XXX




The syntheses of compounds **95**, and **96** are shown below in Scheme XXXI.

Scheme XXXI



The synthesis of compound **97** is shown below in Scheme XXXII.


Scheme XXXII



- 83 -


The syntheses of compounds **98**, and **99** are shown below in Scheme XXXIII.

Scheme XXXIII




The synthesis of compound **100** is shown below in Scheme XXXIV.

Scheme XXXIV




The syntheses of compounds **101**, and **102** are shown below in Scheme XXXV.

Scheme XXXV



The syntheses of compounds **103**, and **104** are shown below in Scheme XXXVI.


Scheme XXXVI



- 87 -


The syntheses of compounds 106, 107, and 108 are shown below in Scheme XXXVII.

Scheme XXXVII




The syntheses of compounds 109, and 110 are shown below in Scheme XXXVIII.

Scheme XXXVIII




The syntheses of compounds 111, and 112 are shown below in Scheme XXXIX.

Scheme XXXIX




- 90 -

The synthesis of compound **113** is shown below in Scheme XL.

Scheme XL


- 91 -

The syntheses of compounds 114, 115, 116, and 117 are shown below in Scheme XLI.




The synthesis of compound 118 is shown below in Scheme XLII.

Scheme XLII




The syntheses of compounds 123 and 124 are shown below in Scheme XLIII.

Scheme XLIII




The syntheses of compounds 131 and 132 are shown below in Scheme XLIV.



5 The syntheses of compounds 134 and 135 are shown below in Scheme XLV.

- 95 -


Scheme XLV



The synthesis of compound 137 is shown below in Scheme XLVI.


- 96 -

Scheme XLVI



The syntheses of compounds 139 and 140 are shown below in Scheme XLVII.

Scheme XLVII




5

The synthesis of compound 142 is shown below in Scheme XLVIII.


- 97 -

Scheme XLVIII



The synthesis of compound 144 is shown below in Scheme XLIX.

Scheme XLIX




5

The synthesis of compound 146 is shown below in Scheme L.


- 98 -

Scheme L



The synthesis of compound 148 is shown below in Scheme LI.

Scheme LI




5

The syntheses of compounds 149-152 are shown below in Scheme LII.

- 99 -


Scheme LII



The syntheses of compounds 153-156 are shown below in Scheme LIII.

- 100 -


Scheme LIII



The syntheses of compounds 157-159 are shown below in Scheme LIV.

- 101 -


Scheme LIV



The syntheses of compounds 160-163 are shown below in Scheme LV.

- 102 -


Scheme LV



The syntheses of compounds 164-166 are shown below in Scheme LVI.

- 103 -


Scheme LVI



The syntheses of compounds 167-168 are shown below in Scheme LVII.


- 104 -

Scheme LVII



The syntheses of compounds 169-171 are shown below in Scheme LVIII.


Scheme LVIII



The syntheses of compounds 172-173 are shown below in Scheme LIX.

- 106 -


Scheme LIX



The syntheses of compounds 174-176 are shown below in Scheme LX.

- 107 -


Scheme LX



The syntheses of compounds 177-178 are shown below in Scheme LXI.


- 108 -

Scheme LXI



The syntheses of compounds 179-180 are shown below in Scheme LXII.


Scheme LXII



5

The syntheses of compounds 181-182 are shown below in Scheme LXIII.


Scheme LXIII



The syntheses of compounds 187-188 are shown below in Scheme LXIV.


- 110 -

Scheme LXIV




The syntheses of compounds 193 and 194 are shown below in Scheme LXV.

Scheme LXV




The syntheses of compounds 199-200 are shown below in Scheme LXVI.

Scheme LXVI




The syntheses of compounds 205-206 are shown below in Scheme LXVII.

Scheme LXVII




The syntheses of compounds 207-210 are shown below in Scheme LXVIII.



5 The syntheses of compounds 211-212 are shown below in Scheme LXIX.


Scheme LXIX



The syntheses of compounds 213-215 are shown below in Scheme LXX.


- 116 -

Scheme LXX




The syntheses of compounds 216-217 are shown below in Scheme LXXI.

Scheme LXXI



The syntheses of compounds 218-219 are shown below in Scheme LXXII.


Scheme LXXII



The synthesis of compounds 221 is shown below in Scheme LXXIII.

5


Scheme LXXIII



The synthesis of compound 222 is shown below in Scheme LXXIV.


- 118 -

Scheme LXXIV



The synthesis of compound 223 is shown below in Scheme LXXV.

Scheme LXXV




5

The synthesis of compound 224 is shown below in Scheme LXXVI.


- 119 -

Scheme LXXVI




The synthesis of compound 229 is shown below in Scheme LXXVII.

Scheme LXXVII




The syntheses of compounds 230-233 are shown below in Scheme LXXVIII.

Scheme LXXVIII




The syntheses of compounds 239-241 are shown below in Scheme LXXIX.

Scheme LXXIX




The syntheses of compounds 242-243 are shown below in Scheme LXXX.

Scheme LXXX




The syntheses of compounds 248-250 are shown below in Scheme LXXXI.

Scheme LXXXI



The syntheses of compounds 123 and 124 are shown below in Scheme LXXXII.


Scheme LXXXII



The syntheses of compounds 258-260 are shown below in Scheme LXXXIII.


- 125 -

Scheme LXXXIII



The syntheses of compounds 261-263 are shown below in Scheme LXXXIV.


Scheme LXXXIV



The syntheses of compounds **264-265** are shown below in Scheme LXXXV.


- 127 -

Scheme LXXXV



The synthesis of compound 266 is shown below in Scheme LXXXVI.

Scheme LXXXVI



5

EXAMPLES

10 Proton NMR spectra were obtained on a Bruker AC 300 spectrometer at 300 MHz or a Bruker 500 MHz spectrometer and were referenced to tetramethylsilane as an internal standard. The IR spectrometer used was a single beam Perkin-Elmer Spectrum 1000 FT-IR. All IR spectra obtained were prepared in a

pressed disc of KBr. All IR spectra obtained were acquired with a total of 4 accumulations at a resolution of 4.00 cm^{-1} . Melting points were obtained on a Mel-Temp II apparatus and are uncorrected. Mass spectra were obtained on either a Shimadzu QP-5000 or a PE Sciex API 150 Mass Spectrometer.

5

Example 1 - Preparation of Compound 2

To the starting material 1 (1.0 g, 5.29 mmol) was added 4-bromobenzylamine (2.53 g, 11.4 mmol), and EtOH (11 mL). The mixture was stirred and heated at $50\text{ }^{\circ}\text{C}$ in a round-bottomed flask and then H_2O (1 mL) and EtOH (10 mL) were added to dissolve the solids. The mixture was refluxed for 1 h. Hünig's base (3.68 mL, 21.2 mmol) was added and refluxed overnight, during which time a precipitate formed. The solution was filtered to provide a light yellow solid. The solid was dried *in vacuo* (1.08 g, 60%): ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.75 (bs, 1 H), 8.15 (s, 1 H), 7.52 (d, 2 H), 7.30 (d, 2 H), 4.63 (bs, 2 H); CI MS m/z = 340 $[\text{C}_{12}\text{H}_9\text{BrClN}_5+\text{H}]^+$.

15

Example 2 - Preparation of Compound 3

To the starting material 2 (1.08 g, 3.19 mmol) was added DMSO (11 mL), K_2CO_3 (2.20 g, 15.95 mmol), and 2-iodopropane (1 mL, 9.57 mmol). The solution was stirred overnight then poured into H_2O (75 mL) and stirred. Additional H_2O (25-50 mL) was added to the mixture to form a yellow solid. The stirring was continued at $0\text{ }^{\circ}\text{C}$. The solid was filtered *in vacuo*. The crude product was purified by silica gel chromatography to provide 3 (0.66 g, 50%) as a white solid: mp $136\text{-}140\text{ }^{\circ}\text{C}$; ^1H NMR (300 MHz, CDCl_3) δ 7.78 (s, 1 H), 7.49 (d, 2 H), 7.28 (d, 2 H), 6.12 (bs, 1 H), 4.90-4.70 (m, 3 H), 1.61 (d, 6 H).

20

25

Example 3 - Preparation of Compound 4

30

To starting material 3 (1.44 g, 3.78 mmol) was added 2-amino-1-butanol (5.06 g, 56.7 mmol) and ethanol (5 mL) and the mixture was heated in a sealed tube in an oil bath at $150\text{-}160\text{ }^{\circ}\text{C}$ for 48 h. The cooled solution was transferred to a round-bottomed flask and the ethanol was removed *in vacuo*. The crude product

was purified by flash column chromatography on silica gel to give 4 (0.90 g, 55%):

^1H NMR (300 MHz, CDCl_3) δ 7.44-7.41 (m, 3 H), 7.23 (d, 2 H), 6.22 (s, 1 H), 5.06 (s, 1 H), 4.90 (d, 1 H), 4.78-4.68 (m, 2 H), 4.65-4.55 (m, 1 H), 3.91-3.80 (m, 2 H), 3.66-3.60 (m, 1 H), 1.66-1.47 (m, 8 H), 1.04-0.99 (t, 3 H).

5

Example 4 - Preparation of Compound 5

To starting material 4 (0.13 g, 0.29 mmol) was added 3-acetamidophenylboronic acid (0.21 g, 1.19 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (0.08 g, 0.07 mmol),
10 Na_2CO_3 (2M, 0.60 mL), and toluene (5 mL). The solution was degassed with argon for 10 min then heated at 130 °C for 6 h. The cooled solution was diluted with water and then extracted with CH_2Cl_2 (3 x 50 mL). The combined organic phases were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated to yield a viscous orange oil. The oil was purified by flash column chromatography on silica
15 gel and then the product crystallized upon standing to give 5 (0.06 g, 41%) as a pale yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 8.01-7.21 (m, 9 H), 6.48 (s, 1 H), 4.97 (d, 1 H), 4.82-4.70 (m, 2 H), 4.65-4.53 (m, 1 H), 3.98-3.25 (m, 2 H), 3.20-3.05 (m, 1 H), 2.20 (s, 3 H), 1.69-1.45 (m, 8 H), 1.07-0.98 (t, 3 H).

Example 5 - Preparation of Compound 7

To 4-iodobenzoic acid (52.2 g, 0.21 mol) was added CH_2Cl_2 (500 mL) and DMF (2 drops) at room temperature. Oxalyl chloride (32 g, 0.25 mol) was added dropwise in 0.5 h and stirred for 2 d. The volatiles were removed *in vacuo* to a
25 volume of 150 mL to give the acid chloride and CH_2Cl_2 . To a mixture of ice (500 mL) and NH_4OH (29%; 100 mL) was added the CH_2Cl_2 solution during 15 min. The resulting solids were collected, washed with CH_2Cl_2 , and dried *in vacuo*. The solids were slurried in H_2O for 1 h. The solids were collected by filtration, washed in water and acetone, and dried *in vacuo* to give 7 (48 g; 92%): mp 213-216 °C.

30

Example 6 - Preparation of Compound 8

To a suspension of 7 (11 g, 45 mmol) in THF (50 mL) was added $\text{BH}_3\text{-THF}$ (1M, 22.5 mL, 22.5 mmol). The resulting solution was heated under reflux

- 130 -

overnight. The reaction was cooled in an ice bath and MeOH-HCl (60 mL) was slowly added dropwise. The resulting precipitate was filtered and dried to give 8 (10.8 g, 88%) as a white solid: mp 256-262 °C dec.; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.55 (bs, 3 H), 7.79 (d, 2 H), 7.32 (d, 2 H), 3.98 (s, 2 H).

5

Example 7 - Preparation of Compound 9

To compound 1 (7.63 g, 40.4 mmol) was added compound 8 (10.8 g, 40.4 mmol), water (123 mL), and Hünig's base (14 mL, 81 mmol). The mixture was heated to reflux for 5 h and stirred overnight at room temperature to give a pale yellow solution. An additional quantity of water (150 mL) was added, refluxed for 3 h, then cooled overnight. A pale yellow solid was formed which was filtered, washed with water, rinsed with EtOH (2 x), and dried *in vacuo* to give yield 9 (13.3 g, 80%): ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.68 (bs, 1 H), 8.28 (s, 1 H), 7.68 (d, 2 H), 7.50 (d, 2 H), 5.08 (bs, 1 H), 4.50 (d, 2 H).

15

Example 8 - Preparation of Compound 10

To compound 9 (12.2 g, 31.7 mmol) was added K₂CO₃ (35 g, 0.25 mol), 2-iodopropane (13 g, 0.13 mol) and DMSO (210 mL). The reaction mixture was stirred under N₂ at room temperature overnight, then poured into H₂O (1.5 L) and stirred for 2 d. The precipitate was collected as an off-white solid and washed with Et₂O. The aqueous layer was extracted with EtOAc (2 x) and the combined organic phases were washed with brine, dried over Na₂SO₄, filtered, and evaporated to give an off-white foam (6.4 g). This off-white foam was combined with the precipitate and washed with Et₂O to give 10 (11.0 g): ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.91 (m, 1 H), 8.38 (s, 1 H), 7.74 (d, 2 H), 7.21 (d, 2 H), 5.11 (bs, 1 H), 4.68 (m, 1 H), 4.60 (d, 2 H), 1.48 (d, 6 H).

25

Example 9 - Preparation of Compound 11

Compound 10 (1.52 g, 3.55 mmol), *trans*-1,4-diaminocyclohexane (6.35 g, 55.60 mmol), and EtOH (18 mL) were placed in a sealed tube. The reaction mixture was heated at 120-190 °C for 24 h. The reaction was then allowed to cool to

30

room temperature. The reaction mixture was filtered and the filtrate evaporated. The residue was purified by column chromatography, and dried *in vacuo* for 16 h to yield 11 (1.60 g, 89%) as a yellow sticky oil: ^1H NMR (300 MHz, CDCl_3) δ 7.62 (d, 2 H), 7.44 (s, 1 H), 7.08 (d, 2 H), 6.14 (br, 1 H), 4.75-4.63 (m, 2 H), 4.63-4.54 (m, 2 H), 3.75-3.63 (m, 1 H), 2.72-2.57 (m, 2 H), 2.18-2.00 (m, 2 H), 2.00-1.75 (m, 4 H), 1.54 (d, 6 H), 1.39-1.00 (m, 3 H); API MS m/z = 506 $[\text{C}_{21}\text{H}_{28}\text{IN}_7+\text{H}]^+$.

Example 10 - Preparation of Compound 12

To compound 11 (0.133 g, 0.26 mmol) was added DME (2.5 mL) and 3-thiopheneboronic acid (0.12 g, 0.97 mmol) in a round-bottomed flask and equipped with a condenser purged with argon. To this was added DME (3 mL) followed by tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol) and PPh_3 (0.04 g, 0.15 mmol). Na_2CO_3 (2M, 0.6 mL) and DME (1 mL) was added to the reaction mixture and the reaction mixture was allowed to reflux for 18.5 h, then stirred at room temperature under argon for 46 h. The reaction mixture was diluted with H_2O and extracted with CH_2Cl_2 . The combined organic phases were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography to yield 12 (0.050 g, 41%) as a tan solid: ^1H NMR (300 MHz, CDCl_3) δ 7.56-7.50 (m, 4 H), 7.44-7.35 (m, 3 H), 6.02 (br, 1 H), 4.78 (d, 2 H), 4.69-4.54 (m, 2 H), 3.75 (br, 1 H), 2.69 (br, 1 H), 2.15 (br, 2 H), 1.88 (br, 3 H), 1.54 (d, 7 H), 1.33-0.97 (m, 4 H); API MS m/z = 462 $[\text{C}_{25}\text{H}_{31}\text{N}_7\text{S}+\text{H}]^+$.

Example 11 - Preparation of Compound 13

DME (3 mL), tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol), and PPh_3 (0.04 g, 0.15 mmol) were placed in a round-bottomed flask fitted with a condenser and maintained under argon. Compound 11 (0.13 g, 0.26 mmol), and 4-methylbenzeneboronic acid (0.13 g, 0.98 mmol) dissolved in Na_2CO_3 (2M, 0.6 mL) and DME (1 mL) were added to the reaction mixture. The reaction mixture was refluxed for 19.5 h and stirred at room temperature for 4 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2 . The combined organic phases were washed with brine, dried over anhydrous Na_2SO_4 , and evaporated. The crude product was purified by column chromatography and dried *in vacuo* for 22 h to yield the

desired product 13 (54 mg, 44%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.56-7.41 (m, 7 H), 7.23 (s, 1 H), 5.92 (br, 1 H), 4.83 (d, 2 H), 4.74-4.58 (m, 2 H), 3.77 (br, 1 H), 2.70 (br, 1 H), 2.40 (s, 3 H), 2.16 (d, 3 H), 1.88 (d, 3 H), 1.55 (d, 7 H), 1.33-0.97 (m, 4 H); API MS m/z = 470 $[\text{C}_{28}\text{H}_{35}\text{N}_7+\text{H}]^+$.

5

Example 12 - Preparation of Compound 14

DME (3 mL), tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol), and PPh_3 (0.04 g, 0.15 mmol) were placed in a round-bottomed flask with a
10 condenser under argon. Compound 11 (0.13 g, 0.25 mmol) and 3-chloro-4-fluoroboronic acid (0.15 g, 0.88 mmol) were dissolved in Na_2CO_3 (2M, 0.6 mL) and DME (1 mL) were added to the reaction mixture, refluxed for 19 h then stirred at room temperature for 2 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2 . The combined organic phases were washed with brine, dried over
15 anhydrous Na_2SO_4 , and evaporated. The crude product was purified by repeated column chromatography to yield 14 (0.019 g, 15%): ^1H NMR (300 MHz, CDCl_3) δ 7.59-7.53 (m, 1 H), 7.47-7.35 (m, 4 H), 7.26-7.14 (m, 3 H), 5.81 (br, 1 H), 4.81 (d, 2 H), 4.72-4.54 (m, 2 H), 3.72 (br, 1 H), 2.69 (br, 1 H), 2.21-2.03 (m, 3 H), 1.94-1.78 (m, 3 H), 1.54 (d, 6 H), 1.33-1.12 (m, 4 H); API MS m/z = 508 $[\text{C}_{27}\text{H}_{31}\text{ClFN}_7+\text{H}]^+$.

20

Example 13 - Preparation of Compound 16

A solution of 15 (2.5 g, 15.8 mmol) and ether was cooled to -78°C . In a separate flask, $n\text{-BuLi}$ (15.8 mmol) was also cooled to -78°C . The solution of 15
25 was added to the $n\text{-BuLi}$ solution *via* cannula to give a dark red solution. The reaction mixture was stirred for 5 min prior to the rapid addition of $(n\text{-Bu})_3\text{SnCl}$ (6.2 g, 19 mmol). The resulting bright yellow solution was stirred at -78°C for 2 h, allowed to warm to room temperature, and stirred for another 10 min. The solution was then diluted with H_2O (80 mL) and extracted with ethyl acetate (3 x 50 mL). The
30 organic extracts were combined, washed with brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo* to yield the crude product as a yellow oil. Purification by column chromatography gave the product 16 (4.89 g, 84%) as a pale yellow liquid: ^1H NMR (300 MHz, CDCl_3) δ 8.72 (d, 1 H), 7.48-7.46 (m, 1 H), 7.40-7.38 (m, 1 H),

7.11-7.09 (m, 1 H), 1.61-1.50 (m, 6 H), 1.38-1.26 (m, 6 H), 1.14-1.09 (m, 6 H), 0.97-0.77 (t, 9 H).

Example 14 - Preparation of Compound 17

5

To compound 16 (0.18 g, 0.48 mmol) was added compound 4 (0.14 g, 0.33 mmol), Pd(PPh₃)₄ (0.05 g, 0.49 mmol), and toluene (10 mL) in a sealed tube under an argon atmosphere. The solution was degassed with argon and heated at 135 °C in an oil bath for 3 h. The solution was cooled to room temperature, diluted with saturated NaHCO₃, and extracted with CH₂Cl₂ (3 x 30 mL). The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a light brown oil. The residue was purified by flash column chromatography using MeOH/CH₂Cl₂ (10%) to afford 17 as a white solid. The sample was dissolved into hexane/CH₂Cl₂/MeOH and then precipitated with diethyl ether, filtered, and rinsed several times with ether to provide in 17 (30.3 mg): mp 95-100 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.68 (d, 1 H), 7.96 (d, 2 H), 7.77-7.69 (m, 2 H), 7.49-7.45 (m, 3 H), 7.24-7.20 (m, 1 H), 5.99 (s, 1 H), 5.11 (s, 1 H), 4.88-4.83 (m, 3 H), 4.65-4.56 (m, 1 H), 3.91-3.80 (m, 2 H), 3.65-3.60 (m, 1 H), 1.66-1.52 (m, 8 H), 1.05-0.99 (t, 3 H); IR (KBr) 3411, 2968, 1601, 1489 cm⁻¹; CI MS *m/z* = 432 [C₂₄H₂₉N₇+H]⁺.

20

Example 15 - Preparation of Compound 19

To a solution of *n*-BuLi (2.5M hexane solution, 10.9 mL, 27.4 mmol) in ethyl ether 28 mL at -78 °C was added 2-bromopyridine (4.33 g, 27.4 mmol) in ethyl ether (15 mL). After stirring for 30 min, a solution of trimethylstannylchloride (6.0 g, 30 mmol) in THF (10 mL) was added. Stirring was continued at -78 °C for 2 h and the mixture was then warmed up to room temperature and filtered. The precipitate was washed with ether and the combined the ether filtrates were concentrated to give the crude product: ¹H NMR (500 Hz, CDCl₃) δ 8.69-8.68 (d, 1 H), 7.47-7.07 (m, 3 H), 0.30 (s, 9 H).

30

Example 16 - Preparation of Compound 21

A mixture of 4-bromobenzonitrile (1.68 g, 9.2 mmol), crude 2-trimethylstannylpyridine (3.33 g, 13.8 mmol), and $\text{PdCl}_2(\text{PPh}_3)_2$ (321 mg, 0.46 mmol) in DMF (25 mL) was heated at 150-155 °C in pressure tube for 24 h. The DMF was distilled off under reduced pressure and the residue was filtered through a short column of basic alumina and washed with ethyl acetate and then concentrated. Flash chromatography of the residue on silica gel gave the product (41%) as a white solid: mp 99-100 °C; ^1H NMR (500 Hz, CDCl_3) δ 8.74 (dd, $J_1 = 1$ Hz, $J_2 = 1.7$ Hz, 1 H), 8.12 (d, $J = 8.6$ Hz, 2 H), 7.83-7.76 (m, 4 H), 7.32 (m, 1 H).

Example 17 - Preparation of Compound 22

To LiAlH_4 (8 mmol) in THF (25 mL) was added 21 (0.96 g, 5.3 mmol) in THF (15 mL) slowly while the flask was cooled with ice. The mixture was stirred at room temperature for 10-30 min then stirred at reflux for 4 h under nitrogen. The mixture was cooled in an ice bath and aqueous sodium hydroxide solution (0.5 mL, 10%) was added. The mixture was stirred until the residue became white and the solid was filtered and washed with methylene chloride (4 x 5 mL). The methylene chloride solution was dried with anhydrous sodium sulfate, concentrated, and the crude product was chromatographed on silica gel to give the product as a yellow liquid. A small amount of ethanol was added and the pure amine 22 was obtained as a white solid (74%) after filtration: mp 114-117 °C; ^1H NMR (500 Hz, CDCl_3) δ 8.66 (d, $J = 4.4$ Hz, 1 H), 7.94 (d, $J = 8.1$ Hz, 2 H), 7.70 (m, 2 H), 7.39 (d, $J = 8.0$ Hz), 7.19 (m, 1 H), 3.90 (s, 2 H), 1.98 (s, 2 H).

Example 18 - Preparation of Compound 23

A mixture of 2,6-dichloropurine (1, 0.19 g, 1 mmol), amine 22 (0.39 g, 2.15 mmol) in ethanol (13 mL), and water (3.4 mL) was heated at 100-110 °C under nitrogen for 24 h and then it was cooled to room temperature. The mixture was concentrated and water (5 mL) was added. A solid was filtered and washed with water (2 x 5 mL) and dried under vacuum to give the product (93%) as yellow solid: mp 260 °C (dec); ^1H NMR (500 Hz, $\text{DMSO}-d_6$) δ 12.4 (bs, 1 H), 8.76 (m, $J = 1$ Hz, 1

H), 8.28 (s, 1 H), 8.16 (d, $J = 8.1$ Hz, 2 H), 8.03 (d, $J = 7.8$ Hz, 1 H), 7.97 (m, 1 H), 7.58 (d = 8.6 Hz, 2 H), 7.45 (m, 1 H), 4.82 (s, 2 H).

Example 19 - Preparation of Compound 24

5

To the solution of compound 23 (0.33 g, 1 mmol) in DMSO (5.2 mL), added potassium carbonate (0.7 g, 5 mmol) and 2-iodopropane (0.5 g, 3 mmol). The mixture was stirred at ambient temperature under nitrogen for 24 h and poured into ice water (30 mL). After filtration, the solid was washed with water (4 x 5 mL), dried
10 under vacuum to give the crude product as a yellow solid. Flash column chromatography of the crude product on silica gel and recrystallization provided the pure product (76%) as white crystals: mp 178-179 °C; ^1H NMR (500 Hz, CDCl_3) δ 8.68 (m, 1 H), 7.96 (d, $J = 8$ Hz, 2 H), 7.76-7.70 (m, 2 H), 7.73 (s, 1 H), 7.47 (d, $J = 8$ Hz, 2 H), 7.22 (m, 1 H), 4.89 (s, 1 H), 4.79 (m, 1 H), 1.54 (d, $J = 6.8$ Hz, 6 H); CI MS
15 $m/z = 379$ [$\text{C}_{20}\text{H}_{19}\text{ClN}_6 + \text{H}$] $^+$. Anal. Calcd. for $\text{C}_{20}\text{H}_{19}\text{ClN}_6$: C, 63.41; H, 5.05; N, 22.18. Found: C, 63.07; H, 5.01; N, 22.01.

Example 20 - Preparation of Compound 17

20

To compound 24 (0.7 g, 1.8 mmol) was added (*R*)-(-)-2-amino-1-butanol (3.5 g, 3.9 mmol) stirred in a sealed tube for 2 h at 190 °C. The reaction mixture was allowed to cool and then was partitioned between EtOAc and brine. The EtOAc was separated, washed with saturated brine (4 x), dried with Na_2SO_4 , and concentrated. The product was air dried to give an oil, then dissolved in EtOAc. The
25 EtOAc solution was cooled again, and the precipitate collected, washed with cold EtOAc (2 x), air dried, and heated *in vacuo* for 2 h to give 17 (0.54 g, 67%): mp 98-100 °C; ^1H NMR (300 MHz, CDCl_3) δ 8.00-7.85 (m, 2 H), 7.75-7.55 (m, 2 H), 7.50-7.35 (m, 3 H), 7.30-7.15 (m, 1 H), 6.40-6.20 (bs, 1 H), 5.00-4.82 (m, 1 H), 4.80-4.68 (bs, 3 H), 4.60 (heptuplet, 1 H), 3.98-3.70 (m, 2 H), 3.70-3.54 (dd, 1 H), 2.10 (bs, 1
30 H), 1.75-1.53 (m, 2 H), 1.51 (d, 6 H), 1.00 (t, 3 H); IR (KBr) 3406, 2969, 1601, 1490, 1389, 1254, 779 cm^{-1} ; API MS $m/z = 432$ [$\text{C}_{24}\text{H}_{29}\text{N}_7\text{O} + \text{H}$] $^+$.

Example 21 - Preparation of Compound 25

To compound 4 (0.14 g, 0.33 mmol) was added 3-(tributylstannyl)pyridine (0.15 g, 0.33 mmol), Pd(PPh₃)₄ (0.06 g, 0.41 mmol), and toluene (10 mL). The solution was degassed with argon for 8 min in a sealed tube, and heated in an oil bath for 3 h at 130 °C. The cooled reaction mixture was diluted with saturated NaHCO₃ and extracted with CH₂Cl₂ (3 x 50 mL). The combined organic extracts were washed with brine and dried over Na₂SO₄. The reaction mixture was purified by column chromatography on silica gel to give the desired coupling product. The product was dissolved in acetonitrile and washed with hexane (3 x 10 mL) to remove a portion of the tin contaminants. The reaction mixture was again purified by column chromatography on reversed phase silica gel to give compound 25 (0.04 g): ¹H NMR (300 MHz, CDCl₃) δ 8.83 (s, 1 H), 8.58 (d, 1 H), 7.88-7.83 (m, 1 H), 7.56-7.46 (m, 5 H), 7.38-7.33 (m, 1 H), 5.99 (s, 1 H), 5.11 (s, 1 H), 4.90-4.83 (m, 2 H), 4.63-4.56 (m, 1 H), 3.92-3.81 (m, 2 H), 3.67-3.60 (m, 1 H), 1.69-1.49 (m, 8 H), 1.05-1.00 (t, 3 H); CI MS *m/z* = 432 [C₂₄H₂₉N₇O+H]⁺.

Example 22 - Preparation of Compound 27

A mixture of diethyl(3-pyridyl)borane (26, 540 mg, 3.67 mmol), 4-bromobenzonitrile (803 mg, 4.41 mmol) and Pd(PPh₃)₄ (144 mg, 0.13 mmol) in toluene (9 mL), ethanol (1.3 mL) and 2M aqueous sodium carbonate solution (4.1 mL, 8.2 mmol) was heated at 90-100 °C under nitrogen for 27 h. The mixture was cooled to room temperature and water (10 mL) was added. The organic layer was separated and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The combined organic layers were washed with brine (2 x 15 mL) and dried over anhydrous sodium sulfate. Flash chromatography of the crude product on silica gave the product as a white solid (80%): mp 95-96 °C.

Example 23 - An Alternative Preparation of 27 is Described Below

A flask charged with 4-bromobenzonitrile (360 mg, 2.0 mmol), bis(pinacolato)diboron (560 mg, 2.2 mmol), potassium acetate (590 mg, 6.0 mmol) and PdCl₂(dppf) (49 mg, 0.06 mmol) was flushed with nitrogen and DMF (12 mL)

was added. The mixture was heated at 80-85 °C for 4 h and then cooled to room temperature at which time PdCl₂(dppf) (49 mg, 0.06 mmol), 3-bromopyridine (385 δL, 3.40 mmol), and 2M aqueous sodium carbonate solution (5 mL, 10 mmol) was added. The mixture was stirred at 80-85 °C for 24 h and extracted with ethyl ether (3
5 x 30 mL) and then washed with brine (3 x 15 mL) and dried with anhydrous sodium sulfate. Flash chromatography of the crude product on silica gel gave the product as white crystals (56%): mp 96-97 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.55 (dd, J₁ = 1 Hz, J₂ = 1.4 Hz, 1 H), 8.66 (m, 1 H), 7.90-7.87 (m, 1 H), 7.77 (d, J = 7.8 Hz, 2 H), 7.69 (d, J = 8.8 Hz, 2 H), 7.42 (m, 1 H).

10

Example 24 - Preparation of Compound 28

To LiAlH₄ (8 mmol) in THF (25 mL) was added 27 (0.96 g, 5.3 mmol) in THF (25 mL) slowly while the flask was cooled with ice. The mixture was stirred
15 at room temperature for 10-30 min then stirred at reflux for 4 h under nitrogen. The mixture was cooled in an ice bath and aqueous sodium hydroxide solution (0.5 mL, 10%) was added. The mixture was stirred until the residue became white and the solid was filtered and washed with methylene chloride (4 x 5 mL). The methylene chloride solution was dried with anhydrous sodium sulfate, concentrated, and the
20 crude product was chromatographed on silica gel to give the product as a yellow liquid. A small amount of ethanol was added and the pure amine 28 was obtained as a white solid (46%) after filtration: mp 94-96 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.74 (d, J = 2.4 Hz, 1 H), 8.48 (dd, J₁ = 1.5 Hz, J₂ = 4.7 Hz, 1 H), 7.77 (m, 1 H), 7.45 (d, J = 8.10 Hz, 2 H), 7.33 (d, J = 8.0 Hz, 2 H), 7.25 (m, 1 H), 3.83 (s, 2 H), 2.25 (s, 2 H).

25

Example 25 - Preparation of Compound 29

A mixture of 2,6-dichloropurine (1, 0.19 g, 1 mmol), amine 28 (0.4 g, 2.15 mmol) in ethanol (13 mL), water (3 mL) was heated at 100-110 °C under
30 nitrogen for 24 h and then it was cooled to room temperature. The mixture was concentrated and water (5 mL) was added. A solid was filtered and washed with water (2 x 5 mL) and dried under vacuum to give the product (92%) as a yellow solid: mp 219 °C (dec); ¹H NMR (500 Hz, DMSO-*d*₆) δ 13.2 (bs, 1 H), 8.99 (s, 1 H), 8.66

(d, $J = 3.5$ Hz, 1 H), 8.28 (s, 1 H), 8.16 (d, $J = 7.3$ Hz, 1 H), 7.80 (d, $J = 7.6$ Hz, 2 H), 7.60-7.57 (m, 3 H).

Example 26 - Preparation of Compound 30

5 To a solution of 29 (0.3 g, 1 mmol) in DMSO (5 mL), was added potassium carbonate (0.7 g, 5 mmol) and 2-iodopropane (0.5 g, 3 mmol). The mixture was stirred at ambient temperature under nitrogen for 24 h and poured into ice water (30 mL). After filtration, the solid was washed with water (4 x 5 mL), dried
10 under vacuum to give the crude product as a yellow solid. Flash column chromatography of the crude product on silica gel and recrystallization provided the pure product (76%) as white crystals: mp 178-179 °C; ^1H NMR (500 Hz, CDCl_3) δ 8.82 (d, $J = 1.3$ Hz, 1 H), 8.59-8.58 (m, 1 H), 7.86-7.84 (m, 1 H), 7.72 (s, 1 H), 7.56-7.48 (m, 4 H), 7.37-7.34 (m, 1 H), 4.88 (s, 2 H), 4.82 (m, 1 H), 1.56 (d, $J = 0.7$ Hz, 3
15 H), 1.55 (d, $J = 0.8$ Hz, 3 H); CI MS $m/z = 379$ [$\text{C}_{20}\text{H}_{19}\text{ClN}_6 + \text{H}$] $^+$. Anal. Calcd. for $\text{C}_{20}\text{H}_{19}\text{ClN}_6$: C, 63.41; H, 5.05; N, 22.18. Found: C, 63.24; H, 4.97; N, 21.93.

Example 27 - Preparation of Compound 32

20 To a mixture of 4 (0.05 g, 0.11 mmol) was added 4-(tributylstannyl)pyridine (0.06 g, 0.16 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.02 g, 0.02 mmol), and toluene (2.5 mL). The reaction mixture was degassed and heated in a sealed tube at 125 °C for 3 h. The reaction mixture was cooled to room temperature then saturated NaHCO_3 (30 mL) was added followed by extraction with CH_2Cl_2 (3 x 30). The
25 organic layer was washed with brine (50 mL), dried with MgSO_4 , and concentrated. The reaction mixture was purified by column chromatography on silica gel to give 32: ^1H NMR (300 MHz, CDCl_3) δ 8.65 (s, 2 H), 7.60-7.57 (m, 2 H), 7.49-7.45 (m, 5 H), 6.20 (s, 1 H), 4.93 (d, 1 H), 4.84 (s, 2 H), 4.65-4.57 (m, 1 H), 3.92-3.80 (m, 2 H), 3.68-3.51 (m, 1 H), 1.68-1.58 (m, 2 H), 1.52 (d, 6 H), 1.05-0.99 (t, 3 H).

30

Example 28 - Preparation of Compound 33

To compound 4 (0.18 g, 0.43 mmol) was added 4-vinylphenylboronic acid (0.19 g, 1.28 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.09 g, 0.08 mmol), Na_2CO_3 (2M, 0.85 mL),

was added toluene (5 mL). The mixture was degassed with argon for 10 min. The resulting solution was heated in a sealed tube at 135 °C for 4.5 h. The cooled solution was diluted with water and extracted with CH₂Cl₂ (3 x 50 mL). The combined organic extracts were washed with brine and dried over Na₂SO₄. The solution was purified by flash column chromatography (2 x) on silica gel to give the desired product 33 as a yellow solid (0.09 g): mp 130-131 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.57-7.42 (m, 9 H), 6.80-6.70 (dd, 1 H), 5.98 (s, 1 H), 5.79 (d, 1 H), 5.27 (d, 1 H), 4.88 (d, 1 H), 4.84-4.72 (m, 2 H), 4.63-4.56 (m, 1 H), 3.92-3.81 (m, 2 H), 3.66-3.60 (m, 1 H), 1.68-1.52 (m, 8 H), 1.05-1.00 (t, 3 H); IR (CH₂Cl₂) 3293, 2968, 1601, 1489, 1390 cm⁻¹; CI MS *m/z* = 457 [C₂₇H₃₂N₆O+H]⁺.

Example 29 - Preparation of Compound 34

To compound 33 (0.008 g, 0.016 mmol) was added OsO₄ (0.007 g, 0.026 mmol), pyridine (0.08 mL), and toluene (0.75 mL). The reaction mixture was stirred at room temperature in the dark for 1 h, concentrated *in vacuo*, and then slurried in methanol/water (9:1). Sodium metabisulfite (0.07 g) was added and the reaction was stirred for 1 h. The mixture was washed with brine, extracted with CH₂Cl₂ (3 x 10 mL), dried over Na₂SO₄, and concentrated. The product was purified by column chromatography on silica gel to give compound 34 (0.003 g) as a tan solid: ¹H NMR (300 MHz, CDCl₃) δ 7.51 (s, 1 H), 7.43-7.35 (m, 6 H), 7.25-7.22 (m, 2 H), 6.51 (s, 1 H), 4.98 (d, 1 H), 4.35-4.25 (m, 2 H), 4.64-4.54 (m, 1 H), 3.93-3.80 (m, 3 H), 3.74-3.59 (m, 3 H), 1.68-1.58 (m, 2 H), 1.52 (d, 6 H), 1.06-0.99 (t, 3 H).

Example 30 - Preparation of Compound 36

To compound 4 (0.12 g, 0.27 mmol) was added 3-aminophenylboronic acid hydrochloride (0.12 g, 0.69 mmol), and Pd(PPh₃)₄ (0.09 g, 0.75 mmol) in a sealed tube filled with argon. To this mixture was added toluene (5 mL) and Na₂CO₃ (2M, 0.55 mL). The resulting solution was degassed with argon for 5 min and placed in a 130 °C oil bath for 6 h. The cooled solution was diluted with water and extracted with CH₂Cl₂ (3 x 50 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated. The solution was purified by column chromatography on silica gel to yield 36 (0.04 g, 36%): ¹H NMR (300 MHz, CDCl₃)

δ 7.52-7.46 (m, 3 H), 7.39 (d, 2 H), 7.23-7.18 (m, 1 H), 6.96 (d, 1 H), 6.88 (t, 1 H), 6.68-6.66 (m, 1 H), 6.12 (s, 1 H), 4.90 (d, 1 H), 4.79 (s, 2 H), 4.62-4.57 (m, 1 H), 3.92-3.76 (m, 4 H), 3.66-3.60 (m, 1 H), 1.65-1.48 (m, 8 H), 1.04-0.99 (t, 3 H); CI MS $m/z = 446$ [$C_{25}H_{31}N_7O+H$] $^+$.

5

Example 31 - Preparation of Compound 38

To a suspension of $Pd(PPh_3)_4$ (0.02 g, 0.01 mmol) in anhydrous DME (8 mL) was added 4 (0.12 g, 0.27 mmol) and the mixture stirred at room temperature for 10 min. To this solution was added 3-(trifluoromethyl)phenylboronic acid (37; 0.12 g, 0.65 mmol) in a minimum of EtOH, followed by Na_2CO_3 (2M, 0.27 mL), and the resulting mixture was heated at reflux for 20 h. The cooled reaction mixture was diluted with water and extracted with CH_2Cl_2 (3 x 50 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , and concentrated. The reaction mixture was purified by column chromatography on normal phase silica gel followed by reversed phase column chromatography to obtain 38 (0.04 g, 33%) as an off white solid: mp 60-67 °C; 1H NMR (300 MHz, $CDCl_3$) δ 7.81 (s, 1 H), 7.74 (d, 1 H), 7.58-7.45 (m, 7 H), 5.98 (s, 1 H), 4.90-4.83 (m, 3 H), 4.63-4.59 (m, 1 H), 3.90-3.81 (m, 2 H), 3.66-3.60 (m, 1 H), 1.68-1.51 (m, 8 H), 1.05-1.00 (t, 3 H); IR (KBr) 3406, 2969, 1602, 1489, 1335 cm^{-1} ; CI MS $m/z = 499$ [$C_{26}H_{29}FN_7O+H$] $^+$.

20

Example 32 - Preparation of Compound 40

A mixture of 4 (0.13 g, 0.31 mmol), 2-naphthaleneboronic acid (39; 0.11 g, 0.62 mmol) and $Pd(PPh_3)_4$ (0.09 g, 0.08 mmol) was placed in a sealed tube that was filled with argon. To the mixture was added toluene (5 mL) and Na_2CO_3 (2M, 0.62 mL). The tube was quickly sealed and heated at 125 °C in an oil bath for 6 h. The cooled solution was diluted with water and extracted with CH_2Cl_2 (3 x 50 mL). The organic layers were washed with brine, dried over Na_2SO_4 , and concentrated. The reaction mixture was purified by column chromatography on normal phase silica gel, followed by reversed phase chromatography to give 40 (0.04 g, 28%): mp 70-75 °C; 1H NMR (300 MHz, $CDCl_3$) δ 8.02 (s, 1 H), 7.92-7.84 (m, 3 H), 7.74-7.67 (m, 3 H), 7.51-7.44 (m, 5 H), 5.96 (s, 1 H), 4.89-4.84 (m, 3 H), 4.66-

30

4.57 (m, 1 H), 3.93-3.82 (m, 2 H), 3.67-3.61 (m, 1 H), 1.76-1.50 (m, 8 H), 1.06-1.01 (t, 3 H); IR (KBr) 3422, 2927, 1601, 1491, 1388 cm^{-1} .

Example 33 - Preparation of Compound 43

5

To compound 4 (0.14 g, 0.33 mmol) was added 4-methoxyphenylboronic acid (**42**, 0.11 g, 0.71 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.10 g, 0.087 mmol), Na_2CO_3 (2M, 0.66 mL), and toluene (7 mL). The solution was degassed for 8 min with argon and heated in an oil bath at 125 °C for 6 h. The cooled solution was diluted with water and extracted with CH_2Cl_2 (3 x 50 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The reaction mixture was purified by normal phase column chromatography followed by reversed phase chromatography to give **43** (0.05 g, 28%) as a white solid: mp 128-130 °C; ^1H NMR (300 MHz, CDCl_3) δ 7.52-7.50 (m, 5 H), 7.41 (d, 2 H), 6.97 (d, 2 H), 5.93 (s, 1 H), 4.89-4.79 (m, 3 H), 4.63-4.56 (m, 1 H), 3.92-3.81 (m, 5 H), 3.67-3.60 (m, 1 H), 1.68-1.49 (m, 8 H), 1.05-1.00 (t, 3 H); IR (KBr) 3417, 2931, 1610, 1499, 1389 cm^{-1} ; CI MS m/z = 461 $[\text{C}_{26}\text{H}_{32}\text{N}_6\text{O}_2 + \text{H}]^+$.

15

Example 34 - Preparation of Compound 45

20

To a solution of *s*-BuLi (5 mL, 6.24 mmol) and TMEDA (1 mL) in anhydrous THF (35 mL) at -75 °C under argon was added dropwise a solution of *N,N*-diethylbenzamide (0.98 g, 5.57 mmol) in THF (5 mL). The mixture was stirred for 50 min and then treated with trimethylborate (2 mL, 17 mmol). The solution was allowed to warm to room temperature overnight. The colorless solution was cooled to 0 °C and acidified to pH = 6 with 2N HCl. The THF was removed *in vacuo* and the residue was diluted with water. This was extracted with CH_2Cl_2 (3 x 50 mL) and the combined organic layers were washed with brine, dried over Na_2SO_4 , concentrated *in vacuo*, followed by removal of trace solvent on the vacuum pump to give **45** as an off-white foamy solid: ^1H NMR (300 MHz, CD_3OD) δ 7.67-7.39 (m, 4 H), 3.88-3.69 (q, 4 H), 1.41-1.30 (t, 6 H).

30

Example 35 - Preparation of Compound 46

To compound 4 (0.14 g, 0.31 mmol) was added 2-(diethylcarbamoyl)phenylboronic acid (45, 0.29 g, 1.31 mmol), Pd(PPh₃)₄ (0.1 g, 0.09 mmol), Na₂CO₃ (2M, 0.63 mL), toluene (5 mL), and the mixture degassed with argon for 10 min. The mixture was heated in an oil bath for 5 h at 135 °C. The cooled solution was diluted with water and extracted with CH₂Cl₂ (3 x 50 mL). The organic layers were combined, washed with brine, dried over Na₂CO₃, and concentrated. The reaction mixture was purified by normal phase column chromatography on silica gel, followed by reversed phase chromatography to give 46 (0.03 g, 18%) as a yellow solid: ¹H NMR (300 MHz, CDCl₃) δ 7.49-7.36 (m, 9 H), 6.18 (s, 1 H), 4.93 (d, 1 H), 4.78 (s, 2 H), 4.64-4.55 (m, 1 H), 3.92-3.60 (m, 4 H), 3.06-2.92 (m, 2 H), 2.69-2.64 (m, 1 H), 1.68-1.51 (m, 8 H), 1.04-0.99 (t, 3 H), 0.91-0.86 (t, 3 H), 0.77-0.72 (t, 3 H); CI MS *m/z* = 530 [C₃₀H₃₉N₇O₂+H]⁺.

Example 36 - Preparation of Compound 48

To a suspension of Pd(PPh₃)₄ (0.08 g, 0.69 mmol) in DME was added 4 (0.129 g, 0.30 mmol) and the mixture stirred for 10 min at room temperature. To this was added 3-nitrophenylboronic acid (47, 0.157 g, 0.94 mmol) and Na₂CO₃ (2 M, 0.59 mL). The solution was heated at reflux under argon overnight. The cooled solution was diluted with water and extracted with CH₂Cl₂ (3 x 50 mL). The organic layers were combined, washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. The solution was purified by flash column chromatography on silica gel to give 48 (0.04 g, 29%) as a bright yellow solid: mp 73-77 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.43 (s, 1 H), 8.20 (d, 1 H), 7.89 (d, 1 H), 7.63-7.43 (m, 6 H), 6.01 (s, 1 H), 4.95-4.76 (m, 3 H), 4.68-4.58 (m, 1 H), 3.98-3.80 (m, 2 H), 3.68-3.60 (m, 1 H), 1.71-1.40 (m, 8 H), 1.02-0.98 (t, 3 H); IR (KBr) 3405, 2930, 1713, 1602, 1490, 1351 cm⁻¹; CI MS *m/z* = 476 [C₂₅H₂₉N₇O₃+H]⁺.

Example 37 - Preparation of Compound 50

To a suspension of Pd(PPh₃)₄ (0.09 g, 0.08 mmol) in DME (5 mL) was added 4 (0.14 g, 0.32 mmol) and the mixture stirred at room temperature for 15 min.

To this was added benzo[b]furan-2-boronic acid (49, 0.153 g, 0.94 mmol) and Na_2CO_3 (2 M, 0.63 mL). The solution was heated at reflux under argon overnight. The reaction mixture was cooled, diluted with water, extracted with CH_2Cl_2 (3 x 50 mL). The organic layers were combined, washed with brine, dried over Na_2SO_4 , and concentrated *in vacuo*. The solution was purified by flash column chromatography on silica gel followed by flash column chromatography on reversed phase silica to give 50 (0.09 g, 60%) as a white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.82 (d, 2 H), 7.58-7.42 (m, 5 H), 7.30-7.19 (m, 2 H), 7.01 (s, 1 H), 6.11 (s, 1 H), 4.91 (d, 1 H), 4.81 (s, 2 H), 4.62-4.58 (m, 1 H), 3.92-3.80 (m, 2 H), 3.66-3.60 (m, 1 H), 1.66-1.48 (m, 8 H), 1.04-0.99 (t, 3 H); CI MS m/z = 471 [$\text{C}_{27}\text{H}_{30}\text{N}_6\text{O}_2 + \text{H}$] $^+$.

Example 38 - Preparation of Compound 52

To compound 4 (0.46 g, 1.20 mmol) was added 1-amino-1-cyclopentanemethanol (51, 1.0 g, 8.61 mmol) and EtOH (2 mL) and the mixture was heated in an oil bath at 150 °C for 60 h. The brown solution was cooled and heated again at 150 °C for 48 h. The reaction mixture was cooled and concentrated *in vacuo*. The reaction mixture was purified by flash column chromatography on silica gel to give 52 (0.39 g, 71%) as a tan solid: ^1H NMR (300 MHz, CDCl_3) δ 7.48-7.40 (m, 3 H), 7.29-7.20 (m, 2 H), 6.88 (s, 1 H), 6.25 (s, 1 H), 5.10 (s, 1 H), 4.72 (s, 2 H), 4.63-4.51 (m, 1 H), 3.78 (s, 2 H), 2.10-1.65 (m, 8 H), 1.54 (d, 6 H); CI MS m/z = 459 [$\text{C}_{21}\text{H}_{27}\text{BrN}_6\text{O} + \text{H}$] $^+$.

Example 39 - Preparation of Compound 53

To a suspension of $\text{Pd}(\text{PPh}_3)_4$ (0.07 g, 0.06 mmol) in DME (5 mL) was added 52 (0.102 g, 0.22 mmol) and stirred at room temperature for 15 min. To this was added phenylboronic acid (0.098 g, 0.80 mmol) and Na_2CO_3 (2 M, 0.44 mL). The solution was heated at reflux under argon for 18 h. The reaction mixture was diluted with water, extracted with CH_2Cl_2 (3 x 50 mL), washed with brine, and dried over Na_2SO_4 . The solution was purified by flash column chromatography on silica gel followed by flash column chromatography on reversed phase silica gel to give 53 (0.02 g, 20%): ^1H NMR (300 MHz, CDCl_3) δ 7.59-7.31 (m, 10 H), 6.95 (s, 1 H), 5.95

(s, 1 H), 5.10 (s, 1 H), 4.79 (s, 2 H), 4.61-4.52 (m, 1 H), 3.76 (s, 2 H), 2.01-1.61 (m, 8 H), 1.54 (d, 6 H); CI MS m/z = 457 $[\text{C}_{27}\text{H}_{32}\text{N}_6\text{O}+\text{H}]^+$.

Example 40 - Preparation of Compound 54

5

To compound 3 (0.26 g, 0.67 mmol) was added *trans*-4-aminocyclohexanol hydrochloride (0.62 g, 4.11 mmol), Et_3N (0.58 mL, 4.16 mmol), and ethanol (5 mL). The mixture was heated for 5 h at 135 °C in an oil bath. The temperature increased to 150 °C and heating was continued for a further 48 h. The solution was cooled and evaporated to give a yellow oil: CI MS m/z = 459 $[\text{C}_{21}\text{H}_{27}\text{BrN}_6\text{O}+\text{H}]^+$.

10

Example 41 - Preparation of Compound 55

15

To compound 3 (0.50 g, 1.31 mmol) was added *cis*-1,2-diaminocyclohexane (1.57 mL, 13.1 mmol) and EtOH (4 mL). The mixture was heated in an oil bath at 150 °C for 6 h. The reaction mixture was concentrated *in vacuo*. The reaction mixture was purified by column chromatography on silica gel to give 55 (0.49 g, 82%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 7.43-7.40 (m, 3 H), 7.23 (d, 2 H), 6.21 (s, 1 H), 5.04 (d, 1 H), 4.72 (s, 2 H), 4.67-4.58 (m, 1 H), 4.08-4.05 (m, 1 H), 3.17-3.15 (m, 1 H), 2.08 (s, 2 H), 1.65-1.38 (m, 14 H); CI MS m/z = 458 $[\text{C}_{21}\text{H}_{28}\text{BrN}_7+\text{H}]^+$.

20

Example 42 - Preparation of Compound 56

25

To compound 55 (0.10 g, 0.22 mmol) was added 2-(tributylstannyl)pyridine (0.10 g, 0.27 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.05 g, 0.04 mmol), and toluene (5 mL). The solution was degassed with argon for 8 min and heated at 135 °C for 3 h. The cooled solution was diluted with water, extracted with CH_2Cl_2 (3 x 50 mL), and the combined organic extracts were washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The solution was followed by flash column chromatography (2 x) to give the desired product 56 (0.03 g, 36%) yellow crystalline solid: ^1H NMR (300 MHz, CDCl_3) δ 8.68 (d, 1 H), 7.96 (d, 2 H), 7.78-7.69 (m, 2 H), 7.49 (s, 1 H), 7.44 (d, 2 H), 7.23-7.18 (m, 1 H), 6.10 (s, 1 H), 5.10-5.00 (m, 1 H), 4.83

30

(s, 2 H), 4.69-4.60 (m, 1 H), 4.20-4.10 (m, 1 H), 3.27-3.13 (m, 1 H), 2.48 (s, 2 H), 1.78-1.42 (m, 14 H); CI MS m/z = 457 [$C_{26}H_{32}N_8+H$]⁺.

Example 43 - Preparation of Compound 57

5

To compound 1 (0.50 g, 1.31 mmol) was added *trans*-1,2-diaminocyclohexane (2.52 mL, 21 mmol), and EtOH (6 mL). The reaction mixture was placed in an oil bath and heated to 190 °C for 25 h. The reaction mixture was removed from the heat and cooled to room temperature, concentrated for purification.

10 The reaction mixture was purified by column chromatography on silica gel to yield 57 (520 mg, 87%) as an off white foam: ¹H NMR (300 MHz, DMSO) δ 7.95 (bs, 1 H), 7.85 (s, 1 H), 7.50 (d, 2 H), 7.34 (d, 2 H), 6.17 (d, 1 H), 4.70-4.40 (m, 1 H), 2.00-1.71 (m, 4 H), 1.70-1.52 (m, 2 H), 1.41 (d, 6 H), 1.30-0.92 (m, 4 H); API MS m/z = 460 [$C_{21}H_{28}N_7Br+H$]⁺.

15

Example 44 - Preparation of Compound 58

Compound 57 (0.15 g, 0.32 mmol) was added to a suspension of Pd(PPh₃)₄ (0.11 g, 0.1 mmol) in DME (7 mL) and stirred at room temperature for 15 min. Phenylboronic acid (0.14 g, 1.14 mmol) was added followed by the Na₂CO₃ (2M, 0.62 mmol). The reaction mixture was refluxed under argon for 18 h and allowed to stir at room temperature for 51 h. It was then diluted with water, extracted with CH₂Cl₂, washed with brine, and then extracted with CH₂Cl₂. The organic layer was evaporated, dried over anhydrous Na₂SO₄, purified by column chromatography, and placed *in vacuo* for 18 h to give 58 (0.10 g, 72%) as a white solid: ¹H NMR (300 MHz, CDCl₃) δ 7.62-7.35 (m, 10 H), 5.92 (br, 1 H), 4.83 (br, 2 H), 4.74-4.56 (m, 2 H), 3.77-3.55 (m, 1 H), 2.55-2.43 (m, 1 H), 2.16-1.91 (m, 2 H), 1.73 (br, 2 H), 1.52 (d, 6 H), 1.37-1.09 (m, 6 H); API MS m/z = 456 [$C_{27}H_{33}N_7+H$]⁺.

30 **Example 45 - Preparation of Compound 59**

To compound 57 (460 mg, 1.0 mmol) in solution with CH₂Cl₂ (2 mL) was added acetic anhydride (0.44 mL, 4.6 mmol), catalytic DMAP, and pyridine (0.5 mL). The mixture was stirred at room temperature for 2.5 h. The mixture was diluted

- 146 -

with CH₂Cl₂, washed with 2N HCl, and the combined organics were then washed with NaHCO₃. The organics were then washed with brine, dried over Na₂SO₄, filtered, and concentrated to give **59** (472 mg, 94%) as an off white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.76 (s, 1 H), 7.42 (d, 2 H), 7.29 (d, 2 H), 4.68-4.40 (m, 1 H), 4.10 (s, 3 H), 3.61-3.40 (m, 2 H), 2.15-1.80 (m, 2 H), 1.74-1.55 (m, 4 H), 1.45 (d, 6 H), 1.35-1.05 (m, 4 H); API MS *m/z* = 500 [C₂₃H₃₀BrN₇O+H]⁺.

Example 46 - Preparation of Compound 60

To a suspension of Pd(PPh₃)₄ (0.11 g, 0.1 mmol) in DME (7 mL) was added compound **59** (0.15 g, 0.3 mmol) and stirred at room temperature for 15 min under argon. Phenylboronic acid (0.13 g, 1.06 mmol) was added, followed by Na₂CO₃ (2M, 0.62 mL). The reaction mixture was refluxed under argon for 18 h. The reaction mixture was then diluted with H₂O, extracted with CH₂Cl₂, washed with brine, and extracted with CH₂Cl₂. The organic layer was dried over anhydrous Na₂SO₄, purified by column chromatography, concentrated *in vacuo* for 18 h to yield **60** (61 mg, 42%): ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.96 (s, 1 H), 7.72 (s, 1 H), 7.51 (t, 3 H), 7.40-7.28 (m, 3 H), 7.28-7.13 (m, 2 H), 5.84 (br, 1 H), 4.46 (br, 3 H), 3.47 (br, 2 H), 1.83 (br, 1 H), 1.62 (s, 4 H), 1.43 (d, 6 H), 0.12 (s, 3 H); API MS *m/z* = 498 [C₂₉H₃₅N₇O+H]⁺.

Example 47 - Preparation of Compound 61

To compound **3** (0.58 g, 1.53 mmol) was added *trans*-1,4-diaminocyclohexane (1.78 g, 15.6 mmol), and EtOH (4 mL). The mixture was heated in an oil bath at 150 °C for ca. 60 h. The reaction mixture was purified by column chromatography on silica gel to yield **61** (0.48 g, 68%) as an off white solid: mp 122-125 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.43 (s, 1 H), 7.40 (d, 2 H), 7.20 (d, 2 H), 6.27 (s, 1 H), 4.75-4.68 (m, 2 H), 4.67-4.58 (m, 2 H), 3.81-3.68 (m, 1 H), 3.45 (s, 2 H), 2.88-2.75 (m, 1 H), 2.18-2.05 (m, 2 H), 2.05-1.89 (m, 2 H), 4.52 (d, 6 H), 1.45-1.13 (m, 4 H); CI MS *m/z* = 459 [C₂₁H₂₈BrN₇+H]⁺.

Example 48 - Preparation of Compound 62

Amine 61 (53 mg, 0.12 mmol) was dissolved in CH₂Cl₂ (2 mL) and pyridine (5 mL). Acetic anhydride (0.05 g, 0.53 mmol) and DMAP (few crystals) were added. The reaction mixture was allowed to stir at room temperature for 2.25 h. The reaction mixture was diluted with CH₂Cl₂, washed with 2N HCl, NaHCO₃, dried over MgSO₄, filtered, and evaporated to yield 62 (0.05 g, 78%) as a white solid: ¹H NMR (300 MHz, CDCl₃) δ 7.50-7.20 (m, 5 H), 6.02 (br, 1 H), 5.29-5.20 (m, 1 H), 4.72 (d, 2 H), 4.66-4.54 (m, 2 H), 3.72 (br, 2 H), 2.18-2.06 (m, 2 H), 2.06-1.91 (m, 2 H), 1.97 (s, 3 H), 1.54 (d, 6 H), 1.36-1.15 (m, 4 H); API MS *m/z* = 500 [C₂₃H₃₀BrN₇O+H]⁺.

Example 49 - Preparation of Compound 64

Compound 61 (0.05 g, 0.11 mmol) was dissolved in CH₂Cl₂ (3 mL) and Et₃N (2 mL) and placed in an ice bath for 10 min. Compound 63 (0.06 g, 0.22 mmol) was dissolved in CH₂Cl₂ (2 mL), added dropwise, and rinsed with CH₂Cl₂ (1.5 mL). The ice bath was removed after 20 min and the reaction was allowed to stir for 7 d. The reaction mixture was diluted with CH₂Cl₂, washed with 2N HCl until the aqueous layer was acidic, washed with NaHCO₃, dried over MgSO₄, and evaporated. The desired product was isolated by column chromatography and dried *in vacuo* to yield 64 (0.04 g, 50%) as a green solid: ¹H NMR (300 MHz, CDCl₃) δ 8.53 (d, 1 H), 8.32-8.20 (m, 2 H), 7.59-7.35 (m, 4 H), 7.23-7.11 (m, 4 H), 6.02 (br, 1 H), 4.69-4.45 (m, 5 H), 3.57 (br, 1 H), 3.12 (br, 1 H), 2.87 (s, 1 H), 1.97 (br, 2 H), 1.75 (br, 2 H), 1.48 (d, 6 H), 1.27-0.97 (m, 4 H); API MS *m/z* = 693 [C₃₃H₃₉BrN₈O₂S+H]⁺.

Example 50 - Preparation of Compound 65

Compound 61 (0.05 g, 0.11 mmol) was dissolved in CH₂Cl₂ (3 mL) and Et₃N (2 mL) and placed in an MeOH/ice bath. Methanesulfonyl chloride (0.012 mg, 0.11 mmol) in CH₂Cl₂ (2.3 mL) was slowly added. The reaction mixture and ice bath was allowed to come to room temperature. After 1.5 h, the reaction mixture was diluted with CH₂Cl₂, washed with 2N HCl until the aqueous layer was acidic. The organic layer was washed with NaHCO₃, dried over MgSO₄, filtered, and evaporated.

- 148 -

The product was purified by column chromatography, and dried *in vacuo* for 14 h to yield **65** (13 mg, 24%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.50-7.17 (m, 5 H), 5.90 (br, 1 H), 4.75-4.57 (m, 3 H), 4.11 (d, 1 H), 3.69 (br, 1 H), 3.30 (br, 1 H), 2.99 (s, 3 H), 2.18-2.03 (m, 4 H), 1.69 (d, 6 H), 1.42-1.15 (m, 5 H); API MS m/z = 538 [$\text{C}_{22}\text{H}_{30}\text{BrN}_7\text{O}_2\text{S}+\text{H}$] $^+$.

Example 51 - Preparation of Compound 66

Compound **61** (0.05 g, 0.11 mmol) was dissolved in toluene (4 mL). 2-Acetylphenylisocyanate (0.024 g, 0.15 mmol) diluted with toluene (1 mL) and added to compound **61**. Toluene (6 mL) was added to the reaction mixture. The reaction mixture was placed under reflux for 19 h. The product was purified by column chromatography, concentrated, and dried *in vacuo* for 23 h to yield **66** (42 mg, 62%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.87-7.20 (m, 9 H), 6.41 (s, 1 H), 5.86 (br, 1 H), 4.75-4.54 (m, 4 H), 3.69 (br, 1 H), 2.60 (s, 3 H), 2.12 (br, 4 H), 1.51 (d, 6 H), 1.42-1.15 (m, 5 H); API MS m/z = 619 [$\text{C}_{30}\text{H}_{35}\text{BrN}_8\text{O}_2+\text{H}$] $^+$.

Example 52 - Preparation of Compound 67

Compound **61** (0.04 g, 0.10 mmol) was dissolved in CH_2Cl_2 (2 mL) and pyridine (0.5 mL). Cyclopropanecarbonyl chloride (0.05 g, 0.44 mmol) was added along with DMAP (small amount). The reaction mixture was allowed to stir at room temperature for 2.25 h. The reaction mixture was diluted with CH_2Cl_2 , washed with 2N HCl, saturated NaHCO_3 , dried over MgSO_4 , filtered, and evaporated. The product was isolated by column chromatography to yield **67** (0.03 g, 63%) as a white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.50-7.20 (m, 5 H), 5.96 (br, 1 H), 5.41 (d, 1 H), 4.72 (d, 2 H), 4.66-4.54 (m, 2 H), 3.72 (br, 2 H), 2.18-1.97 (m, 4 H), 1.51 (d, 6 H), 1.36-1.15 (m, 5 H), 1.06-0.88 (m, 2 H), 0.79-0.67 (m, 2 H); API MS m/z = 526 [$\text{C}_{25}\text{H}_{32}\text{BrN}_7\text{O}+\text{H}$] $^+$.

Example 53 - Preparation of Compound 69

To a solution of 4-biphenylcarboxaldehyde (1.0 g, 5.49 mmol) in MeOH (20 mL) was added NaBH_3CN (0.69 g, 11.0 mmol), and NH_4OH (15 mL) and

the mixture was stirred at room temperature overnight. To this added HCl and extracted with CHCl_3 . The resulting aqueous layer was brought to $\text{pH} > 7$ with sodium bicarbonate and then extracted with CHCl_3 . The solution was dried with MgSO_4 , filtered, and evaporated to give **69** (200 mg) as a white solid: EI MS $m/z =$
5 183 $[\text{C}_{13}\text{H}_{13}\text{N}]^+$.

Example 54 - Preparation of Compound 69

To compound **70** (2.75 g, 13.9 mmol) was added anhydrous THF (60
10 mL), heated to reflux, and kept under nitrogen. 1M Borane-THF (69.7 mL) was added dropwise to **70** through an addition funnel resulting in a homogeneous solution. The solution was refluxed for 18 h. The reaction mixture was cooled in an ice water bath and quenched with H_2O , 2N HCl (20 mL), followed by 3N NaOH (60 mL). The reaction mixture was extracted with EtOAc (3 x). The organic extracts were washed
15 with brine, and dried over sodium sulfate. The crude product was concentrated, dissolved in MeOH, and HCl gas was bubbled through the solution. The solution was filtered *in vacuo* to give **69** as a white solid: ^1H NMR (300 MHz, CD_3OD) δ 7.71 (d, 2 H), 7.63 (d, 2 H), 7.52 (d, 2 H), 7.47-7.30 (m, 3 H), 4.13 (s, 2 H).

Example 55 - Preparation of Compound 71

To compound **1** (6.8 g, 36.0 mmol) and **69** (8.0 g, 36.5 mmol) was added H_2O (60 mL) and Hünigs base (9.0 g, 70.0 mmol). The mixture was stirred and heated to reflux for 5 h during which time H_2O (50 mL) was added as the reaction
25 continued to thicken. The crude product was collected by filtration, washed with H_2O (500 mL) and EtOH (2 x 30 mL), air dried, and dried *in vacuo* to give **71** (11.1 g, 92%): mp 267-269 °C.

Example 56 - Preparation of Compound 72

30 Compound **71** (4.7 g, 14.0 mmol), K_2CO_3 (15.0 g, 109 mmol), DMSO (80 mL), and 2-iodopropane (9.4 g, 55.0 mmol) were combined and stirred overnight. H_2O and EtOAc were added. The EtOAc layer was separated and washed with brine

(3 x). The EtOAc solution was dried with MgSO_4 , concentrated, and crystallized from EtOAc to give 72 (3.5 g, 66%): mp 139-140 °C.

Example 57 - Preparation of Compound 73

5 Compound 72 (2.00 g, 5.30 mmol) and (*R*)-(-)-2-amino-1-butanol (10.8 g, 121 mmol) were combined in a sealed tube, and heated in an oil bath at 190 °C for 2 h. The solution was cooled to 60 °C, diluted in EtOAc, washed with brine (4 x), dried with Na_2SO_4 , and concentrated. Purification by column chromatography on
10 SiO_2 gave the desired product 73 (1.72 g, 75%) as a foam: ^1H NMR (300 MHz, CDCl_3) δ 7.65-7.10 (m, 9 H), 6.40-6.10 (bs, 1 H), 5.05-4.85 (m, 1 H), 4.85-4.67 (m, 1 H), 4.60 (heptuplet, 1 H), 4.00-3.70 (dd, 2 H), 3.76-3.50 (m, 1 H), 1.95 (bs, 1 H), 1.80-1.55 (m, 2 H), 1.51 (d, 6 H), 1.03 (t, 3 H); IR (CH_2Cl_2) 3301, 2969, 1601, 1488, 1389, 1255, 762, 698 cm^{-1} ; API MS m/z = 431 [$\text{C}_{25}\text{H}_{30}\text{N}_6\text{O}+\text{H}$] $^+$.

15

Example 58 - Preparation of Compound 74

 Compound 72 (0.23 g, 0.60 mmol), *cis*-1,2-diaminocyclohexane (0.72 mL, 6.0 mmol), and ethanol (2 mL) were combined in a sealed tube and heated in an
20 oil bath at 155 °C for 5 d. The ethanol was removed *in vacuo* and the crude reaction mixture was filtered through a silica plug. The reaction mixture was chromatographed on silica gel, the resulting orange solid was dissolved in CH_2Cl_2 and a portion of activated charcoal was added. The solution was filtered through a pad of celite and concentrated to give 74 as a yellow solid (0.04 g, 27%): ^1H NMR (300
25 MHz, CDCl_3) 7.59-7.31 (m, 10 H), 6.00 (s, 1 H), 5.09 (d, 1 H), 4.83 (s, 2 H), 4.68-4.62 (m, 1 H), 4.11 (s, 1 H), 3.70-3.65 (m, 2 H), 3.18-3.16 (m, 1 H), 2.02 (s, 2 H), 1.67-1.42 (m, 12 H); CI MS m/z = 456 [$\text{C}_{27}\text{H}_{33}\text{N}_7+\text{H}$] $^+$.

Example 59 - Preparation of Compound 75

30

 Compound 72 (0.17 g, 0.45 mmol), *trans*-1,4-diaminocyclohexane (0.53 g, 4.69 mmol), and EtOH (5 mL) were combined in a sealed tube and heated at 155 °C for 5 d. The EtOH was removed *in vacuo* and the crude mixture was subjected to flash chromatography on silica gel. Recrystallization from $\text{CHCl}_3/\text{MeOH}$

gave 75 (5.8 mg) as an off-white crystalline solid: mp 110-112 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.58-7.31 (m, 10 H), 5.95 (s, 1 H), 4.88-4.78 (m, 2 H), 4.69-4.60 (m, 2 H), 3.88-3.78 (m, 1 H), 3.07-2.98 (m, 1 H), 2.26-2.10 (m, 4 H), 1.62-1.52 (m, 8 H), 1.29-1.15 (m, 4 H); CI MS *m/z* = 456 [C₂₇H₃₃N₇+H]⁺.

5

Example 60 - Preparation of Compound 76

Compound 75 (0.05 g, 0.11 mmol) was dissolved in CH₂Cl₂ and the solution cooled to 0 °C under an argon atmosphere. A catalytic amount of DMAP, triethylamine (50 L, 0.36 mmol), followed by the acetyl chloride (25 L, 0.36 mmol) were added to the reaction mixture. The solution was warmed to room temperature and washed with NaHCO₃ (5%), water, and brine. The solution was dried over Na₂SO₄ and concentrated. Purification by flash chromatography on silica gel gave 76 (0.028 g, 53%) as a pale yellow solid: mp 224-225 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.59-7.31 (m, 10 H), 5.93 (s, 1 H), 5.26 (d, 1 H), 4.81 (s, 2 H), 4.65-4.58 (m, 1 H), 3.78-3.75 (m, 2 H), 2.18-1.99 (m, 4 H), 1.95 (s, 3 H), 1.77 (s, 1 H), 1.53 (d, 6 H), 1.32-1.22 (m, 4 H); CI MS *m/z* = 498 [C₂₉H₃₅N₇O+H]⁺.

Example 61 - Preparation of Compound 77

20

Compound 72 (0.15 g, 0.40 mmol), *trans*-4-aminocyclohexanol hydrochloride (0.31 g, 1.99 mmol), Et₃N (0.11 mL, 0.8 mmol), and EtOH (5 mL) were combined and heated in a sealed tube at 155 °C for 4 d. Additional *trans*-4-aminocyclohexanol hydrochloride (0.34 g, 2.2 mmol) and triethylamine (0.60 mL, 4.3 mmol) were added and the heat was resumed at 155 °C overnight. The crude product was purified by flash column chromatography to give 77 (0.036 g, 20%) as an off-white solid: mp 196-200 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.58-7.30 (m, 10 H), 5.97 (s, 1 H), 4.83-4.81 (m, 2 H), 4.66-4.60 (m, 2 H), 3.82-3.77 (m, 1 H), 3.69-3.62 (m, 1 H), 2.17-2.13 (m, 2 H), 2.01-1.97 (m, 2 H), 1.68 (s, 1 H), 1.53 (d, 6 H), 1.49-1.20 (m, 4 H); CI MS *m/z* = 457 [C₂₇H₃₃N₆O+H]⁺.

30

Example 62 - Preparation of Compound 78

To compound 61 (0.12 g, 0.26 mmol), was added compound 16 (0.12 g, 0.33 mmol), and Pd(PPh₃)₄ (0.06 g, 0.056 mmol) and toluene (5 mL). The resulting mixture was degassed for 10 min with argon. The mixture was heated at 140 °C for 3 h. The cooled solution was diluted with saturated NaHCO₃ and extracted with CH₂Cl₂ (3 x 50 mL). The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered, and concentrated to give a pale yellow oil which crystallized upon standing at room temperature. The crude product was purified by column chromatography and concentrated to give a white solid. The solid was precipitated with acetonitrile, filtered, washed with ether and hexane to give 78 (0.02 g, 18%): ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.63 (d, 1 H), 8.01 (d, 1 H), 7.93-7.83 (m, 2 H), 7.59-7.44 (m, 4 H), 7.34-7.29 (m, 1 H), 6.25 (s, 1 H), 4.70-4.60 (m, 2 H), 4.57-4.49 (m, 2 H), 3.65-3.52 (m, 1 H), 2.98-2.88 (m, 1 H), 1.98-1.90 (m, 4 H), 1.48 (d, 6 H), 1.42-1.18 (m, 6 H); CI MS *m/z* = 457 [C₂₆H₃₂N₈+H]⁺.

Example 63 - Preparation of Compound 78

To compound 24 (200 mg, 0.53 mmol) was added *trans*-1,4-diaminocyclohexane (2.00 g, 17 mmol) and EtOH (4 mL). The reagents were heated in a sealed tube in an oil bath at 170 °C for 18 h. The mixture was cooled to 60 °C and partitioned between EtOAc and brine. The EtOAc layer was separated, washed with brine (3 x), dried with Na₂SO₄, and concentrated to give 78 (0.12 g, 50%): mp 135-138 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.03-7.82 (m, 2 H), 7.80-7.58 (m, 3 H), 7.57-7.30 (m, 3 H), 7.30-7.05 (m, 1 H), 6.20 (bs, 1 H), 5.95-4.73 (m, 2 H), 4.73-4.45 (m, 2 H), 3.90-3.60 (m, 1 H), 2.80-2.52 (m, 1 H), 2.25-1.80 (m, 4 H), 1.80-1.60 (bs, 3 H), 1.52 (d, 6 H), 1.38-1.05 (m, 4 H); IR (KBr) 3422, 2927, 1599, 1489, 1253, 779 cm⁻¹; API MS *m/z* = 457 [C₂₆H₃₂N₈+H]⁺.

Example 64 - Preparation of Compound 79

Compound 78 (50 mg, 0.11 mmol) was dissolved in CH₂Cl₂ (2 mL) and stirred at room temperature. Pyridine (0.5 mL), Ac₂O (0.5 mL, 4.9 mmol), and DMAP (few crystals) were added to the reaction mixture and stirred for 2 h. The

solution was diluted in CH_2Cl_2 and washed in 2N HCl. The HCl layer was concentrated, CH_2Cl_2 was added and the aqueous phase neutralized with saturated NaHCO_3 . The CH_2Cl_2 layer was separated, dried (MgSO_4), and concentrated to give 79 (0.03 g, 55%) as a white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.00-7.80 (m, 2 H), 7.81-7.57 (m, 2 H), 7.56-7.33 (m, 3 H), 7.30-7.05 (m, 2 H), 6.15-5.90 (bs, 1 H), 5.47-5.28 (m, 1 H), 4.96-4.72 (m, 2 H), 4.73-4.45 (m, 2 H), 2.25-1.82 (m, 4 H), 2.00 (s, 3 H), 1.54 (d, 6 H), 1.40-1.00 (m, 4 H); API MS m/z = 499 [$\text{C}_{28}\text{H}_{34}\text{N}_8\text{O}+\text{H}$] $^+$.

Example 65 - Preparation of Compound 80

Compound 74 (0.02 g, 0.05 mmol) was dissolved in dry benzene (5 mL) and stirred under a blanket of argon. The solution was cooled in an ice bath and phenylisocyanate (25 L, 0.23 mmol) was added dropwise. The ice bath was removed and the mixture stirred at room temperature for 0.5 h. The solvent was evaporated *in vacuo* to give a yellow oil. The crude product was purified by flash column chromatography on silica gel to give 80 (0.008 g): ^1H NMR (300 MHz, CDCl_3) δ 7.53-7.30 (m, 10 H), 7.13-7.06 (m, 4 H), 6.98-6.88 (m, 1 H), 6.62 (s, 1 H), 6.02 (s, 1 H), 5.65 (s, 1 H), 5.02 (d, 1 H), 4.85-4.70 (m, 2 H), 4.60-4.52 (m, 1 H), 4.45-4.40 (m, 1 H), 4.36-4.22 (m, 2 H), 4.00 (s, 1 H), 1.91-1.60 (m, 6 H), 1.48-1.43 (m, 6 H).

Example 66 - Preparation of Compound 82

A mixture of 6-chloronicotinamide (2.96 g, 18.9 mmol), phenylboronic acid (2.54 g, 20.8 mmol), and $\text{Pd}(\text{PPh}_3)_4$ (643 mg, 0.565 mmol) in toluene (47 mL), ethanol (7 mL) and 2M aqueous sodium carbonate solution (21 mL, 43 mmol) was stirred and heated at 90-100 °C under nitrogen for 16 h. The mixture was cooled to room temperature and filtered. The resulting solid was washed with water (2 x 20 mL) and dried *in vacuo*. To the dried solid was added methanol (50 mL). The mixture was stirred at reflux, cooled to room temperature, and filtered to give the product (90%) as a powder: mp 218-220 °C; ^1H NMR (500 Hz, $\text{DMSO}-d_6$) δ 9.23 (d, J = 2.5 Hz, 1 H), 8.41 (dd, J_1 = 2.2 Hz, J_2 = 8.3 Hz, 1 H), 8.32 (s, 1 H), 8.27 (d, J = 7.1 Hz, 2 H), 8.20 (d, J = 8.5 Hz, 1 H), 7.74 (s, 1 H), 7.66-7.60 (m, 3 H).

Example 67 - Preparation of Compound 83

To NaBH₄ (0.19 g, 5 mmol) in 1,4-dioxane (4 mL) was added HOAc (0.3 g, 5 mmol) in 1,4-dioxane (2 mL) slowly while the flask was cooled with ice. Compound 82 (0.2 g, 1 mmol) was then added. The mixture was stirred at reflux at 100-110 °C for 4 h and the solvent was evaporated. To this mixture was added water (2 mL) slowly. The mixture was extracted with CH₂Cl₂ (4 x 10 mL), washed with water (3 x 5 mL), dried with anhydrous sodium sulfate, concentrated, and purified by flash chromatography on silica gel to provide the product as a yellow liquid. This was triturated with ethanol (1 mL) to provide a white solid which was collected (60%) and dried: mp 97-99 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.60 (d, J = 2 Hz, 1 H), 7.97-7.95 (m, 2 H), 7.72-7.67 (m, 2 H), 7.47-7.37 (m, 3 H), 3.90 (s, 2 H), 1.77 (bs, 2 H).

Example 68 - Preparation of Compound 84

A mixture of 2,6-dichloropurine (1, 0.19 g, 1 mmol), amine 83 (0.39 g, 2.15 mmol) in ethanol (13 mL), and water (3 mL) was heated at 100-110 °C under nitrogen for 24 h and then cooled to room temperature. The mixture was concentrated and water (5 mL) was added. A solid was filtered and washed with water (2 x 5 mL) and dried under vacuum to give the product (80%) as a yellow solid: mp 260 °C (dec); ¹H NMR (500 Hz, DMSO-*d*₆) δ 13.26 (s, 1 H), 8.79 (s, 1 H), 8.27 (s, 1 H), 8.16 (d, J = 7.1 Hz, 2 H), 8.34 (d, J = 7.3 Hz, 1 H), 7.96 (d, J = 7.6 Hz, 1 H), 7.63-7.52 (m, 3 H), 4.81 (s, 2 H).

Example 69 - Preparation of Compound 85

To a solution of compound 84 (0.34 g, 1 mmol) in DMSO (5 mL), was added potassium carbonate (0.7 g, 5 mmol) and 2-iodopropane (0.5 g, 3 mmol). The mixture was stirred at ambient temperature under nitrogen for 24 h and poured into ice water (30 mL). After filtration, the solid was washed with water (4 x 5 mL), dried under vacuum to give the crude product as a yellow solid. Flash column chromatography of the crude product on silica gel and recrystallization provided the pure product (63%) as ivory colored crystals: mp 138-139 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.70 (d, J = 1.5 Hz, 1 H), 7.97 (m, 2 H), 7.79 (dd, J₁ = 1.7 Hz, J₂ = 8.1 Hz, 1

H), 7.71 (s, 1 H), 7.69 (d, $J = 8.1$ Hz, 1 H), 7.48-7.39 (m, 3 H), 4.87 (s, 2 H), 4.80 (m, 1 H), 1.55 (d, $J = 6.8$ Hz, 6 H); CI MS $m/z = 379$ [$C_{20}H_{19}ClN_6 + H$] $^+$. Anal. Calcd. for $C_{20}H_{19}ClN_6$: C, 63.41; H, 5.05; N, 22.18. Found: C, 63.75; H, 5.09; N, 21.87.

5 **Example 70 - Preparation of Compound 86**

To compound 85 (0.1 g, 0.26 mmol) was added *trans*-1,4-diaminocyclohexane (1 g, 8.8 mmol) and EtOH (2 mL). The reaction mixture was heated in a sealed tube in an oil bath at 120 °C. The crude product was purified by
10 column chromatography to give 86 (0.08 g, 67%): 1H NMR (300 MHz, $CDCl_3$) δ 8.68 (d, 1 H), 7.83-7.97 (m, 2 H), 7.70-7.83 (m, 1 H), 7.55-7.73 (m, 1 H), 7.30-7.55 (m, 4 H), 6.35 (bs, 1 H), 4.72-4.95 (m, 2 H), 4.50-4.72 (m, 2 H), 3.63-3.85 (m, 1 H), 2.65-2.90 (m, 1 H), 2.37-2.63 (bs, 2 H), 1.80-2.20 (dd, 4 H), 1.53 (d, 6 H), 0.72-1.42 (m, 4 H); API MS $m/z = 457$ [$C_{26}H_{22}N_8 + H$] $^+$.

15

Example 71 - Preparation of Compound 87

Compound 86 (0.08 g, 0.18 mmol) was stirred at room temperature in CH_2Cl_2 (3 mL). Pyridine (100 mg, 0.82 mmol) was added followed by Ac_2O (100
20 mg, 0.98 mmol) and DMAP (few crystals). After 2 h, more CH_2Cl_2 (3 mL) was added and the mixture was washed carefully with 2N HCl (10 drops), and saturated $NaHCO_3$. After separation of the CH_2Cl_2 layer, the organic phase was then dried with Na_2SO_4 and concentrated to give 87 (80 mg, 92%): 1H NMR (300 MHz, $CDCl_3$) δ 8.72 (s, 1 H), 8.30-7.03 (m, 9 H), 5.75-5.38 (m, 1 H), 5.02 (bs, 1 H), 4.83 (bs, 2 H),
25 4.72-4.40 (m, 1 H), 3.73 (bs, 2 H), 2.52-1.83 (m, 4 H), 1.98 (s, 3 H), 1.52 (d, 6 H), 1.50-1.00 (m, 4 H); API MS $m/z = 499$ [$C_{28}H_{34}N_8O + H$] $^+$.

Example 72 - Preparation of Compound 88

30 Compound 85 (0.05 g, 0.13 mmol) and (*R*)-(-)-2-amino-1-butanol (0.50 g, 5.6 mmol) were combined in a sealed tube and heated in an oil bath at 190 °C for 2 h then cooled to room temperature. The mixture was partitioned between EtOAc and brine, washed with brine (3 x), dried with Na_2SO_4 , and concentrated. The mixture was allowed to stand over the weekend and then purified by column

chromatography on SiO₂ to give **88** (0.01 g, 17%) as a foam: ¹H NMR (300 MHz, CDCl₃) δ 8.70 (s, 1 H), 8.05-7.82 (m, 2 H), 7.82-7.55 (m, 2 H), 7.57-7.30 (m, 4 H), 6.55 (bs, 1 H), 5.00-4.88 (s, 1 H), 4.78 (s, 2 H), 4.60 (heptuplet, 1 H), 3.98-3.83 (m, 1 H), 3.84-3.70 (m, 1 H), 3.70-3.50 (m, 1 H), 2.90 (bs, 1 H), 1.75-1.55 (m, 2 H), 1.53 (d, 6 H), 1.00 (t, 3 H); API MS *m/z* = 432 [C₂₄H₂₉N₇O+H]⁺.

Example 73 - Preparation of Compound 89

A mixture of 6-chloronicotinamide (2.5 g, 16 mmol), crude 2-trimethylstannylpyridine (5.8 g, 24 mmol), and PdCl₂(PPh₃)₂ (560 mg, 0.8 mmol) in DMF (35 mL) was heated at 150-160 °C in a pressure tube for 17 h. The DMF was distilled off under reduced pressure and the residue was extracted with ethyl acetate (6 x 30 mL) and concentrated. The residue was treated with methanol (15 mL) and a solid separated which was filtered and dried to give the product (40%) as a powder: mp 237-240 °C; ¹H NMR (500 Hz, DMSO-*d*₆) 9.22 (d, *J* = 2.2 Hz, 1 H), 8.83 (m, 1 H), 8.57-8.53 (m, 2 H), 8.48-8.46 (m, 1 H), 8.38 (s, 1 H), 8.11-8.07 (m, 1 H), 7.78 (s, 1 H), 7.63-7.60 (m, 1 H).

Example 74 - Preparation of Compound 90

To NaBH₄ (0.2 g, 5 mmol) in 1,4-dioxane (4 mL) was added HOAc (0.29 g, 5 mmol) in 1,4-dioxane (2 mL) slowly while the flask was cooled with ice. Compound **89** (0.199 g, 1 mmol) was then added. The mixture was stirred at reflux at 100-110 °C for 4 h and the solvent was evaporated. To this mixture was added water (2 mL) slowly. The mixture was extracted with CH₂Cl₂ (4 x 10 mL), washed with water (3 x 5 mL), dried with anhydrous sodium sulfate, filtered, concentrated, and purified by flash chromatography on silica gel to provide the product as a yellow liquid. This was triturated with ethanol (1 mL) and a white solid (32%) was collected and dried: mp 109-112 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.63 (m, 1 H), 8.58 (s, 1 H), 8.32 (m, 2 H), 7.77 (m, 2 H), 7.25 (m, 1 H), 3.91 (s, 2 H), 1.94 (s, 2 H).

Example 75 - Preparation of Compound 91

A mixture of 2,6-dichloropurine (1, 0.2 g, 1 mmol), compound 90 (0.4 g, 2.2 mmol) in ethanol (13 mL), and water (3 mL) was heated at 100-110 °C under nitrogen for 24 h and then cooled to room temperature. The mixture was concentrated and water (5 mL) was added. A solid was filtered and washed with water (2 x 5 mL) and dried under vacuum to give the product (83%) as a yellow solid: mp 248 °C (dec); ¹H NMR (500 Hz, DMSO-*d*₆) δ 13.27 (s, 1 H), 8.81 (s, 1 H), 8.78 (d, J = 4.1 Hz, 1 H), 8.47 (m, 2 H), 8.28 (s, 1 H), 8.06-8.01 (m, 2 H), 7.50 (m, 1 H), 4.84 (s, 2 H).

Example 76 - Preparation of Compound 92

To the solution of compound 91 (0.35 g, 1 mmol) in DMSO (5 mL), added potassium carbonate (0.68 g, 5 mmol) and 2-iodopropane (0.49 g, 3 mmol). The mixture was stirred at ambient temperature under nitrogen for 24 h and poured into ice water (30 mL). After filtration, the solid was washed with water (4 x 5 mL), dried under vacuum to give the crude product as a yellow solid. Flash column chromatography of the crude product on silica gel and recrystallization provided the pure product (64%) as white crystals: mp 150-151 °C; ¹H NMR (500 Hz, CDCl₃) δ 8.71 (d, J = 1.9 Hz, 1 H), 8.67 (m, 1 H), 8.38-8.36 (m, 2 H), 7.86-7.79 (m, 2 H), 7.75 (s, 1 H), 7.30 (m, 1 H), 4.91 (s, 2 H), 4.82 (m, 1 H), 1.57 (d, J = 6.8 Hz, 6 H); CI MS *m/z* = 380 [C₁₉H₁₈ClN₇+H]⁺. Anal. Calcd. for C₁₉H₁₈ClN₇: C, 60.08; H, 4.78; N, 25.81. Found: C, 59.76; H, 4.72; N, 25.57.

Example 77 - Preparation of Compound 93

Compound 92 (150 mg, 0.39 mmol), *trans*-1,4-diaminocyclohexane (1.50 g, 13.1 mmol), and EtOH (30 mL) were heated to 120 °C for 26 h in a sealed tube. The mixture was cooled, additional EtOAc was added, washed with brine, dried over Na₂SO₄, and concentrated to give 93 (170 mg, 94%) as a waxy solid: ¹H NMR (300 MHz, CDCl₃) δ 8.77-8.60 (m, 1 H), 8.44-8.27 (m, 2 H), 7.90-7.75 (m, 2 H), 7.50 (s, 1 H), 7.36-7.22 (m, 2 H), 6.27 (bs, 1 H), 4.96-4.73 (m, 2 H), 4.73-4.52 (m, 2 H), 3.84-3.60 (m, 1 H), 2.80-2.57 (m, 1 H), 2.22-2.00 (m, 2 H), 2.00-1.67 (m, 5 H), 1.54 (d, 6 H), 1.38-1.05 (m, 4 H); API MS *m/z* = 458 [C₂₅H₃₁N₉+H]⁺.

Example 78 - Preparation of Compound 94

Compound 93 (0.15 g, 0.33 mmol) was dissolved in CH₂Cl₂ (6 mL) and then pyridine (0.200 g, 1.64 mmol) followed by Ac₂O (0.200 g, 1.96 mmol) and DMAP (few crystals) were added. The reaction mixture was stirred for 2 h, washed with 2N HCl and NaHCO₃, extracted with CH₂Cl₂, dried with Na₂SO₄, and concentrated to give 94 (0.17 g, 69%) as a solid: mp 141-145 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.80-8.63 (m, 1 H), 8.45-8.25 (t, 2 H), 7.95-7.73 (m, 1 H), 7.52 (s, 1 H), 7.35-7.20 (m, 2 H), 6.20 (bs, 1 H), 5.50-5.30 (m, 1 H), 4.98-4.75 (m, 2 H), 4.75-4.50 (m, 2 H), 3.84-3.60 (m, 2 H), 2.27-1.87 (m, 4 H), 2.00 (s, 3 H), 1.52 (d, 6 H), 1.40-1.10 (m, 4 H); API MS *m/z* = 499 [C₂₇H₃₃N₉O+H]⁺.

Example 79 - Preparation of Compound 95

DME (3 mL), tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol), and PPh₃ (0.04 g, 0.15 mmol) were added to a round bottomed flask equipped with a condensor and maintained under an argon atmosphere. To the solution was added compound 11 (0.13 g, 0.25 mmol). 3-Fluorobenzene boronic acid (0.123 g, 0.9 mmol) was dissolved in a solution of 2M Na₂CO₃ (0.6 mL) and DME (1 mL), and added to the reaction mixture. The mixture was stirred under argon and refluxed for 19 h then stirred at room temperature for 22 h. The reaction mixture was diluted with H₂O, extracted with CH₂Cl₂, washed with brine. The organic layer was dried over Na₂SO₄ and evaporated. The reaction mixture was purified twice by column chromatography and dried under high vacuum to give a white solid (17 mg, 14%): ¹H NMR (300 MHz, CDCl₃) δ 7.56-7.32 (m, 8 H), 7.08-6.99 (m, 1 H), 5.86 (br, 1 H), 4.83 (d, 2 H), 4.71-4.56 (m, 1 H), 3.77 (br, 2 H), 2.70 (br, 1 H), 2.12 (d, 1 H), 1.88 (d, 1 H), 1.51 (d, 6 H), 1.22 (d, 5 H), 0.94-0.70 (m, 3 H); API MS *m/z* = 474 [C₂₇H₃₂FN₇+H]⁺.

Example 80 - Preparation of Compound 96

A stock solution of acetic anhydride was made by mixing CH₂Cl₂ (16 mL), pyridine (4 mL), and Ac₂O (0.16 mL). To this stock solution (1.5 mL) was

added compound 95 (0.01 g, 0.02 mmol) followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 26 h. The reaction mixture was then diluted with CH_2Cl_2 , washed with 2N HCl until the aqueous layer was acidic, washed with NaHCO_3 , dried over MgSO_4 , evaporated, and dried *in vacuo* for 15 h to give a white solid (11 mg, 92%): ^1H NMR (300 MHz, CDCl_3) δ 8.65 (br, 1 H), 7.77-7.17 (m, 8 H), 7.11-6.99 (m, 1 H), 5.14 (br, 2 H), 4.90 (br, 1 H), 4.69 (br, 1 H), 3.78 (br, 2 H), 2.09 (br, 3 H), 1.94 (s, 2 H), 1.57 (d, 6 H), 1.42 (br, 4 H), 1.24 (s, 2 H), 0.94-0.76 (m, 1 H); CI MS m/z = 516 $[\text{C}_{29}\text{H}_{34}\text{FN}_7\text{O}+\text{H}]^+$.

10 Example 81 - Preparation of Compound 97

A stock solution of acetic anhydride was made by mixing CH_2Cl_2 (16 mL), pyridine (4 mL), and Ac_2O (0.16 mL). To this stock solution (1.5 mL) was added compound 13 (0.01 g, 0.02 mmol) followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 2 h. The reaction mixture was then diluted with CH_2Cl_2 , washed with 2N HCl until it was acidic, washed with NaHCO_3 , dried over MgSO_4 , and evaporated to give a white solid (8 mg, 89%): ^1H NMR (300 MHz, CDCl_3) δ 8.78 (d, 1 H), 8.44 (t, 1 H), 7.95 (t, 2 H), 7.69-7.45 (m, 5 H), 5.30 (br, 2 H), 4.84 (br, 1 H), 4.68 (br, 1 H), 3.78 (br, 2 H), 2.39 (s, 3 H), 2.10 (br, 4 H), 1.96 (s, 2 H), 1.57 (br, 10 H), 1.25 (s, 2 H), 0.88 (br, 1 H); API MS m/z = 512 $[\text{C}_{30}\text{H}_{37}\text{N}_7\text{O}+\text{H}]^+$.

Example 82 - Preparation of Compound 98

DME (3 mL), tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol), and PPh_3 (0.04 g, 0.15 mmol) were added to a round bottom flask equipped with condensor and maintained under an argon atmosphere. Iodide 11 (0.13 g, 0.26 mmol), and 3-chlorobenzene boronic acid (0.15 g, 0.93 mmol) was dissolved in 2M Na_2CO_3 (0.6 mL) and DME (1 mL). This was then added to the reaction mixture and refluxed for 19.5 h then stirred at room temperature for 30 h. The reaction mixture was then diluted with H_2O , extracted with CH_2Cl_2 , washed with brine, dried over Na_2SO_4 , filtered, and evaporated. The reaction mixture was purified by column chromatography (3 x) and evaporated. The product was triturated in hexanes, filtered, and dried *in vacuo* for 1 h to give a white solid (16 mg): ^1H NMR (300 MHz, CDCl_3)

δ 7.56-7.38 (m, 9 H), 6.01 (br, 1 H), 4.80 (d, 2 H), 4.71-4.62 (m, 1 H), 3.77 (br, 2 H), 2.73 (br, 1 H), 2.19-2.04 (m, 1 H), 1.94-1.85 (m, 1 H), 1.51 (d, 6 H), 1.24 (d, 5 H), 0.91-1.76 (m, 3 H); API MS m/z = 490 [$C_{27}H_{32}ClN_7+H$]⁺.

5 **Example 83 - Preparation of Compound 99**

A stock solution of acetic anhydride was made by mixing CH_2Cl_2 (16 mL), pyridine (4 mL), and Ac_2O (0.16 mL). To this solution (1.5 mL) was added compound 98 (0.01 g, 0.02 mmol), followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2 , washed with 2N HCl until the aqueous layer was acidic, washed with $NaHCO_3$, dried over $MgSO_4$, filtered, and evaporated to give a white solid (0.01 g, 83%): 1H NMR (300 MHz, $CDCl_3$) δ 7.65-7.35 (m, 8 H), 7.26-7.14 (m, 1 H), 5.23 (br, 1 H), 4.66 (br, 1 H), 3.78 (br, 2 H), 2.18-2.00 (m, 4 H), 1.94 (s, 3 H), 1.54 (d, 6 H), 1.24 (s, 5 H), 0.94-0.69 (m, 3 H); API MS m/z = 532 [$C_{29}H_{34}ClN_7O+H$]⁺.

Example 84 - Preparation of Compound 100

A stock solution of acetic anhydride was made by mixing CH_2Cl_2 (16 mL), pyridine (4 mL), and Ac_2O (0.16 mL). To compound 14 (0.02 g, 0.03 mmol) was added this solution (2 mL), followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 3 h. The reaction mixture was diluted with CH_2Cl_2 , washed with 2N HCl until the aqueous layer was acidic, washed with $NaHCO_3$, filtered, and evaporated to give a white solid (8 mg, 44%): 1H NMR (300 MHz, $CDCl_3$) δ 7.41-7.32 (m, 7 H), 7.26-7.14 (m, 1 H), 5.96 (br, 1 H), 5.23 (d, 1 H), 4.84 (br, 2 H), 4.69-4.54 (m, 1 H), 3.75 (br, 1 H), 2.21-2.12 (m, 1 H), 2.09-1.96 (m, 1 H), 1.97 (s, 3 H), 1.54 (d, 6 H), 1.36-1.15 (m, 5 H), 0.85 (br, 3 H); API MS m/z = 550 [$C_{29}H_{33}ClFN_7O+H$]⁺.

30 **Example 85 - Preparation of Compound 101**

DME (3 mL), tris(dibenzylideneacetone)dipalladium (0.01 g, 0.01 mmol), and PPh_3 (0.04 g, 0.15 mmol) were added to a round bottomed flask equipped with a condensor and maintained under an argon atmosphere. Compound 10 (0.13 g,

- 161 -

0.26 mmol) and 4-fluorobenzene boronic acid (0.13 g, 0.95 mmol) was dissolved in 2M Na₂CO₃ (0.6 mL) and DME (1 mL). This was then added to the reaction mixture and refluxed for 19 h then stirred at room temperature for 72 h. The reaction mixture was then diluted with H₂O, extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄, filtered, and evaporated. The reaction mixture was purified by column chromatography on silica gel to give a white solid (17 mg, 14%): ¹H NMR (300 MHz, CDCl₃) δ 7.56-7.38 (m, 8 H), 7.11 (t, 1 H), 5.81 (br, 1 H), 4.81 (d, 2 H), 4.69-4.57 (m, 1 H), 3.78 (br, 2 H), 2.69 (br, 1 H), 2.12 (br, 1 H), 1.88 (br, 1 H), 1.54 (d, 6 H), 1.33-1.12 (m, 5 H), 0.85 (br, 3 H); API MS *m/z* = 474 [C₂₇H₃₂FN₇+H]⁺.

10

Example 86 - Preparation of Compound 102

A stock solution of acetic anhydride was made by mixing CH₂Cl₂ (16 mL), pyridine (4 mL), and Ac₂O (0.16 mL). To the solution (1.4 mL) was added compound 101 (0.01 g, 0.02 mmol), followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 2.5 h. The reaction mixture was diluted with CH₂Cl₂, washed with 2N HCl until the aqueous layer was acidic, and washed with saturated NaHCO₃. The organic layer was dried over MgSO₄ and evaporated to give a product (3 mg). The NaHCO₃ layer was further extracted with EtOAc (2 x), the organic layers were combined, dried over MgSO₄, evaporated to give product 102 (2 mg). The products were combined using EtOAc, evaporated, and dried *in vacuo* for 15 h to give product 102 (5 mg, 50%): ¹H NMR (300 MHz, CDCl₃) δ 7.71-7.08 (m, 9 H), 5.29 (br, 2 H), 4.84 (br, 1 H), 4.66 (br, 1 H), 3.78 (br, 2 H), 2.09 (br, 4 H), 1.97 (s, 1 H), 1.57 (br, 3 H), 1.24 (d, 6 H), 0.87 (br, 5 H); API MS *m/z* = 516 [C₂₉H₃₄FN₇O+H]⁺.

25

Example 87 - Preparation of Compound 103

Compound 30 (0.10 g, 0.27 mmol) and *trans*-1,4-diaminocyclohexane (0.48 g, 4.2 mmol) were combined with EtOH (2 mL) in a sealed tube and heated at 190 °C for 24 h, and then stirred at room temperature for 46 h. The reaction mixture was purified by column chromatography and dried *in vacuo* to give 103 as a white solid (0.10 g, 81%): ¹H NMR (300 MHz, CDCl₃) δ 8.83 (d, 1 H), 8.58 (t, 1 H), 7.87-7.83 (m, 1 H), 7.55-7.47 (m, 5 H), 7.38-7.33 (m, 1 H), 5.96 (br, 1 H), 4.82 (d, 2 H),

30

- 162 -

4.68-4.59 (m, 1 H), 3.75 (br, 2 H), 2.69 (br, 1 H), 2.14 (d, 2 H), 1.86 (d, 2 H), 1.54 (d, 6 H), 1.31-1.18 (m, 5 H); API MS m/z = 457 $[\text{C}_{26}\text{H}_{32}\text{N}_8+\text{H}]^+$.

Example 88 - Preparation of Compound 104

5 A stock solution of acetic anhydride was made by mixing CH_2Cl_2 (16 mL), pyridine (4 mL), and Ac_2O (0.16 mL). To the solution (3.1 mL) was added compound 103 (0.02 g, 0.04 mmol), followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 2.5 h. The reaction mixture was
10 evaporated, dried *in vacuo* for 19 h, and purified by column chromatography to give a white solid (0.02 g): ^1H NMR (300 MHz, CDCl_3) δ 8.83 (d, 1 H), 8.59 (t, 1 H), 7.85 (d, 1 H), 7.55-7.47 (m, 5 H), 7.38-7.34 (m, 1 H), 5.89 (br, 1 H), 5.25 (d, 2 H), 4.85 (br, 1 H), 4.66-4.61 (m, 1 H), 3.77 (br, 2 H), 2.15 (br, 2 H), 2.05 (br, 2 H), 1.97 (s, 2 H), 1.54 (d, 6 H), 1.33-1.25 (m, 5 H), 0.88 (br, 1 H); API MS m/z = 499
15 $[\text{C}_{28}\text{H}_{34}\text{N}_8\text{O}+\text{H}]^+$.

Example 89 - Preparation of Compound 106

Compound 72 (0.30 g, 0.80 mmol) and compound 105 (1.15 g, 6.50
20 mmol) (Gardiner, J.M., et al. Tetrahedron, 42(11):515 (1995), which is hereby incorporated by reference, were combined with EtOH (7 mL) and allowed to reflux for 23 h. Triethylamine (1 mL) was added and the reaction was refluxed further for another 21 h. The reaction mixture was then transferred to a sealed tube and EtOH (3 mL) was added. The reaction mixture was heated further at 100 °C for 3 h. The
25 mixture was purified by column chromatography to give 105 (0.13 g): ^1H NMR (300 MHz, CDCl_3) δ 7.57-7.26 (m, 10 H) 5.58 (br, 1 H), 5.10 (br, 1 H), 4.83 (br, 1 H), 4.69-4.62 (m, 2 H), 3.36-2.91 (m, 5 H), 2.82-2.65 (m, 2 H), 1.53 (d, 2 H), 1.44 (s, 9 H), 1.25 (d, 1 H), 1.13 (d, 3 H); CI MS m/z = 416 $[\text{C}_{29}\text{H}_{39}\text{N}_7\text{O}-\text{Boc}+\text{H}]^+$.

Example 90 - Preparation of Compound 107

To compound 106 (0.10 g, 0.18 mmol) was added Et_2O (2 mL), CH_2Cl_2 (1 mL) and MeOH (1 mL). During 16 h HCl/ether (1M, 5 mL) was added while stirring. The resulting precipitate was collected by filtration and dried *in vacuo*

- 163 -

for 30 min to provide 106 as an off-white solid (60 mg, 81%): ^1H NMR (300 MHz, DMSO) δ 8.48 (br, 2 H), 8.15 (br, 1 H), 7.67-7.27 (m, 10 H), 4.79 (br, 1 H), 3.60-3.42 (m, 3 H), 3.18-3.06 (m, 2 H), 3.03-2.91 (m, 2 H), 1.52 (d, 2 H), 1.27 (d, 6 H); CI MS $m/z = 416$ [$\text{C}_{24}\text{H}_{29}\text{N}_7 + \text{H}$] $^+$.

5

Example 91 - Preparation of Compound 108

A stock solution of acetic anhydride was made by mixing CH_2Cl_2 (16 mL), pyridine (4 mL), and Ac_2O (0.16 mL). To this solution (5.6 mL) was added
10 compound 107 (0.04 g, 0.09 mmol), followed by DMAP (few crystals). The reaction mixture was allowed to stir at room temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2 , washed with 2N HCl until acidic, the organic layer was washed with NaHCO_3 , dried over MgSO_4 , filtered, and evaporated to give a white solid (16 mg). The product was purified by column chromatography to provide 108 as a white
15 solid (0.01 g, 18%): ^1H NMR (300 MHz, CDCl_3) δ 7.58-7.43 (m, 10 H), 6.60 (br, 1 H), 5.91 (br, 1 H), 5.04 (t, 1 H), 4.84 (br, 2 H), 4.72-4.59 (m, 1 H), 4.10-4.02 (m, 1 H), 3.59-3.47 (m, 2 H), 1.80 (s, 3 H), 1.57 (d, 6 H), 1.19 (d, 3 H); CI MS $m/z = 458$ [$\text{C}_{26}\text{H}_{31}\text{N}_7\text{O} + \text{H}$] $^+$.

Example 92 - Preparation of Compound 109

Compound 61 (1.0 g, 2.18 mmol), 3-chlorophenylboronic acid (1.3 g, 8.16 mmol), PPh_3 (0.3 g, 1.26 mmol), 2M Na_2CO_3 (5.0 mL), and DME (54 mL) were added to a three-necked round-bottomed flask. The mixture was degassed with argon
25 and heated to reflux for 40 min, cooled to room temperature, and then $\text{Pd}_2(\text{dba})_3$ (0.08 g, 0.08 mmol) was added. The reaction mixture was heated at reflux for 7 h. 3-Chlorophenylboronic acid (0.6 g) and $\text{Pd}_2(\text{dba})_3$ (0.08 g) was then added and reflux continued for 12 h. The reaction mixture was cooled to room temperature, diluted with H_2O (50 mL), and extracted with CH_2Cl_2 (3 x 50 mL). The combined organic
30 phases were washed with H_2O (50 mL), brine (50 mL), dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography and concentrated *in vacuo* to obtain compound 109 (950 mg, 89%): mp 178-181 $^\circ\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ 7.56 (s, 1 H), 7.42-7.54 (m, 6 H), 7.26-7.35 (m, 2 H), 6.08 (bs, 1 H), 4.81 (bs, 2 H), 4.59-4.64 (m, 2 H), 3.75-3.81 (m, 1 H), 2.65-2.72 (m, 1

- 164 -

H), 2.12 (d, 2 H), 1.88 (d, 2 H), 1.53 (d, 6 H), 1.18-1.27 (m, 4 H); CI MS m/z = 490 $[\text{C}_{27}\text{H}_{32}\text{ClN}_7+\text{H}]^+$.

Example 93 - Preparation of Compound 110

5

Compound 109 (500 mg, 1.02 mmol) was dissolved in anhydrous CH_2Cl_2 (30 mL), cooled with an ice-water bath, followed by the addition of DMAP (12.2 mg, 0.1 mmol), pyridine (124 μL , 1.53 mmol), and Ac_2O (106 μL , 1.12 mmol). The reaction mixture was stirred for 30 min at 0 °C an ice-water bath then stirred
10 another 2 h at room temperature. The reaction mixture was then concentrated *in vacuo* and the residue was purified by column chromatography on silica gel. After removal of the solvent, the residue was dried *in vacuo* to give 110 (339 mg, 63%):
mp 198-200 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.57 (s, 1 H), 7.39-7.53 (m, 6 H),
7.27-7.37 (m, 2 H), 6.31 (bs, 1 H), 5.28 (d, 1 H), 4.78 (bs, 2 H), 4.70 (d, 1 H), 4.58-
15 4.67 (m, 1 H), 3.72-3.83 (m, 1 H), 2.18 (d, 2 H), 2.00 (d, 2 H), 1.90 (s, 3 H), 1.51 (d, 6 H), 1.18-1.31 (m, 4 H); CI MS m/z = 532 $[\text{C}_{29}\text{H}_{34}\text{ClN}_7\text{O}+\text{H}]^+$.

Example 94 - Preparation of Compound 111

20

Compound 61 (1.0 g, 2.18 mmol), 2-thiopheneboronic acid (1.0 g, 8.16 mmol), PPh_3 (0.3 g, 1.26 mmol), 2M Na_2CO_3 (5.0 mL), $\text{Pd}_2(\text{dba})_3$ (0.08 g, 0.08 mmol), and DME (54 mL) were added to a round-bottomed flask and purged with argon. The reaction mixture was heated at reflux for 24 h. 2-Thiopheneboronic acid (0.5 g), $\text{Pd}_2(\text{dba})_3$ (0.1 g), and 2M Na_2CO_3 (2 mL) were added and heated to reflux for
25 another 24 h. The reaction mixture was cooled to room temperature, diluted with H_2O (50 mL) and extracted with CH_2Cl_2 (3 x 50 mL). The organic phase was washed with H_2O (50 mL) and brine (50 mL), dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was repeatedly chromatographed on silica gel to obtain 111 (574 mg, 59%): mp 109-110 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.56 (d, 2 H), 7.54 (s, 1
30 H), 7.46 (d, 2 H), 7.24-7.37 (m, 2 H), 7.06 (t, 1 H), 6.04 (bs, 1 H), 4.78 (bs, 2 H), 4.59-4.69 (m, 2 H), 3.75-3.81 (m, 1 H), 2.67-2.74 (m, 1 H), 2.14 (d, 2 H), 1.87 (d, 2 H), 1.52 (d, 6 H), 1.17-1.29 (m, 4 H); CI MS m/z = 462 $[\text{C}_{25}\text{H}_{31}\text{N}_7\text{S}+\text{H}]^+$.

Example 95 - Preparation of Compound 112

Compound 111 (410.0 mg, 0.89 mmol) was dissolved in anhydrous CH_2Cl_2 (30 mL) and purged with N_2 and cooled with an ice-water bath. Pyridine (108 mg, 1.34 mmol) and DMAP (10.9 mg, 0.09 mmol) followed by Ac_2O (92 μL , 0.98 mmol) were added slowly. The reaction mixture was stirred for 30 min in an ice-water bath followed by 2 h at room temperature. The reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel to give 112 (325 mg, 73%): mp 237-244 $^\circ\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ 7.54 (d, 2 H), 7.50 (s, 1 H), 7.36 (d, 2 H), 7.24-7.37 (m, 2 H), 7.08 (t, 1 H), 6.06 (bs, 1 H), 5.34 (s, 1 H), 4.78 (bs, 2 H), 4.58-4.70 (m, 2 H), 3.78 (bs, 2 H), 2.17 (d, 2 H), 2.04 (d, 2 H), 1.96 (s, 3 H), 1.56 (d, 6 H), 1.18-1.32 (m, 4 H); CI MS m/z = 504 [$\text{C}_{27}\text{H}_{33}\text{N}_7\text{OS}+\text{H}$] $^+$.

Example 96 - Preparation of Compound 113

Compound 12 (600 mg, 1.30 mmol) was dissolved in anhydrous CH_2Cl_2 (40 mL), purged with N_2 , and cooled to 0 $^\circ\text{C}$ followed by an addition of DMAP (15.9 mg, 0.13 mmol), pyridine (165.3 mg, 1.95 mmol), and Ac_2O (135 mg, 1.43 mmol). The mixture was stirred 30 min at 0 $^\circ\text{C}$ then 2 h at room temperature. The reaction mixture was concentrated *in vacuo*. The residue was chromatographed on silica gel to give 113 (495 mg, 76%): mp 248-253 $^\circ\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ 7.54 (d, 2 H), 7.46 (s, 1 H), 7.35-7.41 (m, 5 H), 6.13 (bs, 1 H), 5.28 (d, 1 H), 4.78 (br, 2 H), 4.61-4.63 (m, 2 H), 3.75 (bs, 2 H), 2.14 (d, 2 H), 1.97 (d, 2 H), 1.95 (s, 3 H), 1.52 (d, 6 H), 1.15-1.37 (m, 4 H); CI MS m/z = 504 [$\text{C}_{27}\text{H}_{33}\text{N}_7\text{OS}+\text{H}$] $^+$.

Example 97 - Preparation of Compound 114

To compound 61 (1.0 g, 2.18 mmol) was added PPh_3 (330 mg, 1.26 mmol), $2\text{M Na}_2\text{CO}_3$ (5 mL), DME (54 mL), and 4-carboxyphenylboronic acid (1.0 g, 6.03 mmol). The mixture was purged with N_2 for 45 min then $\text{Pd}_2(\text{dba})_3$ (366 mg, 0.4 mmol) was added and the mixture was heated at reflux for 3 d. The reaction mixture was diluted with H_2O (100 mL). The aqueous layer was separated, and washed with CH_2Cl_2 (3 x 40 mL). The aqueous layer was adjusted the pH to 5.8 by using 1N HCl. Some precipitate appeared. The mixture was stored in a freezer overnight. The

- 166 -

precipitate was collected and dried to obtain 114 (450 mg, 41%): mp 246-249 °C (dec.); ¹H NMR (500 MHz, CD₃OD+NaOD) δ 7.84 (s, 2 H), 7.64 (s, 1 H), 7.54-7.63 (m, 4 H), 7.39 (s, 2 H), 6.08 (bs, 1 H), 4.85 (bs, 2 H), 4.73 (s, 1 H), 3.76 (m, 1 H), 2.74 (m, 1 H), 1.99 (s, 2 H), 1.88 (s, 2 H), 1.63 (d, 6 H), 1.21-1.36 (m, 4 H); CI MS
5 $m/z = 500$ [C₂₈H₃₃N₇O₂+H]⁺.

Example 98 - Preparation of Compound 115

To a cooled MeOH (20 mL) solution was slowly added TMSCl (253
10 μL, 2.0 mmol). The solution was stirred 20 min, followed by the addition of 114 (100 mg, 0.2 mmol). The reaction mixture was stirred at room temperature for 24 h. The reaction mixture was cooled with an ice-water bath then Et₃N (557 mL) was added. The mixture was concentrated *in vacuo*, to provide the crude product, which was washed with water (2 x 20 mL). The residue was purified by chromatography on a
15 silica gel. After removal of the solvent and drying *in vacuo*, the residue was dissolved in MeOH (5 mL), followed by the addition of ether (10 mL). The precipitate was collected and dried to provide 115 (75 mg, 73%): mp 194-197 °C; ¹H NMR (500 MHz, CD₃OD) δ 8.07 (d, 2 H), 7.80 (s, 1 H), 7.72 (d, 2 H), 7.63 (d, 2 H), 7.46 (d, 2 H), 4.63-4.79 (m, 1 H), 3.91 (s, 3 H), 3.65-3.77 (m, 1 H), 3.07 (bs, 1 H), 2.12 (d, 2 H),
20 2.01 (d, 2 H), 1.55 (d, 6 H), 1.29-1.49 (m, 4 H); API MS $m/z = 514$ [C₂₉H₃₅N₇O₂+H]⁺.

Example 99 - Preparation of Compound 117

To a suspension of compound 114 (250 mg, 0.50 mmol), pyridine (60
25 μL, 0.75 mmol), and DMAP (6.1 mg, 0.05 mmol) in H₂O-dioxane (2:1, 40 mL) was added Ac₂O (57 μL, 0.60 mmol). After stirring 4 h at room temperature, K₂CO₃ (100 mg) was added followed by additional Ac₂O (100 μL). The reaction mixture was stirred 2 h at room temperature. Water (50 mL) was added and the pH was adjusted to 5. The precipitate was collected, washed with water and ether, and dried *in vacuo*.
30 The precipitate (200 mg) was added to a solution of TMSCl (500 μL, 3.94 mmol) in MeOH (25 mL). The reaction mixture was stirred 24 h at room temperature. The mixture was concentrated *in vacuo*. The product was purified by silical gel chromatography to provide 117 (145 mg, 52%): mp 247-250 °C; ¹H NMR (500 MHz,

- 167 -

CDCl₃) δ 8.09 (d, 2 H), 7.64 (d, 2 H), 7.58 (d, 2 H), 7.49 (s, 1 H), 7.45 (d, 2 H), 5.91 (bs, 1 H), 5.18 (d, 1 H), 4.83 (bs, 2 H), 4.61-4.68 (m, 2 H), 3.93 (s, 3 H), 3.67-3.78 (m, 2 H), 3.07 (bs, 1 H), 2.16 (d, 2 H), 2.02 (d, 2 H), 1.95 (s, 3 H), 1.54 (d, 6 H), 1.23-1.32 (m, 4 H); API MS m/z = 556 [C₃₁H₃₇N₇O₃+H]⁺.

5

Example 100 - Preparation of Compound 116

To a solution of compound 117 (90 mg, 0.16 mmol) in MeOH-H₂O (6:1, 23 mL) was added KOH (11 mg, 0.19 mmol) in 5 mL MeOH. The reaction mixture was refluxed for 24 h. After removal of the solvent the residue was dissolved in 15 mL of water and washed with CH₂Cl₂. The aqueous layer was separated and adjusted pH to 4.5 by using 1N HCl. The precipitate was collected and dried to obtain 116 (60 mg, 68%): mp 344-347 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.21 (bs, 1 H), 8.14 (d, 2 H), 7.64-7.88 (m, 6 H), 7.47 (d, 2 H), 6.06 (bs, 1 H), 5.18 (d, 1 H), 4.85 (bs, 2 H), 4.51-4.66 (m, 1 H), 3.62 (bs, 1 H), 3.46 (bs, 1 H), 1.89 (bs, 2 H), 1.77 (bs, 5 H), 1.95 (s, 3 H), 1.47 (d, 6 H), 1.23-1.36 (m, 4 H); API MS m/z = 542 [C₃₀H₃₅N₇O₃+H]⁺.

Example 101 - Preparation of Compound 118

Compound 61 (1.0 g, 2.18 mmol), 3-carboxyphenylboronic acid (1.0 g, 6.03 mmol), 2N Na₂CO₃ (5 mL), and DME/EtOH (50 mL) were mixed together and degassed with N₂ for 1 h. Pd₂(dba)₃ (366.0 mg, 0.4 mmol) and PPh₃ (330.0 mg, 1.26 mmol) were added and the reaction mixture was heated to reflux for 48 h. The reaction mixture was cooled to room temperature, diluted with CH₂Cl₂ (50 mL), and extracted with aqueous 5% Na₂CO₃ (3 x 30 mL). The combined washes were extracted with CH₂Cl₂ (3 x 30 mL) and ether (40 mL). The aqueous phase was neutralized to a pH of 5.8 using 1N HCl and kept in a freezer for 1 h. The precipitate was collected, suspended in MeOH (30 mL) and the insolubles were removed by filtration. To the MeOH solution was added ether (20 mL) to precipitate the product. The white solid was collected and dried *in vacuo* to offer 118 (65 mg, 6%): mp 205-208 °C; ¹H NMR (500 MHz, CD₃OD+NaOD) δ 8.17 (s, 1 H), 7.88 (d, 1 H), 7.80 (s, 1 H), 7.56-7.63 (m, 3 H), 7.35-7.41 (m, 3 H), 6.08 (bs, 1 H), 4.80 (bs, 2 H), 4.59-4.75

(m, 1 H), 3.72-3.82 (m, 1 H), 2.89-3.01 (m, 1 H), 1.90-1.99 (m, 4 H), 1.51 (d, 6 H), 1.29-1.40 (m, 2 H), 1.12-1.23 (m, 2 H); API MS $m/z = 500$ $[\text{C}_{28}\text{H}_{33}\text{N}_7\text{O}_2 + \text{H}]^+$.

Example 102 - Preparation of Compound 119

5 3-Thiopheneboronic acid (4.5 g, 35.2 mmol) and 6-chloronicotinamide (5.0 g, 32.0 mmol) were dissolved in DMA (150 mL), followed by the addition of 2N Na_2CO_3 (23 mL). N_2 gas was passed through the mixture for 1 h. $\text{Pd}(\text{PPh}_3)_4$ (0.74 g, 0.64 mmol) was added and the reaction mixture was heated to reflux for 24 h. The
10 reaction mixture was cooled to room temperature and poured into an ice-water (1 L) and stirred for 10 min. The precipitate was collected and washed with acetone. The collected solid was suspended in EtOAc (150 mL) and heated to reflux for 5 min. The solid was filtered and collected. After drying *in vacuo*, **119** (4.5 g, 69%) was
obtained: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 9.08 (s, 1 H), 8.34 (s, 1 H), 8.28 (d, 1 H),
15 8.20 (bs, 1 H), 7.99 (d, 1 H), 7.81 (d, 1 H), 7.71 (d, 1 H), 7.60 (bs, 1 H).

Example 103 - Preparation of Compound 120

To compound **119** (4.08 g, 20.0 mmol) suspended in THF (50 mL),
20 was added 1M BH_3 -THF (164 mL). The mixture was heated to reflux for 9 h. The mixture was cooled with an ice-water bath and adjusted to a pH of 1-2, and stirred for 1 h at room temperature. The pH was adjusted to 9-10 (2N NaOH) and extracted with EtOAc (3 x 50 mL). The combined organic phases were washed with H_2O (50 mL), brine (50 mL), and dried over Na_2SO_4 . After filtration and removal of the solvent, the
25 residue was dissolved in EtOH (50 mL), followed by the addition of 1M HCl/ether (20 mL). The mixture was concentrated to dryness to provide **120** (2.03 g, 45%): ^1H NMR (500 MHz, CD_3OD) δ 8.93 (s, 1 H), 8.61 (d, 1 H), 8.51 (s, 1 H), 8.43 (d, 1 H), 7.81 (d, 1 H), 7.70 (d, 1 H), 3.30 (t, 2 H).

Example 104 - Preparation of Compound 121

30 Compound **120** (2 g, 8.82 mmol), 2,6-dichloropurine (1.5 g, 8.01 mmol), EtOH (50 mL), and $(i\text{-Pr})_2\text{NEt}$ (3.8 mL, 22 mmol) were heated at reflux for 16 h. The reaction mixture was then cooled with an ice-water bath. The precipitate was

collected and washed with EtOH, H₂O, and ether. The precipitate was dried *in vacuo* to obtain **121** (0.84 g, 31%): ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.02 (bs, 1 H), 8.76 (bs, 1 H), 8.63 (s, 1 H), 8.07 (bs, 2 H), 7.79 (bs, 2 H), 7.71 (d, 1 H), 7.64 (d, 1 H), 4.68 (bs, 2 H).

5

Example 105 - Preparation of Compound 122

Compound **121** (950 mg, 2.77 mmol) was dissolved in DMSO (50 mL), and then K₂CO₃ (2.07 g, 15.0 mmol) was added, followed by the addition of 2-iodopropane (830 L, 8.31 mmol). The reaction mixture then was stirred at room temperature overnight. The reaction mixture was poured into an ice-water bath (400 mL), stirred for 10 min, and extracted with EtOAc (4 x 50 mL). The combined organic phases were washed with H₂O (40 mL), brine (40 mL), and dried over MgSO₄. After filtration and removal of the solvent, the residue was dissolved in hot EtOAc (40 mL), followed by the addition of hexanes (80 mL). The precipitate was collected and dried *in vacuo* to obtain **122** (798 mg, 90%): ¹H NMR (500 MHz, CDCl₃) δ 8.64 (s, 1 H), 7.83 (s, 1 H), 7.70-7.79 (m, 2 H), 7.60 (d, 1 H), 7.55 (d, 1 H), 7.36 (d, 1 H), 6.11 (bs, 1 H), 4.77-4.96 (m, 3 H), 1.53 (d, 6 H).

Example 106 - Preparation of Compound 123

Compound **122** (780.0 mg, 2.03 mmol), *trans*-1,4-diaminocyclohexane (2.3 g, 20.3 mmol), and EtOH (4 mL) were heated in a sealed tube to 150 °C for 20 h. The reaction mixture was poured into ice-water (150 mL) and stirred for 10 min. The resulting precipitate was washed with H₂O (2 x 20 mL) and dried. The solid was chromatographed on a silica gel column. After removal of the solvent and drying *in vacuo*, **123** (765 mg) was obtained: mp 78-81 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.63 (s, 1 H), 7.87 (s, 1 H), 7.72 (d, 1 H), 7.64 (d, 1 H), 7.55 (d, 1 H), 7.04-7.09 (m, 1 H), 6.92 (s, 1 H), 5.95 (bs, 1 H), 4.64 (bs, 2 H), 4.33-4.45 (m, 2 H), 3.74-3.77 (m, 1 H), 2.67-2.76 (m, 1 H), 2.13 (d, 2 H), 1.90 (d, 2 H), 1.63 (bs, 2 H), 1.54 (d, 6 H), 1.19-1.30 (m, 4 H); ¹³C NMR (CDCl₃) δ 159.1, 155.0, 152.7, 151.3, 149.3, 143.3, 142.3, 136.2, 134.8, 133.4, 126.4, 126.4, 123.5, 120.2, 114.8, 50.4, 50.3, 46.5, 42.0, 35.7, 32.3, 22.8; API MS *m/z* = 463 [C₂₄H₃₀N₈S+H]⁺.

Example 107 - Preparation of Compound 124

To an ice-cold solution of compound 123 (420 mg, 0.91 mmol) in
5 CH₂Cl₂ (20 mL) was added pyridine (110 µL, 1.4 mmol), DMAP (11.0 mg, 0.09
mmol) and Ac₂O (94.2 µL, 1 mmol). The reaction mixture was stirred for 30 min at 0
°C, followed by 2 h at room temperature. After removal of the solvent, the residue
was chromatographed on a silica gel column. The resulting solid was recrystallized
with EtOAc/MeOH and dried *in vacuo* to give 124 (350 mg, 79%): mp 249-252 °C;
10 ¹H NMR (500 MHz, CDCl₃) δ 8.61 (s, 1 H), 7.85 (s, 1 H), 7.70 (d, 1 H), 7.62 (d, 1 H),
7.53 (d, 1 H), 7.48 (s, 1 H), 7.38 (d, 1 H), 6.00 (bs, 1 H), 5.25 (d, 1 H), 4.77 (bs, 2 H),
4.53-4.72 (m, 2 H), 3.68-3.77 (m, 2 H), 2.10 (d, 2 H), 2.00 (d, 2 H), 1.94 (s, 3 H), 1.52
(d, 6 H), 1.17-1.28 (m, 4 H); ¹³C NMR (CDCl₃) δ 169.4, 159.0, 155.0, 152.8, 149.2,
142.8, 142.3, 136.1, 134.9, 133.4, 126.5, 126.4, 123.5, 120.2, 114.9, 50.1, 48.3, 46.5,
15 42.2, 32.2, 32.1, 22.8; API MS *m/z* = 505 [C₂₆H₃₂N₈OS+H]⁺.

Example 108 - Alternative Preparation of Compound 71

To a solution of 4-phenylbenzoic acid (5.46 g, 27.6 mmol) in
20 methylene chloride (66 mL) was added 2 drops of DMF and oxalyl chloride (2.80 mL,
30.3 mmol). The reaction mixture was stirred overnight and added to a stirred solution
of ice and ammonium hydroxide. The resulting precipitate was filtered, washed with
methylene chloride, and triturated with water. The product was collected by filtration
and dried *in vacuo* to yield 4-phenylbenzamide (3.88 g, 71%).

25 Under a nitrogen atmosphere, 4-phenylbenzamide (2.01 g, 10.2 mmol)
was dissolved in THF (50 mL) and heated to reflux. To the mixture was added
dropwise 1 M borane in THF (80.0 mL, 80.0 mmol). After refluxing for 18 h, the
reaction mixture was cooled to room temperature and treated with 1 N HCl (40 mL).
The solution was made basic via addition of 3 N NaOH (60 mL) and extracted with
30 ethyl acetate (3 x 370 mL). The extract was washed with brine and dried over sodium
sulfate. Concentration yielded 4-phenylbenzyl amine as a white solid (1.73 g, 93%).

4-Phenylbenzyl amine (1.73 g, 10.1 mmol) and 2,6-dichloropurine
(1.94 g, 10.1 mmol) was dissolved in water (110 mL). To the solution was added *N,N*-

- 171 -

diisopropylethylamine (3.54 mL, 20.2 mmol). The reaction mixture was heated to reflux for 5 h and cooled to room temperature. A precipitate was collected by filtration. The solid was washed with water and ethanol and dried to yield **71** (2.35 g, 69%): ^1H NMR (300 MHz, $(\text{CD}_3)_2\text{SO}$) δ 8.69 (brs, 1 H), 8.15 (s, 1 H), 7.57-7.68 (m, 4 H), 7.30-7.50 (m, 5 H), 4.71 (d, 2 H).

Example 109 - Preparation of Compound 126

To a stirred solution of 3-iodobenzylamine (5.00 g, 21.4 mmol) in water (100 mL) was added 2,6-dichloropurine (4.04 g, 21.4 mmol) and *N,N*-diisopropylethylamine (7.47 mL, 42.5 mmol). The mixture was refluxed for 5 h and stored at room temperature overnight. The resulting suspension was filtered. The filter cake was triturated with water (3 x 25 mL) and ethanol (2 x 15 mL) and dried under high vacuum to yield **126** (7.49 g, 91%): ^1H NMR (300 MHz, $(\text{CD}_3)_2\text{SO}$) δ 8.50-8.80 (brs, 1 H), 8.17 (s, 1 H), 7.75 (s, 1 H), 7.61 (d, 1 H), 7.37 (d, 1 H), 7.14 (t, 1 H), 5.14 (brs, 1 H), 4.61 (d, 2 H).

Example 110 - Preparation of Compound 127

The purine derivative **126** (7.00 g, 18.2 mmol) was dissolved in dimethylsulfoxide (120 mL). To this stirred solution was added potassium carbonate (20.0 g, 145 mmol) and 2-iodopropane (7.28 mL, 72.8 mmol). The reaction mixture was stirred under a nitrogen atmosphere for 20 h before being poured into stirred water (600 mL). After 10 min, the resulting mixture was extracted with ethyl acetate (4 x 95 mL). The combined organic layers were washed with water (25 mL) and brine (3 x 25 mL), dried over sodium sulfate, and concentrated in vacuo. The resulting material was purified by recrystallization from ethyl acetate in hexanes to yield **127** (7.51 g, 97%): mp 147-152 °C; ^1H NMR (300 MHz, $(\text{CD}_3)_2\text{SO}$) δ 8.84 (m, 1 H), 8.31 (s, 1 H), 7.74 (s, 1 H), 7.60 (d, 1 H), 7.36 (d, 1 H), 7.13 (t, 1 H), 4.51-4.75 (m, 3 H), 1.50 (d, 6 H); API MS m/z = 429 $[\text{C}_{15}\text{H}_{15}\text{ClIN}_5+\text{H}]^+$.

Example 111 - Preparation of Compound 128

In a sealed tube, 127 (2.57 g, 6.00 mmol), *trans*-1,4-diaminocyclohexane (6.85 g, 60.0 mmol), and ethanol (10 mL) were combined. The reaction mixture was heated to 160 °C for 24 h. After cooling to room temperature, the mixture was filtered. The filtrate was concentrated and diluted with ethyl acetate (250 mL). The organic solution was washed with water (250 mL) and saturated sodium bicarbonate solution (2 x 250 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. A portion of the resulting crude product was purified via silica gel chromatography to yield 128 (181 mg): ¹H NMR (300 MHz, (CD₃)₂SO) δ 7.87 (brs, 1 H), 7.78 (s, 1 H), 7.70 (s, 1 H), 7.55 (d, 1 H), 7.35 (d, 1 H), 7.09 (t, 1 H), 6.06 (d, 1 H), 4.43-4.70 (m, 3 H), 3.58 (brs, 1 H), 1.66-1.94 (m, 4 H), 1.46 (d, 6 H), 1.00-1.30 (m, 4 H); ESI MS *m/z* = 506 [C₂₁H₂₈IN₇+H]⁺.

Example 112 - Preparation of Compound 129

The amine 127 (6.06 g, 6.00 mmol) was dissolved in a mixture of tetrahydrofuran (45 mL) and water (15 mL). To this stirred mixture were added sodium bicarbonate (2.02 g, 24.0 mmol) and di-*tert*-butyldicarbonate (2.90 g, 13.3 mmol). After 3.5 h, the solution was extracted with methylene chloride (3 x 75 mL). The organic extracts were combined, washed with brine (225 mL), and dried over sodium sulfate. The organic liquid was concentrated and the resulting material was purified via silica gel chromatography (33:67 to 50:50 to 60:40 ethyl acetate/hexanes) to yield 129 (4.94 g, 68%): ¹H NMR (300 MHz, CDCl₃) δ 7.72 (s, 1 H), 7.58 (d, 1 H), 7.46 (s, 1 H), 7.30 (d, 1 H), 7.02 (t, 1 H), 4.50-4.76 (m, 4 H), 4.40 (m, 1 H), 3.70 (brs, 1 H), 3.43 (brs, 1 H), 1.90-2.20 (m, 4 H), 1.51 (d, 6 H), 1.45 (s, 9 H), 1.13-1.35 (m, 4 H); API MS *m/z* = 607 [C₂₆H₃₆IN₇O₂+H]⁺.

Example 113 - Preparation of Compound 130

To a stirred solution of 129 (1.00 g, 1.65 mmol) in ethylene glycol dimethyl ether (40 mL) was added 3-thiopheneboronic acid, triphenylphosphine (250 mg, 0.950 mmol), and 2 M sodium carbonate solution (3.8 mL). The mixture was
5 purged with nitrogen for 10 min and tris(dibenzylideneacetone)dipalladium(0) (64.0 mg, 0.060 mmol) was added. After refluxing overnight under nitrogen, the reaction mixture was cooled to room temperature and diluted with water (100 mL). The resulting solution was extracted with methylene chloride (3 x 50 mL). The organic
10 extracts were combined, washed with brine (30 mL), and dried over sodium sulfate. The organic liquid was filtered and concentrated in vacuo. Purification via silica gel chromatography (50:50 ethyl acetate/hexanes) yielded 130 (0.87 g, 94%): ¹H NMR (300 MHz, CDCl₃) δ 7.23-7.30 (m, 8 H), 6.02 (brs, 1 H), 4.78 (d, 2 H), 4.53-4.68 (m, 1 H), 4.32 (m, 1 H), 3.70 (m, 1 H), 3.28-3.55 (m, 1 H), 1.80-2.19 (m, 4 H), 1.53 (d, 6
15 H), 1.45 (s, 9 H), 1.05-1.32 (m, 4 H); ESI MS *m/z* = 562 [C₃₀H₃₉N₇O₂S+H]⁺.

Example 114 - Preparation of Compound 131

To a solution of 130 in EtOAc was added 1 N HCl. After concentration
20 of the solution, isolate 131 (658 mg, 91%): mp 211-216 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.40 (brs, 1 H), 7.25-7.85 (m, 7 H), 4.60-5.10 (m, 5 H), 3.91 (m, 1 H), 3.16 (m, 1 H), 1.94-2.33 (m, 4 H), 1.32-1.79 (m, 10 H); ESI MS *m/z* = 462 [C₂₅H₃₁N₇S+H]⁺.

Example 115 - Preparation of Compound 132

Following procedures outlined above for acetylation, prepared 132
from 131 (352 mg, 80%): mp 209-211 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.27-7.67
(m, 8 H), 6.25 (brs, 1 H), 5.19 (d, 1 H), 4.79 (d, 2 H), 4.52-4.67 (m, 2 H), 3.69 (m, 2
30 H), 2.09 (m, 2 H), 1.88-1.98 (m, 5 H), 1.50 (d, 6 H), 1.06-1.33 (m, 4 H); ESI MS *m/z*
= 504 [C₂₇H₃₃N₇OS+H]⁺.

Example 116 - Preparation of Compound 133

Following general procedures outlined above for Suzuki Coupling reaction, prepared 133 from 129 (0.61 g, 67%): ^1H NMR (300 MHz, CDCl_3) δ 7.30-7.62 (m, 10 H), 6.00 (brs, 1 H), 4.83 (d, 2 H), 4.54-4.70 (m, 2 H), 4.37 (m, 1 H), 3.71 (m, 1 H), 3.39 (m, 1 H), 2.12 (m, 2 H), 2.47 (m, 2 H), 1.53 (d, 6 H), 1.45 (s, 9 H), 1.19 (m, 4 H); ESI MS m/z = 556 $[\text{C}_{32}\text{H}_{41}\text{N}_7\text{O}_2+\text{H}]^+$.

Example 117 - Preparation of Compound 134

To a stirred solution of 133 (530 mg, 0.950 mmol) in methanol (3 mL) was added 1 N HCl in diethyl ether (9.50 mL, 9.50 mmol). After stirring for 3.5 h, hydrogen chloride gas was gently bubbled through the solution. After 20 min, the solution was concentrated in vacuo. The resulting material was recrystallized from methanol in ether to afford 134 in quantitative yield: ^1H NMR (300 MHz, CD_3OD) δ 8.33 (brs, 1 H), 7.67 (s, 1 H), 7.52-7.64 (m, 3 H), 7.29-7.50 (m, 5 H), 4.80-5.00 (m, 4 H), 4.72 (m, 1 H), 3.87 (m, 1 H), 3.14 (m, 1 H), 2.02-2.25 (m, 4 H), 1.59 (d, 6 H), 1.47 (m, 4 H); ESI MS m/z = 456 $[\text{C}_{27}\text{H}_{33}\text{N}_7+\text{H}]^+$.

Example 118 - Preparation of Compound 135

Following procedures outlined above for acetylation, prepared 135 from 134 (195 mg, 82%): mp 183-185 °C; ^1H NMR (300 MHz, CDCl_3) δ 7.30-7.64 (m, 10 H), 6.19 (brs, 1 H), 5.12 (d, 1 H), 4.81 (d, 2 H), 4.60 (m, 2 H), 3.68 (m, 2 H), 2.09 (m, 2 H), 1.86-1.99 (m, 5 H), 1.51 (d, 5 H), 1.04-1.32 (m, 4 H); ESI MS m/z = 498 $[\text{C}_{29}\text{H}_{35}\text{N}_7\text{O}+\text{H}]^+$.

Example 119 - Preparation of Compound 136

To a mixture of 71 (2.00 g, 5.96 mmol) in dimethylsulfoxide (44 mL) was added potassium carbonate (6.56 g, 47.7 mmol) and iodoethane (2.00 mL, 24.4 mmol). After stirring overnight, the reaction mixture was poured into a stirred solution of water (300 mL). After 2 d, it was filtered. The filtrate was extracted with ethyl acetate (2 x 180 mL). The organic extracts were combined and washed with

brine (150 mL). The organic layer was dried over magnesium sulfate. Concentration afforded 136 (1.90 g, 88%).

Example 120 - Preparation of Compound 137

5 In a sealed tube, 136 (0.60 g, 1.66 mmol), *trans*-1,4-diaminocyclohexane (1.93 g, 16.8 mmol), and potassium iodide (10 mg), were dissolved in ethanol (18 mL). The mixture was heated to 160 °C. After 4 d, the mixture was cooled to room temperature and filtered. The filtrate was dissolved in
10 ethyl acetate and washed with water (2 x 100 mL) and brine (100 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. The material was purified by silica gel chromatography and recrystallization from ethanol in hexanes (1:10) to yield 137 (886 mg, 41%): mp 175-182 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.30-7.75 (m, 10 H), 4.55-4.95 (m, 4 H), 4.10 (q, 2 H), 3.78 (m, 1 H), 2.99 (m, 1 H),
15 2.11 (d, 2 H), 1.99 (d, 2 H), 1.44 (m, 3 H), 1.31 (m 4 H); ESI MS *m/z* = 442 [C₂₆H₃₁N₇+H]⁺.

Example 121 - Preparation of Compound 138

20 Compound 71 (2.02 g, 6.02 mmol), iodomethane (1.50 mL, 24.4 mmol), and potassium carbonate were dissolved in dimethylsulfoxide (44 mL) and stirred overnight. The reaction mixture was poured into 150 mL of stirring water. The organic and aqueous layers were separated. The organic layer was washed with brine (3 x 100 mL) and dried over magnesium sulfate. The solids were removed by
25 filtration and the solution was concentrated in vacuo to afford 138 (1.93 g, 93%).

Example 122 - Preparation of Compound 139

In a sealed tube, 138 (1.80 g, 5.15 mmol) and *trans*-1,4-
30 diaminocyclohexane (5.90 g, 51.7 mmol) were dissolved in ethanol (85 mL). The mixture was heated to 140 °C. After heating overnight, the reaction mixture was cooled to room temperature and concentrated. The resulting solid was dissolved in ethyl acetate (160 mL) and washed with water (160 mL) and brine (2 x 100 mL). The organic layer was dried over magnesium sulfate and filtered. Concentration afforded a
35 solid, which was purified by silica gel chromatography and recrystallization from

- 176 -

ethanol in hexanes (1:20) and ethyl acetate in hexanes to yield **139** (850 mg, 38%): mp 182-184 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.25-7.70 (m, 10 H), 4.64-4.90 (m, 4 H), 3.75 (m, 1 H), 3.65 (s, 3 H), 2.68 (m, 1 H), 2.05 (m, 2 H), 1.88 (m, 2 H), 1.25 (m, 4 H); ESI MS *m/z* = 428 [C₂₅H₂₉N₇+H]⁺.

5

Example 123 - Preparation of Compound 140

The HCl salt of **139** (86.7 mg, 0.162 mmol) was suspended in methylene chloride (20 mL). The suspension was immersed in an ice bath while triethylamine (0.16 mL, 1.12 mmol) and a catalytic amount of DMAP were added. Acetyl chloride (0.04 mL, 0.560 mmol) was added to the mixture. The reaction was quenched by the addition of 5% aqueous NaHCO₃ solution (50 mL). The aqueous layer was extracted with methylene chloride (2 x 50 mL). The extracts were washed with brine (100 mL), dried over magnesium sulfate, filtered, and concentrated. The resulting solid material was dried in vacuo. Purification by silica gel chromatography and recrystallization from ethyl acetate in hexanes (5:60) afforded **140** (6.4 mg, 8%): ¹H NMR (300 MHz, CD₃OD) δ 7.50-7.68 (m, 5 H), 7.25-7.47 (m, 5 H), 5.22 (m, 3 H), 4.20-5.05 (m, 2 H), 3.75 (m, 1 H), 3.65 (s, 3 H), 3.35-3.47 (m, 1 H), 2.78 (m, 1 H), 1.95-2.16 (m, 2 H), 1.80-1.95 (m, 2 H), 1.17-1.40 (m, 4 H).

20

Example 124 - Preparation of Compound 144

Compound **71** (2.03 g, 5.96 mmol), 1-iodopropane (2.25 mL, 24.4 mmol), and potassium carbonate (6.61 g, 47.7 mmol) were dissolved in dimethylsulfoxide (44 mL) and allowed to stir overnight. The reaction mixture was added to 300 mL of stirring water and stirred for 2 d. The resulting precipitate was collected by filtration and dried in vacuo to afford **141** (2.07 g, 92%).

25

Example 125 - Preparation of Compound 142

30

In a sealed tube, **141** (1.82 g, 4.81 mmol) and *trans*-1,4-diaminocyclohexane (5.67 g, 49.7 mmol) were dissolved in ethanol (53 mL). The mixture was heated to 140 °C for 3 d. After cooling to room temperature, the reaction mixture was concentrated and dissolved in ethyl acetate (100 mL). This solution was

- 177 -

washed with water (100 mL) and brine (2 x 100 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography and recrystallizations from ethanol in hexanes (1:20) to yield **142** (523 mg, 24%): mp 133-138 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.25-7.70 (m, 10 H), 4.65-4.85 (m, 4 H), 4.02 (t, 2 H), 3.76 (m, 1 H), 2.85 (m, 1 H), 2.08 (d, 2 H), 1.94 (d, 2 H), 1.86 (q, 2 H), 1.20-1.42 (m, 4 H), 0.93 (t, 3 H); ESI MS *m/z* = 456 [C₂₇H₃₃N₇+H]⁺.

Example 126 - Preparation of Compound 143

Compound **71** (2.01 g, 5.98 mmol), iodocyclopentane (2.80 mL, 24.2 mmol), and potassium carbonate (6.75 g, 48.9 mmol) were dissolved in dimethylsulfoxide (44 mL) and allowed to stir under nitrogen overnight. The reaction mixture was added to 150 mL of stirring water and diluted with 150 mL ethyl acetate. The organic and aqueous phases were separated. The organic phase was washed with brine (3 x 100 mL) and dried over magnesium sulfate. After filtering, the organic liquid was concentrated and the resulting solid was dried in vacuo to afford **143** (1.29 g, 55%).

Example 127 - Preparation of Compound 144

In a sealed tube, **143** (304 mg, 0.749 mmol) and *trans*-1,4-diaminocyclohexane (891 mg, 7.80 mmol) were dissolved in ethanol (10 mL). The mixture was heated to 140 °C for 4 d. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate (160 mL). This solution was washed with water (160 mL) and brine (2 x 100 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated. The material was purified by silica gel chromatography and recrystallizations from ethanol in hexanes (1:20) to yield **144** (299 mg, 19%): mp 144-146 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.28-7.75 (m, 10 H), 4.68-4.85 (m, 4 H), 3.75 (m, 1 H), 2.82 (m, 1 H), 1.70-2.25 (m, 13 H), 1.20-1.43 (m, 4 H); ESI MS *m/z* = 482 [C₂₉H₃₅N₇+H]⁺.

Example 128 - Preparation of Compound 145

Compound 71 (2.01 g, 5.98 mmol), allylbromide (2.10 mL, 24.4 mmol), and potassium carbonate (6.61 g, 47.8 mmol) were dissolved in dimethylsulfoxide (44 mL) and stirred overnight. The reaction mixture was added to 150 mL of stirring water and diluted with 150 mL ethyl acetate. The organic and aqueous phases were separated. The organic phase was washed with brine (3 x 100 mL), dried over magnesium sulfate, filtered, and concentrated. The resulting solid was dried in vacuo to afford 145 (1.98 g, 88%).

Example 129 - Preparation of Compound 146

In a sealed tube, 145 (1.99 mg, 5.29 mmol), *trans*-1,4-diaminocyclohexane (6.21 g, 54.3 mmol), and 2,6-di-*tert*-butylphenol (1.13 g, 5.48 mmol) were dissolved in ethanol (60 mL). The mixture was heated to 140 °C for 4 d. After cooling to room temperature, the reaction was concentrated and diluted with ethyl acetate (175 mL). This organic solution was washed with water (175 mL) and brine (2 x 100 mL) and concentrated. The product was purified by silica gel chromatography and recrystallizations from ethanol in hexanes (1:20) to yield 146 (432 mg, 18%): mp 111-114 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.25-7.72 (m, 10 H), 5.95-6.10 (m, 1 H), 5.21 (d, 1 H), 5.10 (d, 1 H), 4.82 (m, 2 H), 4.77 (s, 2 H), 4.67 (d, 2 H), 3.75 (m, 1 H), 2.87 (m, 1 H), 2.07 (d, 2 H), 1.92 (d, 2 H), 1.15-1.47 (m, 4 H); ESI MS *m/z* = 454 [C₂₇H₃₁N₇+H]⁺.

Example 130 - Preparation of Compound 147

Compound 71 (2.07 g, 6.17 mmol), 2-iodobutane (3.10 mL, 26.9 mmol), and potassium carbonate (6.78 g, 49.1 mmol) were dissolved in dimethylsulfoxide (44 mL) and allowed to stir under nitrogen overnight. The reaction mixture was diluted with ethyl acetate (300 mL). The organic material was washed with water (200 mL) and brine (300 mL) and dried over magnesium sulfate. After filtration, the material was concentrated and the resulting solid was dried in vacuo to afford 147 (1.29 g, 55%).

Example 131 - Preparation of Compound 148

In a sealed tube, **147** (1.29 g, 3.30 mmol) and *trans*-1,4-diaminocyclohexane (3.80 g, 33.2 mmol) were dissolved in ethanol (70 mL). The mixture was heated to 140 °C. After 4 d, the reaction was cooled to room temperature, concentrated, and dissolved in ethyl acetate (160 mL). This solution was washed with water (160 mL) and brine (2 x 100 mL) and dried over magnesium sulfate. The organic liquid was filtered and concentrated. The resulting solid was dried in vacuo and purified by silica gel chromatography and recrystallizations from ethanol in hexanes (1:40) to yield **148** (229 mg, 15%): mp 146-150 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.70 (s, 1 H), 7.25-7.60 (m, 9 H), 4.65-4.90 (m, 4 H), 4.37 (m, 1 H), 3.74 (m, 1 H), 2.86 (m, 1 H), 1.82-2.15 (m, 6 H), 1.50 (d, 3 H), 1.20-1.43 (m, 4 H), 0.87 (t, 3 H).

Example 132 - Preparation of Compound 149

To a solution of **61** (0.90 g, 1.96 mmol) in ethylene glycol dimethyl ether (54 mL), were added 3,5-dimethylphenylboronic acid (0.59 g, 3.93 mmol), triphenylphosphine (0.26 g, 0.99 mmol), and 2 M sodium carbonate solution (10 mL). The solution was refluxed for 20 min and cooled to room temperature. Tris(dibenzylideneacetone)dipalladium(0) (0.66 g, 0.072 mmol) was added and the reaction returned to reflux. After refluxing overnight, the reaction mixture was cooled to room temperature and another 100 mg 3,5-dimethylphenylboronic acid were added. After refluxing for another 5 h, the reaction was quenched with 50 mL water. The aqueous solution was extracted with methylene chloride (3 x 50 mL). The extracts were combined and washed with water (50 mL) and brine (50 mL). The organic solution was dried over sodium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography to yield **149**: mp 86-90 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.34-7.58 (m, 5 H), 7.20 (s, 2 H), 6.97 (s, 1 H), 5.93 (brs, 1 H), 4.54-4.90 (m, 4 H), 3.66-3.85 (m, 1 H), 2.70 (m, 1 H), 2.37 (s, 6 H), 2.05-2.20 (m, 4 H), 1.80-1.95 (m, 2 H), 1.54 (d, 6 H), 1.10-1.35 (m, 4 H); API MS *m/z* = 484 [C₂₉H₃₇N₇+H]⁺.

Example 133 - Preparation of Compound 150

To a stirred, 0 °C solution of 149 (500 mg, 0.97 mmol) in methylene chloride (20 mL), were added pyridine (120 µL), DMAP (11.8 mg, 0.097 mmol), and acetic anhydride (91 µL, 0.097 mmol). After 40 min, the reaction mixture was warmed to room temperature and stirred for 3 h. Another 100 µL acetic anhydride was added. After 1 h, the solution was concentrated and dried in vacuo. The resulting material was purified by silica gel chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to yield 150 (400 mg, 80%): mp 207-210 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.46-7.56 (m, 3 H), 7.36-7.45 (m, 2 H), 7.18 (s, 2 H), 6.98 (s, 1 H), 5.98 (brs, 1 H), 5.27 (d, 1 H), 4.80 (d, 2 H), 4.56-4.70 (m, 1 H), 3.68-3.84 (m, 1 H), 2.37 (s, 6 H), 1.90-2.23 (m, 7 H), 1.54 (d, 6 H), 1.15-1.48 (m, 4 H); API MS *m/z* = 526 [C₃₁H₃₉N₇O+H]⁺.

Example 134 - Preparation of Compound 152

To a solution of 149 (500 mg, 1.03 mmol) in 1,2-dichloroethane (4 mL) was added propionaldehyde (90 µL, 1.24 mmol). After stirring for 10 min, sodium triacetoxyborohydride (306 mg, 1.44 mmol) was added. The reaction mixture stirred under nitrogen for 1.5 h before being quenched with saturated sodium bicarbonate solution (5 mL). The resulting solution was extracted with ethyl acetate (3 x 7 mL). The organic extracts were combined and dried over sodium sulfate. The organic liquid was filtered and concentrated. Purification by silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) yielded 152.

Example 135 - Preparation of Compound 152 ·HCl

To a stirred solution of 152 (120 mg, 0.211 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (127 µL). After 20 min, the solution was concentrated and dried in vacuo to yield the HCl salt of 152: ¹H NMR (300 MHz, CD₃OD) δ 8.48 (s, 1 H), 7.40-7.68 (m, 4 H), 7.20 (s, 2 H), 6.98 (s, 1 H), 4.66-5.07 (m, 3 H), 3.80-4.00 (m, 1 H), 2.90-3.50 (m, 5 H), 2.01-2.45 (m 10 H), 1.36-1.90 (m, 14 H), 1.02 (t, 6 H); API MS *m/z* = 568 [C₃₅H₄₉N₇+H]⁺.

Example 136 - Preparation of Compound 151

To a solution of 149 (302 mg, 0.624 mmol) in 1,2-dichloroethane (2.5 mL) was added propionaldehyde (36.0 μ L, 0.500 mmol). After stirring for 15 min under nitrogen, sodium triacetoxymethylborohydride (93.0 mg, 0.874 mmol) was added. After 10 min, another 93.0 mg (0.874 mmol) sodium triacetoxymethylborohydride were added. The reaction mixture stirred under nitrogen for 1.5 h before being quenched with saturated sodium bicarbonate solution (5 mL). The resulting solution was extracted with ethyl acetate (3 x 7 mL). The organic extracts were combined and dried over sodium sulfate. The organic liquid was filtered and concentrated. Purification by silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) yielded 151.

Example 137 - Preparation of Compound 151 ·HCl

To a stirred solution of 151 (90.0 mg, 0.171 mmol) in ethyl acetate (8 mL) was added 2 M HCl in diethyl ether (103 μ L). After 20 min, the solution was concentrated and dried in vacuo to yield the HCl salt of 151: mp 280-290 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.45 (brs, 1 H), 7.38-7.67 (m, 4 H), 7.29 (s, 2 H), 6.97 (s, 1 H), 4.63-5.40 (m, 4 H), 3.78-3.97 (m, 1 H), 2.85-3.23 (m, 3 H), 2.07-2.44 (m, 8 H), 1.32-1.87 (m, 14 H), 1.02 (t, 3 H); API MS m/z = 526 [C₃₂H₄₃N₇+H]⁺.

Example 138 - Preparation of Compound 153

To a solution of 61 (2.03 g, 4.44 mmol) in ethylene glycol dimethyl ether (100 mL), was added 2,5-dimethoxyphenylboronic acid (2.42 g, 13.3 mmol), tris(dibenzylideneacetone)dipalladium(0) (0.135 g, 0.148 mmol), triphenylphosphine (0.581 g, 2.22 mmol), and 2 M sodium carbonate solution (10 mL). The dispersion was refluxed overnight under nitrogen. After cooling to room temperature, the reaction mixture was diluted with 100 mL water. The aqueous solution was extracted with methylene chloride (3 x 100 mL). The extracts were combined and washed with water (300 mL) and brine (300 mL). The organic solution was dried over sodium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) to yield 153 (900 mg, 39%).

Example 139 - Preparation of 153 ·HCl

To a stirred solution of **153** (100 mg, 0.194 mmol) in ethyl acetate (10 mL), was added 2 M HCl in diethyl ether (116 µL). After 20 min, the solution was concentrated to afford the HCl salt of **153** in quantitative yield: mp 278-288 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.34 (brs, 1 H), 7.36-7.53 (m, 4 H), 6.77-7.01 (m, 3 H), 5.30 (brs, 1 H), 4.65-5.20 (m, 4 H), 3.80-3.95 (m, 1 H), 3.76 (s, 1 H), 3.69 (s, 1 H), 3.05-3.22 (m, 1 H), 2.03-2.30 (m, 4 H), 1.35-1.71 (m, 10 H); API MS *m/z* = 516 [C₂₉H₃₇N₇O₂+H]⁺.

Example 140 - Preparation of Compound 154

To a 0 °C stirred solution of **153** (300 mg, 0.582 mmol) in methylene chloride (12 mL) was added pyridine (70 µL), acetic anhydride (54.0 µL, 0.582 mmol), and DMAP (7.10 mg, 0.582 mmol). The solution was stirred for 40 min before warming to room temperature. After 3 h, another 100 µL acetic anhydride was added. After stirring for 1 h, the reaction mixture was concentrated and dried in vacuo. The material was purified by silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) to yield **154**: mp 185-192 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.78 (s, 1 H), 7.29-7.45 (m, 4 H), 6.87-6.98 (m, 1 H), 6.75-6.86 (m, 2 H), 4.75 (s, 2 H), 4.55-4.68 (m, 1 H), 3.74 (s, 3 H), 3.67 (s, 3 H), 1.97-2.13 (m, 2 H), 1.81-1.96 (m, 5 H), 1.53 (d, 6 H), 1.21-1.40 (m, 4 H); API MS *m/z* = 558 [C₃₁H₃₉N₇O₃+H]⁺.

Example 141 - Preparation of Compound 155

To a solution of **153** (242 mg, 0.469 mmol) in 1,2-dichloroethane (1.5 mL) was added propionaldehyde (27.0 µL, 0.375 mmol). After stirring for 15 min under nitrogen, sodium triacetoxyborohydride (140 mg, 0.657 mmol) was added. The reaction mixture was stirred under nitrogen overnight before being concentrated. Purification by silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) yielded **155**.

Example 142 - Preparation of Compound 155 ·HCl

To a stirred solution of 155 (30.0 mg, 0.054 mmol) in ethyl acetate (6 mL) was added 2 M HCl in diethyl ether (40.0 µL). After 20 min, the solution was concentrated and dried in vacuo to afford the HCl salt of 155 in quantitative yield: mp 264-268 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.41 (brs, 1 H), 7.46 (m, 4 H), 6.98 (d, 1 H), 6.76-6.93 (m, 2 H), 4.63-5.07 (m, 4 H), 3.89 (m, 1 H), 3.77 (s, 3 H), 3.71 (s, 3 H), 3.13 (m, 1 H), 2.97 (m, 2 H), 2.10-2.35 (m, 4 H), 1.35-1.88 (m, 12 H), 1.02 (t, 3 H); API MS *m/z* = 558 [C₃₂H₄₃N₇O₂+H]⁺.

Example 143 - Preparation of Compound 156

To a solution of 153 (242 mg, 0.469 mmol) in 1,2-dichloroethane (1.5 mL) was added propionaldehyde (27.0 µL, 0.375 mmol). After stirring for 15 min under nitrogen, sodium triacetoxyborohydride (140 mg, 0.657 mmol) was added. The reaction mixture stirred under nitrogen overnight before being concentrated. Purification by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) yielded 156.

Example 144 - Preparation of Compound 156 ·HCl

To a stirred solution of 156 (160 mg, 0.267 mmol) in ethyl acetate (8 mL) was added 2 M HCl in diethyl ether (170 µL). After 20 min, the solution was concentrated and dried in vacuo to afford the HCl salt of 156 in quantitative yield: mp 235-243 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.43 (brs, 1 H), 7.47 (m, 4 H), 6.94-7.04 (d, 1 H), 6.80-6.93 (m, 2 H), 4.67-5.10 (m, 4 H), 3.81-3.98 (m, 1 H), 3.76 (s, 3 H), 3.71 (s, 3 H), 2.93-3.48 (m, 5 H), 2.00-2.35 (m, 4 H), 1.36-1.90 (m, 14 H), 1.01 (t, 6 H); API MS *m/z* = 601 [C₃₅H₄₉N₇O₂+H]⁺.

Example 145 - Preparation of Compound 157

To a solution of 61 (1.50 g, 3.27 mmol) in ethylene glycol dimethyl ether (75 mL), was added 5-methyl-2-thiopheneboronic acid (1.40 g, 9.82 mmol), tris(dibenzylideneacetone)dipalladium(0) (100 mg, 0.109 mmol), triphenylphosphine (430 mg, 1.64 mmol), and 2 M sodium carbonate solution (10 mL). The solution was

refluxed under nitrogen for 2 d. After cooling to room temperature, the reaction mixture was diluted with 100 mL water. The aqueous solution was extracted with methylene chloride (3 x 100 mL). The extracts were combined and washed with water (300 mL) and brine (300 mL). The organic solution was dried over sodium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) to yield 157 (1.04 g, 67%).

Example 146 - Preparation of Compound 157 ·HCl

To a stirred solution of 157 (150 mg, 0.269 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (161 µL). After 20 min, the solution was concentrated to afford the HCl salt of 157 in quantitative yield: mp 300 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.35 (brs, 1 H), 7.55 (d, 2 H), 7.39 (d 2 H), 7.17 (d, 1 H), 6.74 (d, 1 H), 4.62-5.40 (m, 5 H), 3.78-3.94 (m, 1 H), 3.03-3.20 (m, 1 H), 2.47 (s, 3 H), 1.98-2.29 (m, 4 H), 1.44-1.74 (m, 10 H); API MS *m/z* = 476 [C₂₆H₃₃N₇S+H]⁺.

Example 147 - Preparation of Compound 159

To a solution of 157 (300 mg, 0.631 mmol) in 1,2-dichloroethane (2.0 mL) was added propionaldehyde (36.0 µL, 0.505 mmol). After stirring for 30 min under nitrogen, sodium triacetoxyborohydride (190 mg, 0.883 mmol) was added. The reaction mixture stirred under nitrogen for 3 h before being concentrated. Purification by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) yielded 159: mp 150-155 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.81 (s, 1 H), 7.51 (d, 2 H), 7.33 (d, 2 H), 7.11 (d, 1 H), 6.73 (d, 1 H), 4.80-5.05 (m, 1 H), 4.55-4.78 (m, 3 H), 3.65-3.80 (m, 1 H), 3.01 (m, 1 H), 2.91 (t, 2 H), 2.47 (s, 3 H), 2.02-2.19 (m, 4 H), 1.60-1.75 (m, 8 H), 1.54 (d, 6 H), 1.19-1.50 (m, 4 H), 1.01 (t, 3 H); API MS *m/z* = 518 [C₂₉H₃₉N₇S+H]⁺.

Example 148 - Preparation of Compound 158

To a 0 °C stirred solution of 157 (300 mg, 0.631 mmol) in methylene chloride (12 mL) was added pyridine (76 µL), acetic anhydride (58 µL, 0.631 mmol), and DMAP (7.7 mg, 0.063 mmol). After 20 min, the reaction mixture warmed to

room temperature. After mixing overnight, the reaction mixture was concentrated and dried in vacuo. The material was purified by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) to yield **158**: mp 225-230 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.39-7.51 (m, 3 H), 7.32 (d, 2 H), 7.07 (d, 1 H), 6.71 (m, 1 H), 6.14 (brs, 1 H), 5.32 (d, 1 H), 4.53-4.82 (m, 4 H), 3.74 (m, 2 H), 2.50 (s, 3 H), 1.90-2.23 (m, 7 H), 1.51 (d, 6 H), 1.12-1.38 (m, 4 H); API MS *m/z* = 518 [C₂₈H₃₅N₇OS+H]⁺.

Example 149 - Preparation of Compound 160

To a suspension of **61** (1.00 g, 2.18 mmol) in ethylene glycol dimethyl ether (50 mL), was added 4-methylthiophene-2-boronic acid (0.93 g, 6.54 mmol), tris(dibenzylideneacetone)dipalladium(0) (67.0 mg, 0.073 mmol), triphenylphosphine (287 mg, 1.09 mmol), and 2 M sodium carbonate solution (10 mL). The solution was refluxed under nitrogen for 2 d. After cooling to room temperature, the reaction mixture was diluted with 100 mL water. The aqueous solution was extracted with methylene chloride (3 x 100 mL). The extracts were combined and washed with water (300 mL) and brine (300 mL). The organic solution was dried over sodium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) to yield **160** (450 mg, 44%).

Example 150 - Preparation of Compound 160 ·HCl

To a stirred solution of **160** (50.0 mg, 0.105 mmol) in ethyl acetate (4 mL) was added 2 M HCl in diethyl ether (100 µL). After 20 min, the solution was concentrated to afford the HCl salt of **160** in quantitative yield: mp 308-315 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.31 (brs, 1 H), 7.60 (d, 2 H), 7.39 (d, 2 H), 7.20 (s, 1 H), 6.94 (s, 1 H), 4.65-5.10 (m, 3 H), 3.86 (m, 1 H), 3.13 (m, 1 H), 2.26 (s, 3 H), 2.00-2.12 (m, 4 H), 1.32-1.72 (m, 10 H); API MS *m/z* = 476 [C₂₆H₃₃N₇S+H]⁺.

Example 151 - Preparation of Compounds 162 and 163

To a stirred solution of **160** (300 mg, 0.631 mmol) in 1,2-dichloromethane (3 mL) was added propionaldehyde (36.0 µL). After stirring under nitrogen for 15 min, sodium triacetoxyborohydride (161 mg, 0.757 mmol) was added. The mixture was stirred overnight. The reaction was concentrated and purified by

silica gel chromatography (95:4.5:1 CHCl₃/methanol/NH₄OH) to yield **162** (80 mg, 31%), and **163** (110 mg). For **162**: ¹H NMR (300 MHz, CD₃OD) δ 7.79 (s, 1 H), 7.54 (d, 2 H), 7.33 (d, 2 H), 7.16 (s, 1 H), 6.89 (s, 1 H), 4.71 (s, 2 H), 4.60 (m, 1 H), 3.70 (m, 1 H), 3.31 (t, 2 H), 2.42-2.62 (m, 3 H), 2.23 (s, 3 H), 1.87-2.10 (m, 4 H), 1.51 (d, 6 H), 1.10-1.33 (m, 4 H), 0.94 (t, 3 H).

Example 152 - Preparation of Compound 162 ·HCl

To a stirred solution of **162** (80.0 mg, 0.155 mmol) in ethyl acetate (6 mL) was added 2 M HCl in diethyl ether (100 μL). After 20 min, the solution was concentrated to afford the HCl salt of **162** in quantitative yield: mp 225-240 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.36 (brs, 1 H), 7.62 (d, 2 H), 7.42 (d, 2 H), 7.22 (s, 1 H), 6.95 (s, 1 H), 4.65-5.05 (m, 4 H), 3.87 (m, 1 H), 3.11 (m, 1 H), 2.98 (t, 2 H), 2.07-2.36 (m, 5 H), 1.32-1.85 (m, 14 H), 1.04 (t, 3 H); API MS *m/z* = 518 [C₂₉H₃₉N₇S+H]⁺.

Example 153 - Preparation of Compound 163 ·HCl

To a stirred solution of **163** (110 mg, 0.196 mmol) in ethyl acetate (8 mL) was added 2 M HCl in diethyl ether (120 μL). After 20 min, the solution was concentrated to afford the HCl salt of **163** in quantitative yield: mp 227-229 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.39 (brs, 1 H), 7.60 (d, 2 H), 7.41 (d, 2 H), 7.22 (s, 1 H), 6.94 (s, 1 H), 4.63-5.10 (m, 3 H), 3.85 (m, 1 H), 2.93-3.47 (m, 5 H), 1.92-2.40 (m, 7 H), 1.32-1.90 (m, 14 H), 1.04 (t, 6 H); API MS *m/z* = 560 [C₃₂H₄₅N₇S+H]⁺.

Example 154 - Preparation of Compound 161

To a 0 °C stirred solution of **160** (100 mg, 0.210 mmol) in methylene chloride (5 mL) were added pyridine (26.0 μL), acetic anhydride (20.0 μL, 0.210 mmol), and DMAP (3.0 mg, 0.021 mmol). After 40 min, the reaction mixture warmed to room temperature. The reaction mixture was concentrated and dried in vacuo. The material was purified by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) to yield **161**: mp 222-223 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.78 (s, 1 H), 7.52 (d, 2 H), 7.35 (d, 2 H), 7.16 (s, 1 H), 6.89 (s, 1 H), 4.83-5.05 (m, 1 H), 4.72 (s, 2 H), 4.64 (m, 1 H), 3.73 (m, 1 H), 3.61 (m, 1 H), 2.25 (s, 3 H),

- 187 -

1.97-2.13 (m, 2 H), 1.81-1.96 (m, 5 H), 1.54 (d, 6 H), 1.19-1.40 (m, 4 H); API MS m/z = 518 [$C_{28}H_{35}N_7OS+H$]⁺.

Example 155 - Preparation of Compound 164

5 To a solution of **61** (1.20 g, 2.62 mmol) in ethylene glycol dimethyl ether (75 mL), was added furan-3-boronic acid (0.88 g, 7.85 mmol), tris(dibenzylideneacetone)dipalladium(0) (80.0 mg, 0.087 mmol), triphenylphosphine (343 mg, 1.31 mmol), and 2 M sodium carbonate solution (10 mL). The solution was
10 refluxed under nitrogen overnight. After cooling to room temperature, the reaction mixture was diluted with 100 mL water. The aqueous solution was extracted with methylene chloride (3 x 150 mL). The extracts were combined and washed with water (450 mL) and brine (450 mL). The organic solution was dried over sodium sulfate, filtered, and concentrated. The product was purified by silica gel chromatography
15 (90:10:1 $CHCl_3$ /methanol/ NH_4OH) to yield **164** (700 mg, 60%).

Example 156 - Preparation of Compound 164 · HCl

To a stirred solution of **164** (100 mg, 0.224 mmol) in ethyl acetate (6
20 mL) was added 2 M HCl in diethyl ether (135 μ L). After 20 min, the solution was concentrated to afford the HCl salt of **164** in quantitative yield: mp 320-330 °C; ¹H NMR (300 MHz, CD_3OD) δ 8.35 (brs, 1 H), 7.90 (s, 1 H), 7.57 (m, 3 H), 7.42 (d, 2 H), 6.79 (s, 1 H), 4.65-5.07 (m, 5 H), 3.87 (m, 1 H), 3.13 (m, 1 H), 2.00-2.30 (m, 4 H), 1.35-1.75 (m, 10 H); API MS m/z = 446 [$C_{25}H_{31}N_7O+H$]⁺.

25

Example 157 - Preparation of Compound 166

To a stirred solution of **164** (200 mg, 0.449 mmol) in 1,2-dichloromethane (2 mL) was added propionaldehyde (26.0 μ L). After stirring under
30 nitrogen for 20 min, sodium triacetoxyborohydride (114 mg, 0.539 mmol) was added. The mixture was stirred overnight. The reaction was concentrated and purified by silica gel chromatography (90:10:1 $CHCl_3$ /methanol/ NH_4OH) to yield **166**.

Example 158 - Preparation of Compound 166 ·HCl

To a stirred solution of 166 (120 mg, 0.227 mmol) in ethyl acetate (7 mL) was added 2 M HCl in diethyl ether (150 µL). After 20 min, the solution was concentrated to afford the HCl salt of 166 in quantitative yield: mp 285-286 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.42 (brs, 1 H), 7.90 (s, 1 H), 7.49-7.62 (m, 3 H), 7.43 (d, 2 H), 6.80 (s, 1 H), 4.76-5.10 (m, 3 H), 3.91 (m, 1 H), 2.93-3.49 (m, 5 H), 2.00-2.40 (m, 4 H), 1.35-1.95 (m, 14 H), 1.03 (t, 6 H); API MS *m/z* = 530 [C₃₁H₄₃N₇O+H]⁺.

Example 159 - Preparation of Compound 165

To a 0 °C stirred solution of 164 (200 mg, 0.449 mmol) in methylene chloride (8 mL) was added pyridine (55.0 µL), acetic anhydride (46.0 µL, 0.449 mmol), and DMAP (6.0 mg, 0.045 mmol). After 40 min, the reaction mixture warmed to room temperature. After stirring overnight, the reaction mixture was concentrated and dried in vacuo. The material was purified by silica gel chromatography (90:10:1 CHCl₃/methanol/NH₄OH) to yield 165 (190 mg, 87%): mp 241-243 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.86 (s, 1 H), 7.80 (s, 1 H), 7.45-7.54 (m, 3 H), 7.32-7.40 (m, 2 H), 6.76 (s, 1 H), 4.71 (s, 2 H), 4.62 (m, 1 H), 3.74 (m, 1 H), 3.61 (m, 1 H), 1.73-2.11 (m, 7 H), 1.54 (d, 6 H), 1.20-1.40 (m, 4 H); API MS *m/z* = 488 [C₂₇H₃₃N₇O₂+H]⁺.

Example 160 - Preparation of Compounds 167 and 168

Compound 75 (0.500 g, 1.10 mmol) was dissolved in 1,2-dichloroethane (10 mL). To this stirred solution was added acetaldehyde (0.054 g, 1.22 mmol) and sodium triacetoxyborohydride (0.360 g, 1.71 mmol). After 1.5 h, the reaction was quenched with saturated sodium bicarbonate solution (10 mL). The mixture was extracted with ethyl acetate (10 mL). The organic layers were combined, dried over sodium sulfate, and concentrated. The resulting material was purified via silica gel chromatography (60:1:1 CH₂Cl₂/methanol/triethylamine) to yield 167 (213 mg, 40%), and 168 (109 mg, 19%).

Example 161 - Preparation of Compound 167 ·HCl

To a stirred solution of 167 (213 mg, 0.440 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.264 mL). The organic liquid was concentrated to afford the HCl salt of 167 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.30 (brs, 1 H), 7.30-7.63 (m, 10 H), 4.71-4.80 (m, 3 H), 3.65-3.72 (m, 1 H), 3.48 (q, 2 H), 3.10 (brs, 2 H), 2.10-2.20 (m, 4 H), 1.60 (d, 6 H), 1.30-1.59 (m, 7 H); API MS *m/z* = 484 [C₂₉H₃₇N₇+H]⁺.

Example 162 - Preparation of Compound 168 ·HCl

To a stirred solution of 168 (109 mg, 0.213 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.128 mL). The organic liquid was concentrated to afford the HCl salt of 168 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 7.25-7.67 (m, 10 H), 4.60-4.80 (m, 3 H), 3.68-3.80 (m, 1 H), 3.48 (q, 4 H), 2.78-2.95 (m, 4 H), 2.02-2.17 (m, 2 H), 1.10-1.59 (m, 10 H), 1.35 (t, 6 H); API MS *m/z* = 512 [C₃₁H₄₁N₇+H]⁺.

Example 163 - Preparation of Compounds 169 and 170

Compound 75 (0.500 g, 1.10 mmol) was dissolved in 1,2-dichloroethane (10 mL). To this stirred solution was added butyraldehyde (0.072 g, 1.00 mmol) and sodium triacetoxyborohydride (0.297 g, 1.40 mmol). After 2.5 h, the reaction was quenched with saturated sodium bicarbonate solution (10 mL). The mixture was extracted with ethyl acetate (10 mL). The organic layers were combined, dried over sodium sulfate, and concentrated. The resulting material was purified via silica gel chromatography (200:10:1 CH₂Cl₂/methanol/NH₄OH) to yield 169 (180 mg, 32%), and 170 (160 mg, 26%).

Example 164 - Preparation of Compound 169 ·HCl

To a stirred solution of 169 (170 mg, 0.332 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.199 mL). The organic liquid was concentrated to afford the HCl salt of 169 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.30-8.34 (m, 1 H), 7.30-7.68 (m, 10 H), 5.30 (brs, 1 H), 4.65-4.90 (m, 4

- 190 -

H), 3.80-3.92 (m, 2 H), 2.94-3.35 (m, 4 H), 2.15-2.32 (m, 4 H), 1.41-1.75 (m, 12 H), 1.00 (t, 3 H).

Example 165 - Preparation of Compound 171

5

To a stirred, 0 °C solution of 169 (125 mg, 0.228 mmol) in methylene chloride (10 mL), was added pyridine (46 µL), DMAP (6.0 mg, 0.046 mmol), and acetic anhydride (24.0 µL, 0.251 mmol). After 1 h under a nitrogen atmosphere, the reaction mixture was warmed to room temperature. After stirring overnight, another 10 2.2 equivalents of acetic anhydride and 0.2 equivalents of DMAP were added and the mixture was heated to reflux. Following concentration, the material was diluted with ethyl acetate (20 mL) and saturated sodium bicarbonate solution (20 mL). The organic layer was concentrated and dried in vacuo. The resulting material was purified via silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) and trituration with 15 hexanes to yield 171 (26 mg): API MS *m/z* = 554 [C₃₃H₄₃N₇O+H]⁺.

Example 166 - Preparation of Compound 170 ·HCl

To a stirred solution of 170 (150 mg, 0.264 mmol) in ethyl acetate (10 20 mL) was added 2 M HCl in diethyl ether (0.158 mL). The organic liquid was concentrated to afford the HCl salt of 170 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.15-8.25 (m, 1 H), 7.31-7.68 (m, 10 H), 4.65-4.90 (m, 3 H), 3.70-3.95 (m, 1 H), 2.95-3.41 (m, 6 H), 2.05-2.32 (m, 4 H), 1.31-1.79 (m, 18 H), 1.00 (t, 6 H).

25 **Example 167 - Preparation of Compounds 172 and 173**

Compound 75 (0.500 g, 1.10 mmol) was dissolved in 1,2-dichloroethane (10 mL). To this stirred solution was added cyclopropanecarboxaldehyde (0.070 g, 1.00 mmol) and sodium triacetoxymethylborohydride 30 (0.297 g, 1.40 mmol). After 3 h, the reaction was quenched with saturated sodium bicarbonate solution (10 mL). The mixture was extracted with ethyl acetate (10 mL). The organic layers were combined, dried over sodium sulfate, and concentrated. The resulting material was purified via silica gel chromatography (200:10:1 CH₂Cl₂/methanol/NH₄OH) to yield 172 (103 mg, 18%), and 173 (160 mg, 26%).

Example 168 - Preparation of Compound 172 ·HCl

To a stirred solution of 172 (103 mg, 0.202 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.121 mL). The organic liquid was concentrated to afford the HCl salt of 172 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.30 (brs, 1 H), 7.30-7.69 (m, 10 H), 4.69-4.92 (m, 4 H), 3.80-3.92 (m, 1 H), 2.84-3.19 (m, 3 H), 2.11-2.28 (m, 4 H), 1.37-1.72 (m, 10 H), 1.05-1.15 (m, 1 H), 0.68-0.74 (m, 2 H), 0.38-0.42 (m, 2 H).

Example 169 - Preparation of Compound 173 ·HCl

To a stirred solution of 173 (160 mg, 0.284 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.170 mL). The organic liquid was concentrated to afford the HCl salt of 173 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.30-8.41 (m, 1 H), 7.30-7.68 (m, 10 H), 5.30 (brs, 1 H), 4.68-4.90 (m, 4 H), 3.55-3.95 (m, 2 H), 3.05-3.20 (m, 4 H), 2.00-2.32 (m, 4 H), 1.10-1.90 (m, 10 H), 0.75-0.80 (m, 4 H), 0.35-0.50 (m, 4 H).

Example 170 - Preparation of Compounds 174 and 175

Compound 75 (1.50 g, 3.29 mmol) was dissolved in 1,2-dichloroethane (30 mL). To this stirred solution was added propionaldehyde (0.174 g, 2.99 mmol) and sodium triacetoxymethylborohydride (0.888 g, 4.19 mmol). After 1.5 h, the reaction was quenched with saturated sodium bicarbonate solution (30 mL). The mixture was extracted with ethyl acetate (30 mL). The organic layers were combined, dried over sodium sulfate, and concentrated. The resulting material was purified via silica gel chromatography (200:10:1 CH₂Cl₂/methanol/NH₄OH) to yield 174 (317 mg, 19%): ¹H NMR (300 MHz, CD₃OD) δ 7.26-7.80 (m, 10 H), 4.62-4.81 (m, 3 H), 3.74 (brs, 1 H), 2.41-2.62 (m, 3 H), 1.90-2.11 (m, 4 H), 1.52 (d, 6 H), 1.12-1.52 (m, 8 H), 0.92 (t, 3 H), and 175 (320 mg, 18%): ¹H NMR (300 MHz, CD₃OD) δ 8.18-8.28 (m, 1 H), 7.28-7.68 (m, 10 H), 4.65-4.90 (m, 3 H), 3.81-3.94 (m, 1 H), 2.94-3.25 (m, 4 H), 2.02-2.31 (m, 6 H), 1.40-1.81 (m, 14 H), 1.05 (t, 6 H).

Example 171 - Preparation of Compound 267

The HCl salt of **167** (10 mg, 0.037 mmol), was dispersed in ethyl acetate and neutralized with sodium bicarbonate. The organic material was dried over magnesium sulfate and concentrated. The solid was dissolved in dry CH₂Cl₂ (10 mL) and cooled to 0 °C. To the solution was added DMAP (9 mg), pyridine (0.074 mL) and acetic anhydride (0.037 mL). The ice bath was removed after 1 h. After being stirred overnight, additional DMAP and Ac₂O was added in portions to consume starting material by TLC analysis. The mixture was heated to reflux for 2 d. Upon cooling, the mixture was concentrated in vacuo, then neutralized with aqueous sodium bicarbonate, extracted with ethyl acetate, dried and concentrated. The residue was purified by chromatography to provide **267**: API MS $m/z = 526$ [C₃₁H₃₉N₇O+H]⁺.

Example 172 - Preparation of Compound 177

To a solution of **61** (1.00 g, 2.21 mmol) and 3-tolylboronic acid (0.33 g, 2.43 mmol) in tetrahydrofuran (5 mL) was added tris(dibenzylideneacetone)dipalladium(0) (0.010 g, 0.011 mmol), tri-*tert*-butylphosphine (5.5 mg, 0.027 mmol), and potassium fluoride (0.42 g, 7.29 mmol). After mixing overnight at room temperature, the reaction mixture was refluxed for 24 h and cooled to room temperature. The reaction mixture was diluted with ether (50 mL) and filtered through Celite. The organic liquid was concentrated and the resulting material was purified via silica gel chromatography (90:10:1 CH₂Cl₂/methanol/NH₄OH) to yield **177** (0.70 g, 71%).

Example 173 - Preparation of Compound 177 · HCl

To a stirred solution of **177** (449 mg, 0.956 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (575 µL). The organic liquid was concentrated to afford the HCl salt of **177** in quantitative yield: mp 186-195 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.82 (s, 1 H), 7.54 (d, 2 H), 7.20-7.48 (m, 5 H), 7.12 (d, 1 H), 4.83-5.10 (m, 2 H), 4.77 (s, 2 H), 4.64 (m, 1 H), 3.77 (m, 1 H), 3.03 (m, 1 H), 2.38 (s, 3 H), 1.93-2.20 (m, 4 H), 1.19-1.70 (m, 10 H); API MS $m/z = 470$ [C₂₈H₃₅N₇+H]⁺.

Example 174 - Preparation of Compound 178

To a stirred solution of 177 (219 mg, 0.466 mmol) in methylene chloride (25 mL) was added acetic anhydride (48 μ L, 0.513 mmol), DMAP (5.7 mg, 0.047 mmol), and pyridine (57.0 μ L, 0.699 mmol). The mixture was placed under a nitrogen atmosphere and immersed in an ice water bath. After 30 min, the reaction mixture was warmed to room temperature and stirred for another 1.5 h. The solution was concentrated and the resulting material was purified via silica gel chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to afford 178.

Example 175 - Preparation of Compound 178 ·HCl

To a stirred solution of 178 (50.0 mg, 0.098 mmol) in ethyl acetate (5 mL) was added 2 M HCl in diethyl ether (59.0 μ L). The organic liquid was concentrated to afford the HCl salt of 178 in quantitative yield: mp 165-174 °C; ¹H NMR (300 MHz, CD₃OD) δ 8.26 (brs, 1 H), 7.61 (d, 2 H), 7.47 (d, 2 H), 7.24-7.41 (m, 3 H), 7.15 (d, 1 H), 4.62-5.08 (m, 4 H), 3.83 (m, 1 H), 3.67 (m, 1 H), 2.39 (s, 3 H), 1.87-2.20 (m, 7 H), 1.60 (d, 6 H), 1.40 (m, 4 H); API MS *m/z* = 512 [C₃₀H₃₇N₇O+H]⁺.

Example 176 - Preparation of Compound 179

To a solution of 61 (2.00 g, 4.36 mmol) and 3-methoxyphenyl boronic acid (0.73 g, 4.79 mmol) in tetrahydrofuran (10 mL) was added tris(dibenzylideneacetone)dipalladium(0) (20 mg, 0.022 mmol), tri-*tert*-butylphosphine (10.0 mg, 0.052 mmol), and potassium fluoride (0.84 g, 14.39 mmol). After refluxing overnight, the reaction mixture was diluted with ether (50 mL) and filtered through Celite. The organic liquid was concentrated and the resulting material was purified via silica gel chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to yield 179 (1.18 g, 56%).

Example 177 - Preparation of Compound 179 ·HCl

To a stirred solution of 179 (980 mg, 2.02 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (1.21 mL). The organic liquid was

concentrated to afford the HCl salt of 179 in quantitative yield: mp 182-189 °C; ¹H NMR (300 MHz, CD₃OD) δ 7.80 (s, 1 H), 7.54 (d, 2 H), 7.40 (d, 2 H), 7.30 (t, 1 H), 7.12 (m, 2 H), 6.87 (m, 1 H), 4.76 (s, 2 H), 4.62 (m, 1 H), 3.67-3.90 (m, 4 H), 2.94 (m, 1 H), 1.90-2.20 (m, 4 H), 1.17-1.65 (m, 10 H); API MS *m/z* = 486
5 [C₂₈H₃₅N₇O+H]⁺.

Example 178 - Preparation of Compound 180

To a stirred solution of 179 (200 mg, 0.412 mmol) in methylene
10 chloride (25 mL) was added acetic anhydride (43 μL, 0.450 mmol), DMAP (5.0 mg, 0.041 mmol), and pyridine (50.0 μL, 0.618 mmol). The mixture was placed under a nitrogen atmosphere and immersed in an ice water bath. After 30 min, the reaction mixture was warmed to room temperature and stirred for another 1.5 h. The solution was concentrated and the resulting material was purified via silica gel
15 chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to afford 180.

Example 179 - Preparation of Compound 180 ·HCl

To a stirred solution of 180 (60.0 mg, 0.114 mmol) in ethyl acetate (5
20 mL) was added 2 M HCl in diethyl ether (68.0 μL). The organic liquid was concentrated to afford the HCl salt of 180 in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.20 (brs, 1 H), 7.62 (d, 2 H), 7.46 (d, 2 H), 7.34 (t, 1 H), 7.10-7.20 (m, 2 H), 6.90 (1 H), 4.60-5.10 (m, 4 H), 3.83 (s, 3 H), 3.64 (m, 1 H), 3.20-3.42 (m, 1 H), 1.87-2.18 (m, 7 H), 1.60 (d, 6 H), 1.22-1.50 (m, 4 H); API MS *m/z* = 528
25 [C₃₀H₃₇N₇O₂+H]⁺.

Example 180 - Preparation of Compound 181

To a solution of 61 (2.00 g, 4.36 mmol) and furan-2-boronic acid (1.50
30 g, 13.1 mmol) in ethylene glycol dimethyl ether (150 mL) was added tris(dibenzylideneacetone)dipalladium(0) (120 mg, 0.130 mmol), tri-*tert*-butylphosphine (570 mg, 2.18 mmol), and 2 M sodium carbonate solution (12.5 mL, 25.3 mmol). After refluxing overnight, 2 more equivalents of furan-2-boronic acid were added. The reaction was refluxed for 24 h, cooled to room temperature, and

- 195 -

diluted with water (50 mL). The aqueous mixture was extracted with methylene chloride (3 x 80 mL). The extracts were combined and washed with water (250 mL) and brine (250 mL). The organic phase was dried over sodium sulfate and filtered. The organic liquid was concentrated and the resulting material was purified via silica gel chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to yield **181**.

Example 181 - Preparation of Compound 181 ·HCl

To a stirred solution of **181** (600 mg, 1.35 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.810 mL). The organic liquid was concentrated to afford the HCl salt of **181** (406 mg, 68%): ¹H NMR (300 MHz, CD₃OD) δ 8.06 (s, 1 H), 7.66 (d, 2 H), 7.53 (s, 1 H), 7.41 (d, 2 H), 6.74 (m, 1 H), 6.49 (m, 1 H), 4.60-5.00 (m, 5 H), 3.82 (m, 1 H), 3.10 (m, 1 H), 1.95-2.20 (m, 4 H), 1.20-1.61 (m, 10 H); API MS *m/z* = 446 [C₂₅H₃₁N₇O+H]⁺.

Example 182 - Preparation of Compound 182

To a stirred solution of **181** (750 mg, 1.68 mmol) in methylene chloride (30 mL) was added acetic anhydride (0.18 mL, 1.85 mmol), DMAP (20.8 mg, 0.17 mmol), and pyridine (0.20 mL, 2.52 mmol). The mixture was placed under a nitrogen atmosphere and immersed in an ice water bath. After 30 min, the reaction mixture was warmed to room temperature and stirred for another 1.5 h. The solution was concentrated and the resulting material was purified via silica gel chromatography (95:5:1 CH₂Cl₂/methanol/NH₄OH) to afford **182** (530 mg, 65%).

Example 183 - Preparation of Compound 182 ·HCl

To a stirred solution of **182** (300 mg, 0.620 mmol) in ethyl acetate (10 mL) was added 2 M HCl in diethyl ether (0.370 mL). The organic liquid was concentrated to afford the HCl salt of **182** in quantitative yield: ¹H NMR (300 MHz, CD₃OD) δ 8.25 (brs, 1 H), 7.56-7.73 (m, 2 H), 7.54 (s, 1 H), 7.30-7.47 (m, 2 H), 6.75 (m, 1 H), 6.49 (m, 1 H), 4.60-5.05 (m, 4 H), 3.73-3.90 (m, 1 H), 3.55-3.73 (m, 1 H), 1.82-2.23 (m, 7 H), 1.15-1.70 (m, 10 H); API MS *m/z* = 488 [C₂₇H₃₃N₇O₂+H]⁺.

Example 184 - Preparation of Compound 183

To a stirred mixture of 6-chloronicotinamide (5.00 g, 31.9 mmol) in ethanol (13 mL) and toluene (80 mL) was added 3-fluorobenzeneboronic acid (4.92 g, 35.1 mmol) and 2 M sodium carbonate solution (32 mL). The suspension was heated to 80 °C and degassed with argon for 1 h. After cooling to room temperature, tetrakis(triphenylphosphine)palladium(0) (1.11 g, 0.958 mmol) was added. The reaction mixture was refluxed under argon for 3 h. After cooling to room temperature, the mixture was diluted with water (100 mL) and filtered. The filter cake was washed with water and dried in vacuo to afford **183** (6.38 g, 92%).

Example 185 - Preparation of Compound 184

To a stirred suspension of **183** (3.00 g, 13.9 mmol) in tetrahydrofuran (25 mL) was added dropwise 1 M borane in THF (97.0 mL, 97.0 mmol). After refluxing for 2 h, the reaction mixture was cooled in an ice bath. The mixture was acidified to pH 1 with 2 N HCl and stirred for 1 h. The pH was raised to a value of 10 by adding 6 N NaOH and the resulting solution was extracted with ethyl acetate (3 x 50 mL). The extractions were combined, washed with brine (150 mL), and dried over sodium sulfate. The suspension was filtered and concentrated. The resulting material was purified by precipitation as the HCl salt from an ethanol solution. The product was recovered by filtration and dried in vacuo to yield **184** (1.69 g, 51%): ¹H NMR (300 MHz, CD₃OD) δ 8.92 (s, 1 H), 8.41 (d, 1 H), 8.27 (d, 1 H), 7.77-7.90 (m, 2 H), 7.35-7.70 (m, 2 H), 4.36 (s, 2 H).

Example 186 - Preparation of Compound 185

The amine **184** (1.84 g, 7.69 mmol), 2,6-dichloropurine (1.31 g, 6.99 mmol), and *N,N*-diisopropylethylamine (2.68 mL, 15.4 mmol) were dissolved in ethanol (65 mL). After refluxing overnight, the solution was immersed in an ice water bath for 20 min. The mixture was filtered and cake was washed with water. The cake was triturated with ethanol and diethyl ether and dried in vacuo to afford **185** (1.12 g, 47%).

Example 187 - Preparation of Compound 186

To a stirred solution of 185 (1.00 g, 2.94 mmol) in dimethylsulfoxide (100 mL) was added potassium carbonate (2.19 g, 15.9 mmol) and 2-iodopropane
5 (0.88 mL, 8.81 mmol). The mixture was placed under an argon atmosphere and stirred overnight. The reaction mixture was poured into stirred water (300 mL) and the resulting solution was extracted with ethyl acetate (3 x 300 mL). The extractions were combined, washed with water (900 mL) and brine (900 mL), and dried over magnesium sulfate. Following filtration, the organic liquid was concentrated. The
10 material was purified by recrystallization from ethyl acetate in hexanes to yield 186 (0.92 g, 82%).

Example 188 - Preparation of Compound 187

In a sealed tube were combined 186 (640 mg, 1.67 mmol), *trans*-1,4-diaminocyclohexane (0.96 g, 8.36 mmol), and ethanol (3.5 mL). The reaction mixture was heated to 150 °C for 4 d and cooled to room temperature. The solution was poured into stirred ice water (5 mL) and the resulting mixture was extracted with methylene chloride (3 x 5 mL). The extractions were combined, washed with water
20 (15 mL) and brine (15 mL), and dried over sodium sulfate. The organic liquid was concentrated. Purification by column chromatography (97:3 CH₂Cl₂/methanol) and trituration with hexanes yielded the free base, 187: ¹H NMR (300 MHz, CDCl₃) δ 8.68 (s, 1 H), 7.60-7.82 (m, 4 H), 7.50 (s, 1 H), 7.42 (q, 1 H), 7.11 (m, 1 H), 6.07 (m, 1 H), 4.85 (d, 2 H), 4.60 (m, 2 H), 3.74 (m, 1 H), 2.69 (m, 1 H), 2.02-2.19 (m, 2 H),
25 1.78-1.96 (m, 2 H), 1.52 (d, 6 H), 1.10-1.36 (m, 4 H); API MS *m/z* = 475 [C₂₆H₃₁FN₈+H]⁺.

Example 189 - Preparation of Compound 188

The free amine 187 (80.0 mg, 0.170 mmol) was dissolved in methylene chloride (4 mL). The solution was immersed in an ice water bath and acetic anhydride (17.5 μL, 0.185 mmol), DMAP (2.0 mg, 0.017 mmol), and pyridine (20.0 μL, 0.252 mmol) were added. After stirring for 30 min, the solution was warmed to room temperature and concentrated. Purification via silica gel chromatography and
35 trituration with hexanes yielded 188 (67 mg, 78%): mp 199-230 °C; ¹H NMR (300

MHz, CDCl₃) δ 8.71 (s, 1 H), 7.62-7.85 (m, 4 H), 7.52 (s, 1 H), 7.42 (m, 1 H), 7.10 (m, 1 H), 6.01 (m, 1 H), 5.28 (d, 1 H), 4.81 (d, 2 H), 4.62 (m, 2 H), 3.72 (m, 2 H), 1.90-2.27 (m, 7 H), 1.53 (d, 6 H), 1.24 (m, 4 H); API MS m/z = 517 [C₂₈H₃₃FN₈O+H]⁺.

5

Example 190 - Preparation of Compound 189

To a stirred mixture of 6-chloronicotinamide (3.00 g, 19.2 mmol) in ethanol (7.6 mL) and toluene (48 mL) were added 3-methoxyphenylboronic acid
10 (3.20 g, 21.1 mmol) and 2 M sodium carbonate solution (19 mL). The suspension was heated to 80 °C and degassed with argon for 1 h. After cooling to room temperature, tetrakis(triphenylphosphine)palladium(0) (664 mg, 0.575 mmol) was added. The reaction mixture was refluxed under argon for 3 h. After cooling to room temperature, the mixture was diluted with water (100 mL) and filtered. The filter cake was washed
15 with water and dried in vacuo to afford 189 (3.62 g, 83%).

Example 191 - Preparation of Compound 190

To a stirred solution of 189 (3.00 g, 13.1 mmol) in tetrahydrofuran (25
20 mL) was added dropwise 1 M borane in THF (92.0 mL, 92.0 mmol). After refluxing for 4 h, the reaction mixture was cooled in an ice bath. The mixture was acidified to pH 1 with 2 N HCl and stirred for 1 h. The pH was raised to a value of 10 by adding 6 N NaOH and the resulting solution was extracted with ethyl acetate (3 x 50 mL). The extractions were combined, washed with brine (150 mL), and dried over sodium
25 sulfate. The suspension was filtered and concentrated. The resulting material was purified by precipitation as the HCl salt from an ethanol solution. The product was recovered by filtration and dried in vacuo to yield 190 (1.81 g, 55%).

Example 192 - Preparation of Compound 191

30

The amine 190 (1.80 g, 7.18 mmol), 2,6-dichloropurine (1.22 g, 6.53 mmol), and *N,N*-diisopropylethylamine (1.86 g, 6.53 mmol) were dissolved in ethanol (82 mL). After refluxing overnight, the dispersion was immersed in an ice water bath for 60 min. The mixture was filtered and cake was washed with water. The cake was

- 199 -

trituated with ethanol and diethyl ether and dried in vacuo to afford **191** (1.04 g, 44%).

Example 193 - Preparation of Compound 192

5 To a stirred solution of **191** (1.04 g, 2.84 mmol) in dimethylsulfoxide (60 mL) was added potassium carbonate (2.12 g, 15.3 mmol) and 2-iodopropane (0.85 mL, 8.52 mmol). The mixture was placed under an argon atmosphere and stirred overnight. The reaction mixture was poured into stirred water (60 mL) and the
10 resulting solution was extracted with ethyl acetate (3 x 60 mL). The extractions were combined, washed with water (180 mL) and brine (180 mL), and dried over magnesium sulfate. Following filtration, the organic liquid was concentrated. The material was purified by recrystallization from ethyl acetate in hexanes (1:40) to yield **192**.

15 **Example 194 - Preparation of Compound 193**

In a sealed tube were combined **192** (400 mg, 1.09 mmol), *trans*-1,4-diaminocyclohexane (1.25 g, 10.9 mmol), and ethanol (4.0 mL). The reaction mixture
20 was heated to 150 °C for 24 h and cooled to room temperature. The solution was filtered and the filtrate was concentrated. Purification by column chromatography (97:3 CH₂Cl₂/methanol) yielded the free base, **193** (240 mg, 45%).

Example 195 - Preparation of Compound 194

25 The free amine **193** (130 mg, 0.27 mmol) was dissolved in methylene chloride (6 mL). The solution was immersed in an ice water bath and acetic anhydride (28.0 µL, 0.294 mmol), DMAP (3.2 mg, 0.026 mmol), and pyridine (33.0 µL, 0.401 mmol) were added. After stirring for 30 min, the solution was warmed to room
30 temperature and concentrated. Purification via prep-TLC (9:1 CH₂Cl₂/methanol) and trituration with ethyl acetate yielded **194**: mp 161-163 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.76 (s, 1 H), 7.77 (d, 1 H), 7.64 (d, 1 H), 7.50 (m, 3 H), 7.35 (t, 1 H), 6.73-7.00 (m, 2 H), 5.33 (d, 1 H), 4.50-4.92 (m, 4 H), 3.56-4.00 (m, 5 H), 1.84-2.22 (m, 7 H), 1.55 (d, 6 H), 1.25 (m, 4 H); API MS *m/z* = 529 [C₂₉H₃₆N₈O₂+H]⁺.

- 200 -

Example 196 - Preparation of Compound 195

To a stirred mixture of 6-chloronicotinamide (2.00 g, 12.8 mmol) in ethanol (5.0 mL) and toluene (32 mL) was added thiophene-2-boronic acid (1.80 g, 14.1 mmol) and 2 M sodium carbonate solution (13 mL). The suspension was heated to 80 °C and degassed with argon for 1 h. After cooling to room temperature, tetrakis(triphenylphosphine)palladium(0) (443 mg, 0.383 mmol) was added. The reaction mixture was refluxed under argon for 3 h. After cooling to room temperature, another 0.950 g thiophene-2-boronic acid and 280 mg tetrakis(triphenylphosphine)palladium(0) were added to the reaction mixture. It was refluxed for 4 h and cooled to room temperature. The mixture was diluted with water (50 mL) and filtered. The filter cake was washed with water and dried in vacuo to afford **195** (1.65 g, 63%).

Example 197 - Preparation of Compound 196

To a stirred solution of **195** (1.40 g, 6.86 mmol) in tetrahydrofuran (23 mL) was added dropwise 1 M borane in THF (48.0 mL, 48.0 mmol). After refluxing for 1 h, the reaction mixture was cooled in an ice bath. The mixture was acidified to pH 1 with 2 N HCl and stirred for 1 h. The pH was raised to a value of 10 by adding 6 N NaOH and the resulting solution was extracted with ethyl acetate (3 x 50 mL). The extractions were combined, washed with brine (150 mL), and dried over sodium sulfate. The suspension was filtered and concentrated. The resulting material was purified by precipitation as the HCl salt from an ethanol solution. The product was recovered by filtration and dried in vacuo to yield **196** (0.87 g, 56%).

Example 198 - Preparation of Compound 197

The amine **196** (210 mg, 1.10 mmol), 2,6-dichloropurine (188 mg, 1.00 mmol), and *N,N*-diisopropylethylamine (286 g, 2.21 mmol) were dissolved in ethanol (12 mL). After refluxing overnight, the suspension was immersed in an ice water bath for 60 min. The mixture was filtered and cake was washed with water. The cake was triturated with ethanol and diethyl ether and dried in vacuo to afford **197** (206 mg, 60%).

- 201 -

Example 199 - Preparation of Compound 198

To a stirred solution of 197 (200 mg, 0.583 mmol) in dimethylsulfoxide (12 mL) was added potassium carbonate (435 mg, 3.15 mmol) and 2-iodopropane (0.18 mL, 1.75 mmol). The mixture was placed under an argon atmosphere and stirred overnight. The reaction mixture was poured into stirred water (15 mL) and the resulting solution was extracted with ethyl acetate (3 x 30 mL). The extractions were combined, washed with water (90 mL) and brine (90 mL), and dried over magnesium sulfate. Following filtration, the organic liquid was concentrated to yield 198 (200 mg, 89%).

Example 200 - Preparation of Compound 199

In a sealed tube were combined 198 (100 mg, 0.260 mmol), *trans*-1,4-diaminocyclohexane (297 mg, 2.60 mmol), and ethanol (2.0 mL). The reaction mixture was heated to 150 °C for 2 d and cooled to room temperature. The solution was filtered and diluted with ethanol. The filtrate was concentrated, and converted to its HCl salt to afford 199: ¹H NMR (300 MHz, CD₃OD) δ 9.00 (s, 1 H), 8.72 (d, 2 H), 8.46 (d, 1 H), 8.27 (m, 1 H), 8.15 (d, 1 H), 7.54 (m, 1 H), 4.89-5.35 (3 H), 4.04 (m, 1 H), 3.35 (m, 1 H), 2.20-2.50 (m, 4 H), 1.55-1.90 (m, 10 H); ESI MS *m/z* = 463 [C₂₄H₃₀N₈S+H]⁺.

Example 201 - Preparation of Compound 200

The free amine 199 (100 mg, 0.216 mmol) was dissolved in methylene chloride (5 mL). The solution was immersed in an ice water bath and acetic anhydride (20.0 μL, 0.216 mmol), DMAP (2.6 mg, 0.021 mmol), and pyridine (33.0 μL, 0.324 mmol) were added. After stirring for 30 min, the solution was warmed to room temperature and concentrated. Purification via prep-TLC (10:1 CH₂Cl₂/methanol) yielded 200 (20 mg): mp 206-208 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.56 (s, 1 H), 7.70 (d, 1 H), 7.58 (d, 1 H), 7.53 (m, 2 H), 7.37 (m, 1 H), 7.10 (m, 1 H), 6.15 (brs, 1 H), 5.33 (d, 1 H), 4.76 (d, 2 H), 4.61 (m, 2 H), 3.70 (m, 2 H), 1.75-2.15 (m, 7 H), 1.52 (d, 6 H), 1.23 (m, 4 H); API MS *m/z* = 505 [C₂₆H₃₂N₈OS+H]⁺.

Example 202 - Preparation of Compound 201

To a stirred solution of 6-chloronicotinamide (2.80 g, 17.9 mmol) in ethanol (7.5 mL) and toluene (48 mL) was added furan-2-boronic acid (3.00 g, 26.8 mmol) and 2 M sodium carbonate solution (18 mL). The suspension was heated to 80 °C and degassed with argon for 1 h. After cooling to room temperature, tetrakis(triphenylphosphine)palladium(0) (619 mg, 0.536 mmol) was added. The reaction mixture was refluxed under argon for 2 d then cooled to room temperature. The mixture was diluted with water (75 mL) and filtered. The filter cake was washed with water and dried in vacuo to afford **201** (1.95 g, 58%).

Example 203 - Preparation of Compound 202

To a stirred solution of **201** (1.69 g, 8.98 mmol) in tetrahydrofuran (34 mL) was added dropwise 1 M borane in THF (50.0 mL, 50.0 mmol). After refluxing for 2 h, the reaction mixture was cooled in an ice bath. The mixture was acidified to pH 1 with 2 N HCl and stirred for 1 h. The pH was raised to a value of 10 by adding 6 N NaOH and the resulting solution was extracted with ethyl acetate (3 x 50 mL). The extractions were combined, washed with brine (150 mL), and dried over sodium sulfate. The suspension was filtered and concentrated. The resulting material was purified by precipitation as the HCl salt from an ethanol solution. The product was recovered by filtration and dried in vacuo to yield **202** (1.12 g, 50%).

Example 204 - Preparation of Compound 203

The amine **202** (177 mg, 1.02 mmol), 2,6-dichloropurine (173 mg, 0.920 mmol), and *N,N*-diisopropylethylamine (267 g, 2.07 mmol) were dissolved in ethanol (11 mL). After refluxing overnight, the suspension was immersed in an ice water bath for 60 min. The mixture was filtered and cake was washed with water. The cake was triturated with ethanol and diethyl ether and dried in vacuo to afford **203** (166 mg, 54%).

Example 205 - Preparation of Compound 204

To a stirred solution of **203** (166 mg, 0.508 mmol) in dimethylsulfoxide (11 mL) was added potassium carbonate (379 mg, 2.74 mmol) and 2-iodopropane (0.150 mL, 1.52 mmol). The mixture was placed under an argon atmosphere and stirred overnight. The reaction mixture was poured into stirred water (15 mL) and the resulting solution was extracted with ethyl acetate (3 x 30 mL). The extractions were combined, washed with water (90 mL) and brine (90 mL), and dried over magnesium sulfate. Following filtration, the organic liquid was concentrated to yield **204** (178 mg, 95%).

Example 206 - Preparation of Compound 205

In a sealed tube were combined **204** (170 mg, 0.461 mmol), *trans*-1,4-diaminocyclohexane (526 mg, 4.61 mmol), and ethanol (2.5 mL). The reaction mixture was heated to 150 °C for 4 d and cooled to room temperature. The solution was filtered and concentrated to afford **205**.

Example 207 - Preparation of Compound 206

The free amine **205** (100 mg, 0.224 mmol) was dissolved in methylene chloride (5 mL). The solution was immersed in an ice water bath and acetic anhydride (21.0 µL, 0.224 mmol), DMAP (2.7 mg, 0.022 mmol), and pyridine (34.0 µL, 0.336 mmol) were added. After stirring for 30 min, the solution was stored at 0 °C overnight. Purification via prep-TLC (9:1 CH₂Cl₂/methanol) yielded **206** (43 mg): mp 216-218 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.63 (s, 1 H), 7.72 (d, 1 H), 7.61 (d, 1 H), 7.50 (m, 2 H), 6.99 (m, 1 H), 6.50 (m, 1 H), 6.15 (brs, 1 H), 5.36 (d, 1 H), 4.79 (d, 2 H), 4.62 (m, 2 H), 3.68 (m, 2 H), 1.78-2.20 (m, 7 H), 1.54 (d, 6 H), 1.22 (m, 4 H); API MS *m/z* = 489 [C₂₆H₃₂N₈O₂+H]⁺.

Example 208 - Preparation of Compound 207

Prepared by reaction of **72** with ethylene diamine by general methods described above (91%): ¹H NMR (300 MHz, (CD₃)₂SO) δ 8.06 (brs, 3 H), 7.65 (d, 4

H), 7.31-7.52 (m, 6 H), 4.68-4.90 (m, 3 H), 3.61 (m, 2 H), 3.01 (m, 2 H), 1.54 (d, 6 H); ESI MS m/z = 402 [$C_{23}H_{27}N_7+H$]⁺.

Example 209 - Preparation of Compound 208

5

Prepared by reaction of 207 under standard acetylation conditions (35%): ¹H NMR (300 MHz, CDCl₃) δ 7.49-7.62 (m, 5 H), 7.30-7.48 (m, 5 H), 6.44 (brs, 1 H), 6.13 (brs, 1 H), 5.05 (t, 1 H), 4.82 (d, 2 H), 4.65 (m, 1 H), 3.58 (m, 2 H), 3.45 (m, 2 H), 1.87 (s, 3 H), 1.54 (d, 6 H); ESI MS m/z = 444 [$C_{25}H_{29}N_7O+H$]⁺.

10

Example 210 - Preparation of Compound 211

Prepared by reaction of 72 and 1,3-propanediamine (28%): ¹H NMR (300 MHz, (CD₃)₂SO) δ 8.00-8.22 (m, 3 H), 7.63-7.70 (d, 4 H), 7.32-7.59 (m, 6 H), 4.84 (m, 2 H), 4.70 (m, 1 H), 4.43 (m, 2 H), 2.88 (m, 2 H), 1.88 (m, 2 H), 1.52 (d, 6 H); ESI MS m/z = 416 [$C_{24}H_{29}N_7+H$]⁺.

15

Example 211 - Preparation of Compound 212

Prepared by reaction of 211 with acetic anhydride under standard conditions (44%): mp 106-107 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.50-7.62 (m, 5 H), 7.28-7.50 (m, 5 H), 5.87 (brs, 2 H), 4.93 (m, 1 H), 4.84 (m, 2 H), 4.66 (m, 1 H), 3.49 (m, 2 H), 3.33 (m, 2 H), 1.91 (s, 3 H), 1.74 (m, 2 H), 1.55 (d, 6 H); ESI MS m/z = 458 [$C_{26}H_{31}N_7O+H$]⁺.

25

Example 212 - Preparation of Compound 213

Prepared by the general methods described above (48%): ¹H NMR (300 MHz, CDCl₃) δ 7.29-7.66 (m, 10 H), 6.40 (brs, 1 H), 4.71-5.03 (m, 3 H), 4.61 (m, 1 H), 3.30-3.55 (m, 2 H), 2.73 (m, 2 H), 2.32 (m, 2 H), 1.38-1.85 (m, 10 H); ESI MS m/z = 430 [$C_{25}H_{31}N_7+H$]⁺.

30

Example 213 - Preparation of Compound 214

Prepared by the general methods described above (45%): ¹H NMR (300 MHz, CDCl₃) δ 7.27-7.65 (m, 10 H), 5.92 (m, 1 H), 5.43 (m, 1 H), 4.83 (m, 3

35

H), 4.66 (m, 1 H), 3.46 (m, 2 H), 3.25 (m, 2 H), 1.93 (s, 3 H), 1.76 (m, 3 H), 1.40-1.70 (m, 8 H); ESI MS m/z = 472 [$C_{27}H_{33}N_7O+H$]⁺.

Example 214 - Preparation of Compounds 209 and 210

5

Prepared by the general methods described above. For 210 (17%): ¹H NMR (300 MHz, (CD₃)₂SO) δ 7.31-7.72 (m, 10 H), 4.64-4.92 (m, 3 H), 3.73 (m, 4 H), 2.84-3.33 (m, 6 H), 1.43-1.79 (m, 10 H), 0.85 (m, 6 H); ESI MS m/z = 486 [$C_{29}H_{39}N_7+H$]⁺. For 209: ¹H NMR (300 MHz, CDCl₃) δ 7.26-7.63 (m, 10 H), 6.46 (brs, 1 H), 5.83 (brs, 1 H), 4.85 (m, 2 H), 4.66 (m, 1 H), 3.79 (d, 2 H), 3.11 (m, 2 H), 2.76 (m, 2 H), 1.76 (m, 2 H), 1.50 (d, 6 H), 0.84 (m, 3 H); ESI MS m/z = 444 [$C_{26}H_{33}N_7+H$]⁺.

10

Example 215 - Preparation of Compound 215

15

Prepared by the general methods described above: ¹H NMR (300 MHz, CDCl₃) δ 7.22-7.60 (m, 10 H), 6.18 (brs, 1 H), 5.11 (brs, 1 H), 4.81 (m, 2 H), 4.64 (m, 1 H), 3.40 (m, 2 H), 2.87 (m, 2 H), 2.77 (m, 2 H), 1.70-2.00 (m, 4 H), 1.55-1.70 (m, 2 H), 1.51 (m, 6 H), 0.92 (t, 3 H); ESI MS m/z = 472 [$C_{28}H_{37}N_7+H$]⁺.

20

Example 216 - Preparation of Compound 216

Prepared by the general methods described above (71%): ¹H NMR (300 MHz, CDCl₃) δ 7.28-7.61 (m, 10 H), 6.01 (brs, 1 H), 4.86 (d, 2 H), 4.66 (m, 1 H), 3.79-3.89 (m, 1 H), 3.70-3.79 (m, 1 H), 3.57-3.70 (m, 2 H), 3.27-3.37 (m, 1 H), 2.10-2.23 (m, 1 H), 1.68-1.82 (m, 1 H), 1.55 (d, 6 H); ESI MS m/z = 428 [$C_{25}H_{29}N_7+H$]⁺.

25

Example 217 - Preparation of Compound 217

30

Prepared by the general methods described above: TLC silica gel R_f = 0.52 (20:1:0.01-CH₂Cl₂/MeOH/NH₄OH).

Example 218 - Preparation of Compound 218

Prepared by the general methods described above: ^1H NMR (300 MHz, $(\text{CD}_3)_2\text{SO}$) δ 8.02 (brs, 1 H), 7.88 (s, 1 H), 7.54-7.68 (m, 4 H), 7.40-7.50 (m, 3 H), 7.30-7.40 (m, 1 H), 7.26 (s, 1 H), 6.77 (s, 1 H), 4.50-4.72 (m, 4 H), 2.78 (t, 2 H), 2.31 (m, 1 H), 1.68 (d, 2 H), 1.49 (d, 6 H); ESI MS $m/z = 470$ $[\text{C}_{27}\text{H}_{31}\text{N}_7\text{O}+\text{H}]^+$.

Example 219 - Preparation of Compound 219

Prepared by the general methods described above (66%): ^1H NMR (300 MHz, CDCl_3) δ 7.28-7.62 (m, 10 H), 5.89 (brs, 1 H), 4.86 (d, 4 H), 4.67 (m, 1 H), 2.81 (t, 2 H), 2.58 (d, 2 H), 1.77 (d, 2 H), 1.40-1.69 (m, 10 H), 1.06-1.31 (m, 2 H); ESI MS $m/z = 456$ $[\text{C}_{27}\text{H}_{33}\text{N}_7+\text{H}]^+$.

Example 220 - Preparation of Compound 221

The compound 220 (100 mg, 0.170 mmol) was dissolved in methanol (25 mL). To the stirred solution was added ammonium formate (100 mg), and Pd/C (10.0 mg). After refluxing for 2 h, more ammonium formate (100 mg) and Pd/C (10.0 mg) were added. The reaction was cooled to room temperature and filtered through Celite. The filtrate was concentrated in vacuo. The resulting material was purified via silica gel chromatography (3:1:0.01 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$) to yield 221 (26.9 mg, 34%): ^1H NMR (300 MHz, CDCl_3) δ 7.29-7.63 (m, 10 H), 5.95 (brs, 1 H), 4.73-4.93 (m, 3 H), 4.64 (m, 1 H), 3.32 (t, 2 H), 3.10 (d, 2 H), 2.59 (t, 2 H), 1.76 (m, 2 H), 1.54 (d, 6 H), 1.28 (m, 3 H), 0.90 (m, 1 H); ESI MS $m/z = 456$ $[\text{C}_{27}\text{H}_{33}\text{N}_7+\text{H}]^+$.

Example 221 - Preparation of Compound 222

Prepared by the general methods described above: ^1H NMR (300 MHz, CDCl_3) δ 7.28-7.61 (m, 10 H), 6.20 (brs, 1 H), 4.70-4.91 (m, 3 H), 4.54-4.70 (m, 1 H), 4.28 (brs, 1 H), 3.75-3.90 (m, 1 H), 3.08-3.11 (m, 1 H), 2.80-2.93 (m, 1 H), 2.28-2.41 (d, 1 H), 1.95-2.10 (m, 1 H), 1.84-1.95 (m, 1 H), 1.70-1.84 (m, 1 H), 1.60-1.70 (m, 1 H), 1.52 (d, 6 H), 1.22-1.41 (m, 2 H), 0.94-1.22 (m, 2 H), 0.89 (t, 1 H); ESI MS $m/z = 456$ $[\text{C}_{27}\text{H}_{33}\text{N}_7+\text{H}]^+$.

Example 222 - Preparation of Compound 223

Prepared by the general Suzuki coupling conditions of 62 with boronic ester as shown in Scheme LXXV (58%): mp 200-206 °C; ¹H NMR (300 MHz, (CD₃)₂SO) δ 7.22-8.00 (m, 9 H), 6.10 (m, 2 H), 4.40-4.76 (m, 4 H), 3.63 (m, 1 H), 1.62-2.01 (m, 7 H), 1.35-1.60 (d, 6 H), 1.08-1.35 (m, 4 H); ESI MS *m/z* = 527 [C₃₀H₃₈N₈O+H]⁺.

Example 223 - Preparation of Compound 224

Prepared by the general Suzuki coupling conditions of 61 and 3,4-dimethylbenzeneboronic acid: ¹H NMR (300 MHz, CDCl₃) δ 7.53 (m, 3 H), 7.41 (d, 2 H), 7.26-7.38 (m, 2 H), 7.19 (d, 1 H), 4.79 (s, 2 H), 4.64 (m, 1 H), 3.80 (m, 1 H), 3.12 (m, 1 H), 2.10-2.36 (m, 10 H), 1.43-1.72 (m, 8 H), 1.27 (m, 4 H); ESI MS *m/z* = 484 [C₂₉H₃₇N₇+H]⁺.

Example 224 - Preparation of 5-Bromo-2-cyanopyridine

2,5-Dibromopyridine (20.0 g, 84.4 mmol) was dissolved in dimethylformamide (422 mL). To the stirred solution was added copper(I) cyanide. After refluxing for 5 h, the mixture was cooled to room temperature and stored overnight. The reaction mixture was diluted with ethyl acetate (1200 mL) and filtered through a Buchner funnel containing sand, Celite, and silica gel layers. The filtrate was concentrated to a volume of 400 mL. This organic liquid was diluted with water (300 mL) and the resulting liquid was extracted with ethyl acetate (2 x 200 mL). The organic extracts were combined, washed with water (2 x 300 mL) and brine (1 x 250 mL), and dried over magnesium sulfate. After concentration, the product was purified via silica gel chromatography (50:50 ethyl acetate/CH₂Cl₂) to afford the title compound (9.79 g).

Example 225 - Preparation of Compound 225

Prepared by reaction of 5-bromo-2-cyanopyridine with benzenboronic acid under standard Suzuki conditions (68%).

Example 226 - Preparation of Compound 226

In a Parr shaker vessel were combined 225 (300 mg, 1.67 mmol), glacial acetic acid (25 mL), and 10% palladium on carbon catalyst (177 mg, 0.167 mmol). The solution was agitated under 45 psig hydrogen gas for 2 h. The resulting dispersion was filtered through a Buchner funnel. The filtrate was concentrated. Purification by acid/base extraction yielded 226 (240 mg, 78%).

Example 227 - Preparation of Compound 229

Following the general schemes outlined above, compound 226 was transformed into 227 (57% yield). Compound 227 was then transformed into 228 in 83% yield. Compound 228 was then converted into compound 229 and then its HCl salt (75%): ^1H NMR (300 MHz, CD_3OD) δ 9.11 (s, 1 H), 8.90 (d, 2 H), 8.23 (d, 1 H), 7.83 (m, 2 H), 7.58 (m, 3 H), 5.23 (m, 2 H), 4.70-5.01 (m, 1 H), 3.72 (m, 1 H), 3.09 (m, 1 H), 1.80-2.15 (m, 4 H), 1.20-1.80 (m, 10 H); ESI MS m/z = 457 [$\text{C}_{26}\text{H}_{32}\text{N}_8 + \text{H}$] $^+$.

Example 228 - Preparation of Compound 230

In a flask immersed in an ice water bath were combined 187 (30.0 mg, 0.055 mmol), BOC-L-alanine (10.4 mg, 0.055 mmol), HATU (25.0 mg, 0.066 mmol), *N,N*-diisopropylethylamine (0.050 mL, 0.274 mmol), and dimethylformamide (0.500 mL) for 10 min then warmed to room temperature. After stirring overnight, the reaction mixture was diluted with methylene chloride (50 mL). The organic material was washed with 1 M citric acid (2 x 50 mL), saturated sodium bicarbonate solution (50 mL), and brine (50 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated in vacuo. To remove remaining dimethylformamide, the resulting material was dissolved in ethyl acetate (50 mL) and rinsed with 5% lithium chloride solution (3 x 50 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated in vacuo to yield 230 (28.0 mg, 79%): ^1H NMR (300 MHz, CDCl_3) δ 8.71 (s, 1 H), 7.62-7.83 (m, 4 H), 7.37-7.60 (m, 2 H), 7.11 (m, 1 H), 6.11 (brs, 1 H), 6.00 (d, 1 H), 5.01 (brs, 1 H), 4.81 (d, 2 H), 4.64 (m, 2 H), 4.08 (m, 1 H), 3.72 (m, 2 H), 2.12 (m, 2 H), 2.00 (m, 2 H), 1.54 (d, 6 H), 1.43 (s, 9 H), 1.15-1.38 (m, 7 H); ESI MS m/z = 646 [$\text{C}_{34}\text{H}_{44}\text{FN}_9\text{O}_3 + \text{H}$] $^+$.

Example 229 - Preparation of Compound 231

To a stirred solution of 230 in methylene chloride (2 mL) was added HCl in ethanol (2 mL). After stirring for 10 min, the solution was concentrated in vacuo to yield 231 (15.4 mg): ¹H NMR (300 MHz, CD₃OD) δ 8.92 (s, 1 H), 8.35-8.67 (m, 2 H), 8.32 (d, 1 H), 7.83 (t, 2 H), 7.69 (m, 1 H), 7.40 (m, 1 H), 4.65-5.20 (m, 3 H), 3.60-4.00 (m, 2 H), 1.80-2.30 (m, 4 H), 2.65 (d, 3 H), 1.08-1.58 (m, 12 H); ESI MS *m/z* = 546 [C₂₉H₃₆FN₉O+H]⁺.

Example 230 - Preparation of Compound 232

In a flask immersed in an ice water bath were combined 187 (30.0 mg, 0.055 mmol), BOC-glycine (9.6 mg, 0.055 mmol), HATU (25.0 mg, 0.066 mmol), *N,N*-diisopropylethylamine (0.05 mL, 0.274 mmol), and dimethylformamide (0.50 mL) for 10 min then warmed to room temperature. After stirring overnight, the reaction mixture was diluted with methylene chloride (50 mL). The organic material was washed with 1 M citric acid (2 x 50 mL), saturated sodium bicarbonate solution (50 mL), and brine (50 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated in vacuo. To remove remaining dimethylformamide, the resulting material was dissolved in ethyl acetate (50 mL) and rinsed with 5% lithium chloride solution (3 x 50 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated in vacuo to yield 232 (33.0 mg): ¹H NMR (300 MHz, CDCl₃) δ 8.65 (s, 1 H), 7.52-7.75 (m, 4 H), 7.44 (s, 1 H), 7.35 (q, 1 H), 7.01 (t, 1 H), 6.05 (brs, 1 H), 4.75 (d, 2 H), 4.57 (m, 2 H), 3.70 (m, 2 H), 2.05 (m, 2 H), 1.92 (m, 2 H), 1.46 (d, 6 H), 1.40 (s, 9 H), 1.20 (m, 6 H); ESI MS *m/z* = 632 [C₃₃H₄₂FN₉O₃+H]⁺.

Example 231 - Preparation of Compound 233

To a stirred solution of 232 in methylene chloride (2 mL) was added HCl in ethanol (2 mL). After stirring for 10 min, the solution was concentrated in vacuo to yield 233 (10.6 mg): ¹H NMR (300 MHz, CD₃OD) δ 8.95 (s, 1 H), 8.28-8.75 (m, 3 H), 7.79 (t, 2 H), 7.69 (m, 1 H), 7.46 (t, 1 H), 4.63-5.20 (m, 3 H), 3.59-3.92 (m, 2 H), 1.96-2.24 (m, 4 H), 1.62 (d, 6 H), 1.05-1.52 (m, 8 H); ESI MS *m/z* = 532 [C₂₈H₃₄FN₉O+H]⁺.

Example 232 - Preparation of Compound 239

Reaction of 234 with 1 under standard conditions provides 236 (90%).
Reaction of 236 with *trans*-1,4-cyclohexanediamine provides 237 (95%). Boc
5 protection of 237 followed by Suzuki coupling provides 238 in 50% yield. Compound
238 was added to a 1:1 mixture of methylene chloride and trifluoroacetic acid. After
stirring for 2 h, the solution was concentrated in vacuo. The resulting material was
purified via silica gel chromatography (94:5:1 CH₂Cl₂:MeOH:NH₄OH) to afford 239:
¹H NMR (300 MHz, CDCl₃) δ 7.51-7.61 (m, 4 H), 7.48 (s, 1 H), 7.41 (t, 2 H), 7.31
10 (m, 3 H), 5.61 (m, 1 H), 4.63 (m, 2 H), 3.87 (m, 2 H), 3.18 (m, 1 H), 3.00 (t, 2 H),
2.20-2.35 (m, 3 H), 1.72 (m, 4 H), 1.52 (d, 6 H), 1.25 (m, 4 H).

Example 233 - Preparation of Compound 240

15 Compound 239 was acetylated under the general conditions described
above to provide 240 (73%). Salt formation occurred in 71% yield: ¹H NMR (300
MHz, CDCl₃) δ 7.28-7.62 (m, 10 H), 5.67 (m, 1 H), 4.64 (m, 2 H), 3.68-3.97 (m, 3
H), 3.00 (t, 2 H), 2.23 (m, 2 H), 1.84-2.11 (m, 7 H), 1.53 (d, 6 H), 1.30 (m, 4 H).

Example 234 - Preparation of Compound 241

20 Reductive amination of 239 with propionaldehyde followed by salt
formation provided 241: ESI MS *m/z* = 512 [C₃₁H₄₁N₇+H]⁺.

Example 235 - Preparation of Compound 242

25 Compound 237 was Boc-protected and then treated with 3-
thiopheneboronic acid under standard Suzuki condition to prepare 242: ¹H NMR (300
MHz, CDCl₃) δ 7.33-7.76 (m, 6 H), 7.28 (m, 2 H), 5.72 (brs, 1 H), 4.64 (m, 1 H), 4.43
30 (m, 1 H), 3.83 (m, 2 H), 3.47 (m, 1 H), 2.97 (t, 2 H), 2.21 (m, 2 H), 2.08 (m, 2 H),
1.53 (d, 6 H), 1.46 (s, 9 H), 1.29 (m, 4 H); ESI MS *m/z* = 576 [C₃₁H₄₁N₇O₂S+H]⁺.

Example 236 - Preparation of Compound 243

35 Compound 242 was deprotected with HCl in methanol to provide 243:
¹H NMR (500 MHz, CDCl₃) δ 7.68 (m, 1 H), 7.54 (d, 2 H), 7.47 (s, 1 H), 7.42 (s, 1

- 211 -

H), 7.37 (s, 1 H), 7.28 (m, 2 H), 5.68 (brs, 1 H), 4.63 (m, 2 H), 3.85 (m, 3 H), 2.99 (t, 2 H), 2.71 (m, 1 H), 2.18 (d, 2 H), 1.90 (d, 2 H), 1.52 (d, 6 H), 1.25 (m, 4 H).

Example 237 - Preparation of Compound 245

5

To a stirred solution of sodium hydride (423 mg, 17.6 mmol) in tetrahydrofuran (12 mL), was added 4-phenylphenol (2.00 g, 11.8 mmol). After 1 h, BOC-2-aminoethylbromide (3.90 g, 17.6 mmol) was added to the solution. After stirring overnight, the reaction mixture was quenched with 2 N potassium hydroxide solution (10 mL). The resulting mixture was extracted with methylene chloride (12 mL). The organic layer was concentrated and the crude material was purified via silica gel chromatography to yield 245.

Example 238 - Preparation of Compound 246

15

The protected amine 245 was added to 10 mL of an 1:1 mixture of methylene chloride and trifluoroacetic acid. After concentration, the material was diluted with 2 N potassium hydroxide solution (10 mL). The aqueous layer was extracted with methylene chloride (2 x 10 mL). The organic extracts were combined, dried over magnesium sulfate, and concentrated in vacuo to afford the product (400 mg). Reaction with 1 under standard conditions provided 246 (91%).

Example 239 - Preparation of Compound 248

25

Compound 246 was transformed into 247 under standard conditions (80%). Reaction of 247 with *trans*-1,4-cyclohexanediamine provided 248. Salt formation provided the target compound (68%): ¹H NMR (500 MHz, CDCl₃) δ 7.53 (m, 4 H), 7.40 (m, 2 H), 7.29 (m, 2 H), 7.00 (d, 2 H), 5.93 (brs, 1 H), 4.61 (m, 2 H), 4.22 (t, 2 H), 4.02 (m, 2 H), 3.78 (m, 1 H), 2.70 (m, 1 H), 2.18 (d, 2 H), 1.90 (d, 2 H), 1.53 (d, 6 H), 1.25 (m, 4 H).

30

- 212 -

Example 240 - Preparation of Compound 250

Reductive amination of 248 with propionaldehyde and salt formation under standard conditions described above provided 250: ESI MS $m/z = 528$

5 $[C_{31}H_{41}N_7O+H]^+$.

Example 241 - Preparation of Compound 249

N-Acetylation of 248 and salt formation under standard conditions
10 provided 249: 1H NMR (300 MHz, $CDCl_3$) δ 6.95-7.60 (m, 10 H), 5.97 (brs, 1 H), 5.24 (d, 1 H), 4.63 (m, 2 H), 4.23 (t, 2 H), 4.02 (m, 2 H), 3.78 (m, 2 H), 2.21 (m, 2 H), 2.04 (m, 2 H), 1.94 (s, 3 H), 1.55 (d, 6 H), 1.30 (m, 4 H).

Example 242 - Preparation of Compound 255

15 Utilizing reaction conditions described in general above, 251 was converted to 252 (100%). Compound 252 was converted to 253 then 254 and then Boc-protected to make 255 (21%).

Example 243 - Preparation of Compound 256

Compound 255 was treated with phenylboronic acid under standard Suzuki condions. The product was dissolved in methanol and immersed in an ice water bath. Hydrogen chloride gas was bubbled through the solution. The solution
25 was concentrated in vacuo and the resulting material was purified via preparatory HPLC (acetonitrile/water/trifluoroacetic acid) to yield 256 (8 mg).

Example 244 - Preparation of Compound 257

30 Compound 255 was treated with 3-thiopheneboronic acid under standard Suzuki conditions. The product was dissolved in methanol and immersed in an ice water bath. Hydrogen chloride gas was bubbled through the solution. The solution was concentrated in vacuo and the resulting material was purified via preparatory HPLC (acetonitrile/water/trifluoroacetic acid) to yield 257: 1H NMR
35 (300 MHz, $CDCl_3$) δ 7.31-7.58 (m, 8 H), 5.99 (brs, 1 H), 5.10-5.50 (m, 1 H), 4.49-

- 213 -

4.69 (m, 2 H), 3.67 (m 1 H), 2.67 (m, 1 H), 2.13 (m, 2 H), 1.90 (m, 2 H), 1.62 (d, 3 H), 1.53 (d, 6 H), 1.21 (m, 4 H).

Example 245 - Preparation of Compound 258

5

Reaction of 75 with propionoyl chloride under standard conditions provides 258 (89%): mp 182-183 °C

Example 246 - Preparation of Compound 259

10

Reaction of 75 with methyl chloroformate under standard conditions provides 259 (68%): mp 148-150 °C.

Example 247 - Preparation of Compound 260

15

Reaction of 75 with methanesulfonyl chloride under standard conditions provides 260 (56%): mp 143-145 °C.

Example 248 - Preparation of Compound 261

20

Reaction of 75 with cyclopropanecarbonyl chloride under standard conditions provides 261 (87%): mp 196-204 °C.

Example 249 - Preparation of Compound 262

25

Compound 75 (250 mg, 0.549 mmol) and succinic anhydride (60.0 mg, 0.600 mmol) were dissolved in xylene (30 mL). A few drops of dimethylformamide were added to the solution. After refluxing for 48 h, the mixture was concentrated in vacuo. The resulting material was purified via silica gel chromatography (99.5:0.5 CH₂Cl₂/MeOH) and recrystallized from CH₂Cl₂ in hexanes (1:10) to yield 262 (30.0 mg, 10%): mp 141-147 °C.

30

Example 250 - Preparation of Compound 263

35

The amine 75 (200 mg, 0.439 mmol) was dissolved in methylene chloride (15 mL). The stirred solution was cooled to -78 °C and *N,N*-diisopropylethylamine (113 mg, 0.878 mmol) and trifluoromethylsulfonylchloride

- 214 -

(81.4 mg, 0.483 mmol) were added. After 30 min, the solution was warmed to room temperature. The mixture was cooled to -78 °C and another 1.10 equivalents of trifluoromethylsulfonylchloride and 1.50 equivalents of *N,N*-diisopropylethylamine were added. After warming to room temperature, the solution was concentrated. The
5 resulting material was purified via silica gel chromatography (99:1 CH₂Cl₂/MeOH) and recrystallization from ether in hexanes to afford **263** (60 mg, 23%): mp 131-136 °C.

Example 251 - Preparation of Compound 264

10

Prepared by standard Suzuki coupling of **61** to provide **264** (65%): mp 186-190 °C.

Example 252 - Preparation of Compound 265

15

N-Acetylation of **264** under standard conditions provides **265** (37%): mp 241-246 °C.

Example 253 - Preparation of Compound 266

20

Suzuki coupling of **61** with 2-chlorobenzenboronic acid provides **266** (13%): API MS *m/z* = 490 [C₂₇H₃₂ClN₇+H]⁺.

Example 254 - Description of Biological Assays

25

A. Immunopurification of CyclinA/cdk2 and CyclinE/cdk2 Complexes.

CyclinA/cdk2 and cyclinE/cdk2 assays were carried out with cyclin/cdk complexes isolated from HeLa S-3 suspension cultures. HeLa cells were
30 grown in spinner flasks at 37 °C in Joklik's modified minimum essential media (MEM) supplemented with 7% horse serum. After growing in medium supplemented with 2 mM thymidine for 16-18 h, cultures were arrested at the G1/S border and cyclinA/cdk2 and cyclinE/cdk2 were isolated from cell lysates by immunoprecipitation with antibodies specifically directed against each cyclin subunit.
35 Rabbit anti-cyclinA (H-432) and the mouse monoclonal antibody against cyclinE

(HE111) were purchased from Santa Cruz Biotechnology. Cells blocked at the appropriate stage of the cell cycle were disrupted in lysis buffer (50 mM Tris, pH 8.0, 250 mM NaCl, 0.5% NP-40 plus protease and phosphatase inhibitors) and centrifuged at 10,000 x g to remove insoluble material. To isolate cyclin/cdk complexes, 1 µg of anti-cyclin antibody was incubated with lysate from 1×10^7 cells for 1 h at 4 °C. Protein A-coated agarose beads were then added for 1 h to collect antibody-bound immune complexes. The immobilized cyclin/cdk complexes were then washed 4 x with lysis buffer to reduce nonspecific protein binding. The complexes were then washed 1 x in kinase assay buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 1 mM DTT) and aliquoted into individual assay tubes.

B. Immunopurification of CyclinB/cdk1 Complex.

HeLa cells are blocked at the G1/S border by culturing in the presence of 2 mM thymidine for 20 h. The cells are then rinsed 3x in phosphate buffered saline and resuspended in regular medium. After 4 h of culture, the mitotic blocker, nocodazole is added to a final concentration of 75 ng/ml. Sixteen hours later, the cells are harvested by centrifugation, washed in PBS, and lysed in cold Lysis Buffer (50 mM Tris pH 8.0, 250 mM NaCl, 0.5% NP-40, 1 mM DTT, 25 µg/ml leupeptin, 25 µg/ml aprotinin, 15 µg/ml benzamidin, 1 mM PMSF, 50 mM sodium fluoride, 1 mM sodium orthovanadate) for 15 min at 1×10^7 cells/ml. The lysate is then clarified by centrifugation at 10,000 x g for 10 min. The supernatant is collected and diluted 1:5 with Lysis Buffer. Monoclonal antibody against cyclinB (GNS1) is added to the supernatant to a final concentration of 5 µg/ml and shaken at 4 °C for 2 h. The immune complexes are then collected by the addition of 200 µl of protein agarose beads for 1 h. The beads are washed 4x in lysis buffer and 1x in kinase assay buffer.

C. Protein Kinase Assays and Determination of IC₅₀ Values.

CyclinA/cdk2 assays were carried out with complexes isolated from 0.5×10^6 cells. CyclinE/cdk2 assays were carried out with complexes isolated from 4×10^6 cells. CyclinB/cdk1 assays were carried out with complexes isolated from 4×10^4 cells. After centrifugation, the wash buffer was removed and the complexes resuspended in 15 µl of kinase assay buffer (kinase wash buffer + 167 µg/ml histone H1). Compounds being tested for inhibition were added prior to the addition of [³²P] ATP to a final concentration of 15 µM. The tubes were incubated at 30 °C for 5

- 216 -

min and the reactions were stopped by the addition of an equal volume of 2 x SDS-PAGE sample buffer. The samples were then subjected to electrophoresis on 10% SDS-PAGE to resolve the histone H1 from other reaction components. The amount of radioactive phosphate transferred to histone H1 was quantified on a Storm
5 Phosphorimager (Molecular Dynamics).

Prior to the protein kinase assay, test compounds were dissolved in DMSO at a concentration of 25 mM and were diluted to produce final concentrations of 0.1, 1.0, and 10.0 μ M in the kinase assays. To eliminate possible effects of differences in DMSO concentration, the DMSO was kept constant at 0.04%, including
10 the control reaction. Duplicate assays were performed at each concentration. The activity was plotted as the percent of activity in the absence of added test compound versus test compound concentration. IC₅₀ values were calculated using GraphPad Prism data analysis software.

D. Measuring the Inhibition of Cell Growth.

15 Growth inhibition (GI₅₀) values were measured with HeLa S-3 cells selected for growth on plastic. The procedure was based on the protocol of Skehan *et al.* (Skehan, P., et al., J. Natl. Cancer Inst., 82:1107-1112 (1990), which is hereby incorporated by reference) HeLa cells were plated at 2×10^4 cells/well in 96 well plates. One day later, a control plate was fixed by addition of TCA to 5%. After five
20 rinses with tap water the plate was air dried and stored at 4 °C. Test compounds were added to the remaining plates at 10-fold dilutions between 0.01 and 100 μ M. Two days later all plates were fixed as described above. Cells were then stained by the addition of 100 μ l per well of 0.4% sulforhodamine B (SRB) in 1% acetic acid for 30 min at 4 °C. Wells were then quickly rinsed 5 x with acetic acid (1%) and allowed to
25 air dry. The SRB was then solubilized by the addition of 100 μ l per well of unbuffered 10 mM Tris base. Dye was quantified by measuring absorbance at 490 nm on a Molecular Devices kinetic microplate reader. Growth at each inhibitor concentration relative to the untreated control was calculated according to the following equation: percent growth = $100 \times (T - T_0) / (C - T_0)$, where T was the average
30 optical density (OD) of the test wells after 2 days of treatment, T₀ was the average OD of the wells in the control plate on day 0 and C was the average OD of untreated

- 217 -

wells. Plots of percent growth versus inhibitor concentration were used to determine the GI_{50} .

The data below shown in Table 2 summarizes the *in vitro* cyclin/cdk inhibition constants (IC_{50}) and growth inhibition constants (GI_{50}) of HeLa Cells for
5 the compounds of the current invention. Replicate experimental results are summarized below.

Table 2: In Vitro Cyclin/cdk Inhibition (IC₅₀) and Growth Inhibition (GI₅₀) of HeLa Cells For Compounds of the Current Invention.

Compound	IC ₅₀ CyclinA/cdk2 (μ M)	IC ₅₀ CyclinE/cdk2 (μ M)	IC ₅₀ CyclinB/cdk1 (μ M)	GI ₅₀ HeLa Cells (μ M)
5	> 10	12	7	5
	0.4	0.6		> 10
12	2	1	3	0.06
	0.7	3		0.003
	0.9	0.5		0.001
	0.2	0.1		0.02
13	4	2	4	3
	1	0.3		2
	0.8	0.9		
14	3	0.4	7	0.4
	3	2		0.03
17				0.03
	1	1	10	0.4
	2	0.9	3	0.6
	1	0.2	11	0.25
	> 10	9		0.4
	10	2		0.3
25				0.4
	1	4	> 10	2
	6	1	> 10	0.4
32	> 10	9		> 1
33	2	3	--	5
	5	0.9		0.7
34	> 10	4	> 10	1
	13	6		2
	8			0.9
36	12	5	> 10	7
	13	2		6
				7
38	> 10	> 10	> 10	20
	> 10	> 10		20
	> 10			> 10
40	> 10	> 10	> 10	0.6
	> 10	> 10		1
				0.6
43	> 10	> 10	> 10	9
	> 10	> 10		25
				> 10
46	> 10	> 10	> 10	4
	> 10	> 10		4
				8
46	> 10	6	> 10	25
	8	3		> 10

- 219 -

48	22 6	1 5	> 10	0.3 0.6 0.5
50	> 10 7	> 10 9	> 10	3 > 10
53	> 10 > 10	15 4	> 10	0.2 0.3 0.5
58	11 4	2 4	12	2 0.5 0.7
60	> 10 0.4	12 > 10	> 10	7 6
73	> 50 14 > 10 > 10	4 12 > 10 > 10	> 10	0.3 0.5 0.3 0.5
74	5 2 1	2 3 2	6	0.2 0.01 0.05 0.03 0.05
75	3	3	6	0.09 0.02 0.005
76	12 11 3	3 5 2	6	0.07 0.01 0.06 0.2 0.04
77	> 10 > 10	4 14	> 10	0.15 0.5 0.3
78	0.9 0.9 0.7	0.6 0.3 0.2	0.8 0.8	0.05 0.025 0.08 0.002
79	10 0.5 1	2 0.1 0.08	3	0.07 0.007 0.004 0.4
80	> 10 > 10	> 10 4 2	> 10	> 100 > 10
86	0.9 0.7 0.4 0.6	0.4 0.2 0.4 0.03	2	0.2 0.03 0.01 0.01 0.2

- 220 -

87	4 2 0.5	1 0.3 0.1	5	0.07 0.01 0.004 0.006 0.03 0.006 0.001 0.0001
88	3 > 10 2	4 > 10 5	> 10	0.1 0.05 0.04 0.005
93	0.2 0.3	0.09 0.1	0.9	0.3 0.08 0.3
94	0.6 0.2	0.3 0.3	0.4	0.1 0.07 0.4
95	1 2	1 0.7	4	0.08 0.003 0.0005
96	8	4	6	0.04 0.01
97	> 10	3	10	3
98	6 2	2 2	> 10	> 10 11
99	> 10	9	> 10	5
100	> 10	4	> 10	0.6
101	3 0.9	1 0.7	4	1 1
102	> 10	4	--	4
103	0.6 0.7	0.2 0.2	1	0.03 0.008 0.02 0.01
104	7 8	1 1	2	0.4 0.2
106	11 4	3 1	--	0.3 0.1
107	1 4	2	--	0.4 0.3
108	10 > 10	> 10 > 10	--	3 5
109	0.6	0.1	--	0.04 < 0.0001
110	0.6	2	--	0.02 0.03 0.02 0.01
111	0.2	0.07	--	0.02 0.0006

- 221 -

112	2	2	--	< 0.001 0.002 0.02 0.006 0.0006
113	0.4	0.3	--	< 0.001 0.00001 0.03 0.001 0.02
114	3	0.7	--	> 10
115	3	0.4	--	3
116	> 10	> 10	--	> 10 > 10
117	> 10	3	--	3
118	6	1	--	> 10 > 10
123	0.2	0.04	--	< 0.001 < 0.001 0.0001
124	2	0.8	--	0.003 < 0.001 < 0.0001
130	--	--	--	> 10 > 10
131	--	--	--	3 2
132	--	--	--	4 3
133	--	--	--	> 10 > 10
134	--	--	--	2 3
135	--	--	--	4 3
137	--	--	--	0.05 0.06 0.05
139	--	--	--	0.2 0.07
140	--	--	--	1 2
142	--	--	--	0.4 0.5
144	--	--	--	0.4 0.4
146	--	--	--	0.7 0.3
148	--	--	--	1 1
149	--	--	--	0.3

- 222 -

				0.2
150	--	--	--	0.3
				0.2
151	--	--	--	0.8
				0.6
152	--	--	--	0.7
				0.3
153	--	--	--	3
				2
154	--	--	--	0.6
				0.9
155	--	--	--	0.5
				0.8
156	--	--	--	3
				2
157	--	--	--	0.4
				0.5
158	--	--	--	0.6
				0.4
159	--	--	--	4
				3
160	--	--	--	0.2
				0.3
161	--	--	--	0.2
				0.4
162	--	--	--	0.2
				0.3
163	--	--	--	2
				3
164	--	--	--	0.2
				0.1
165	--	--	--	0.2
				0.1
166	--	--	--	4
				2
167	--	--	--	2
				0.9
168	--	--	--	4
				3
169	--	--	--	0.5
				0.3
170	--	--	--	4
				2
171	--	--	--	3
				3
172	--	--	--	0.3
				0.3
173	--	--	--	3
				3
174	--	--	--	0.04
				0.03

- 223 -

				0.1 0.06 0.4 0.4
175	--	--	--	0.6 0.3
177	--	--	--	0.2 0.06 0.06
178	--	--	--	0.4 0.2
179	--	--	--	0.1 0.05 0.05
180	--	--	--	0.4 0.3
181	--	--	--	0.04
182	--	--	--	0.3 0.3
187	--	--	--	0.05 0.03
188	--	--	--	0.2 0.07
194	--	--	--	0.06 0.04
199	--	--	--	0.2 0.09
200	--	--	--	0.3 0.2
206	--	--	--	0.2 0.2
207	--	--	--	0.4 0.2
208	--	--	--	4 3
209	--	--	--	2 2
210	--	--	--	3 4
211	--	--	--	0.6 0.3
212	--	--	--	5 3
213	--	--	--	3 2
214	--	--	--	5 5
215	--	--	--	2 3
216	--	--	--	0.5 0.5

- 224 -

217	--	--	--	4 4
218	--	--	--	3 5
219	--	--	--	0.4 0.6
221	--	--	--	2 2
222	--	--	--	1 2
223	--	--	--	0.04 0.1
224	--	--	--	2 2
229	--	--	--	0.4
230	--	--	--	0.3
231	--	--	--	0.04
232	--	--	--	0.3
233	--	--	--	0.5
239	--	--	--	4 6
240	--	--	--	8 8
241	--	--	--	7 4
242	--	--	--	7 >10
243	--	--	--	3 3
248	--	--	--	3 4
249	--	--	--	>10 >10
250	--	--	--	3 6
256	--	--	--	4 3
257	--	--	--	3 3
258	--	--	--	0.2 0.3 0.4
259	--	--	--	0.3 0.4 0.7
260	--	--	--	0.2 0.1 0.2
261	--	--	--	0.3 0.3 0.3





- 225 -

262	--	--	--	0.3 0.2 0.5
263	--	--	--	2 3 4
264	--	--	--	0.3 0.3 0.5
265	--	--	--	0.3 0.3 0.4
266	--	--	--	0.3 0.3 0.5
267	--	--	--	0.8 0.6





The data below shown in Table 3 summarizes the in vitro cyclin/cdk inhibition (IC₅₀) and growth inhibition (GI₅₀) of HeLa Cells for several reference compounds in comparison to several compounds of the current invention. The chemical structures are provided.

5

Table 3: In Vitro cyclin/cdk Inhibition (IC₅₀) and Growth Inhibition (GI₅₀) of HeLa Cells For Reference Compounds in Comparison to Several Compounds of the Current Invention.

Compound	Structure	IC ₅₀ CyclinA/cdk2 (μ M)	IC ₅₀ CyclinE/cdk2 (μ M)	IC ₅₀ CyclinB/cdk1 (μ M)	GI ₅₀ HeLa Cells (μ M)
Olomoucine		0.5-24 (n > 10)	1-14 (n > 10)	7-23 (n > 10)	75
Roscovitine		2.1 4 3	0.04 0.7	—	30 25 30 > 10 25
Flavopiridol		0.06 0.2	0.6 0.04	0.06 (n = 2)	0.18
125		1	0.1	0.6	3

- 227 -

126		0.6 0.8	0.06 0.06	2 0.2	2 4 6
74		5	2	6	0.2 0.01 0.05
127		0.3-2 (n > 15)	0.04-0.07 (n > 15)	0.5-2 (n > 15)	7-15 (n > 5)
88		3	4	> 10	0.1 0.05 0.04

The following data in Tables 4, 5, 6, and 7 summarize the growth inhibition properties of several compounds of the current invention and olomoucine against 60-human transformed cell lines. These data were cooperatively obtained at the National Cancer Institute in their 60-cell line growth inhibition assay according to published procedures (Boyd, M.R., "Anticancer Drug Development Guide," Preclinical Screening, Clinical Trials, and Approval;

Teicher, B. Ed.; Humana Press; Totowa, NJ, 23-42 (1997), which is hereby incorporated by reference).

Table 4: In Vitro Growth Inhibition (GI₅₀) of NCI Human Transformed Cell Lines of Several Compounds of the Current Invention.

Cancer Type	Cell Line	73 GI ₅₀ (μM)	17 GI ₅₀ (μM)	33 GI ₅₀ (μM)	38 GI ₅₀ (μM)
Breast	BT-549	0.25	0.40	51.3	0.32
Breast	HS 578T	0.10	6.31	--	--
Breast	MCF7	0.16	0.16	5.2	0.20
Breast	MDA-MB-231/ATCC	0.50	--	--	0.06
Breast	MDA-MB-435	0.25	0.20	4.9	0.05
Breast	MDA-N	0.13	0.11	--	--
Breast	NCI/ADR-RES	0.40	0.28	6.3	0.32
Breast	T-47D	0.25	0.13	3.9	0.25
CNS	SF-268	0.16	0.04	6.3	0.20
CNS	SF-295	0.25	0.19	7.8	0.50
CNS	SF-539	0.76	0.40	89.1	1.26
CNS	SNB-19	0.43	0.14	38.0	0.50
CNS	SNB-75	0.02	0.02	--	--
CNS	U251	0.32	0.40	3.7	0.20
Colon	COLO 205	0.28	0.05	7.8	0.16
Colon	HCC-2998	0.20	0.03	> 1000	7.94
Colon	HCT-116	0.20	0.16	6.2	0.32
Colon	HCT-15	0.18	0.04	8.9	0.25
Colon	HT29	--	0.10	8.9	0.25
Colon	KM12	0.13	0.03	4.1	0.16
Colon	SW-620	--	0.01	2.9	0.03
Leukemia	CCRF-CEM	0.25	0.16	4.6	0.20
Leukemia	HL-60(TB)	--	--	3.2	0.04
Leukemia	K-562	0.16	0.16	3.1	0.25
Leukemia	MOLT-4	0.32	0.25	3.8	0.25
Leukemia	RPMI-8226	0.03	0.03	1.5	--
Leukemia	SR	--	0.50	4.5	3.98
Melanoma	LOX IMVI	--	0.32	16.6	0.40
Melanoma	M14	0.03	0.03	7.8	0.05
Melanoma	MALME-3M	0.27	19.95	11.7	0.25
Melanoma	SK-MEL-2	0.63	1.00	> 1000	2.00
Melanoma	SK-MEL-28	0.45	0.12	5.9	0.03
Melanoma	SK-MEL-5	0.25	0.32	16.2	0.32
Melanoma	UACC-257	0.16	0.20	75.9	0.50
Melanoma	UACC-62	0.30	0.27	8.3	1.00
Non-Small Cell Lung	A549/ATCC	0.03	0.03	4.6	0.13
Non-Small Cell Lung	EKVX	0.25	2.51	6.9	0.20
Non-Small Cell Lung	HOP-62	0.06	0.20	> 1000	0.32

- 229 -

Non-Small Cell Lung	HOP-92	1.00	1.58	--	0.32
Non-Small Cell Lung	NCI-H226	0.22	0.11	--	--
Non-Small Cell Lung	NCI-H23	0.32	0.16	26.3	0.32
Non-Small Cell Lung	NCI-H322M	0.16	> 1000	38.9	0.40
Non-Small Cell Lung	NCI-H460	0.40	0.41	25.7	3.16
Non-Small Cell Lung	NCI-H522	--	--	4.2	--
Ovarian	IGROV1	0.32	0.20	10.0	0.16
Ovarian	OVCAR-3	0.30	0.65	> 1000	1.00
Ovarian	OVCAR-4	0.32	0.32	31.6	1.26
Ovarian	OVCAR-5	0.25	0.26	> 1000	0.40
Ovarian	OVCAR-8	--	0.13	6.6	0.25
Ovarian	SK-OV-3	0.95	0.40	> 1000	3.98
Prostate	DU-145	7.08	0.63	17.8	1.26
Prostate	PC-3	0.35	0.20	> 1000	0.40
Renal	786-0	0.20	0.25	18.6	0.32
Renal	A498	2.88	1.58	--	1.26
Renal	ACHN	0.32	0.40	5.2	2.00
Renal	CAKI-1	1.66	0.13	4.4	0.20
Renal	RXF 393	0.09	0.02	13.2	0.13
Renal	SN12C	--	0.56	--	--
Renal	TK-10	--	--	8.3	0.40
Renal	UO-31	0.06	0.10	8.1	0.13

Table 5: In Vitro Growth Inhibition (GI₅₀) of NCI Human Transformed Cell Lines of Several Compounds of the Current Invention.

5

Cancer Type	Cell Line	43 GI ₅₀ (μM)	48 GI ₅₀ (μM)	75 GI ₅₀ (μM)	76 GI ₅₀ (μM)
Breast	BT-549	4.0	0.01	< 0.01	< 0.01
Breast	HS 578T	--	0.03	< 0.01	< 0.01
Breast	MCF7	2.7	0.25	< 0.01	< 0.01
Breast	MDA-MB-231/ATCC	3.2	0.09	< 0.01	< 0.01
Breast	MDA-MB-435	2.1	--	--	--
Breast	MDA-N	--	0.02	< 0.01	< 0.01
Breast	NCI/ADR-RES	5.2	0.12	0.48	0.015
Breast	T-47D	2.2	0.15	< 0.01	< 0.01
CNS	SF-268	3.0	< 0.01	< 0.01	< 0.01
CNS	SF-295	4.0	0.24	< 0.01	< 0.01
CNS	SF-539	3.4	0.38	0.02	0.054
CNS	SNB-19	5.0	0.02	< 0.01	< 0.01
CNS	SNB-75	--	< 0.01	< 0.01	< 0.01
CNS	U251	2.3	0.17	< 0.01	0.020
Colon	COLO 205	1.6	0.03	< 0.01	< 0.01
Colon	HCC-2998	3.4	--	--	--
Colon	HCT-116	2.1	0.19	< 0.01	0.014

- 230 -

Colon	HCT-15	3.9	0.02	0.03	< 0.01
Colon	HT29	3.6	< 0.01	< 0.01	< 0.01
Colon	KM12	2.3	0.02	< 0.01	< 0.01
Colon	SW-620	1.6	< 0.01	< 0.01	< 0.01
Leukemia	CCRF-CEM	2.8	0.03	< 0.01	< 0.01
Leukemia	HL-60(TB)	2.1	--	--	--
Leukemia	K-562	3.1	0.16	< 0.01	< 0.01
Leukemia	MOLT-4	2.0	0.05	< 0.01	< 0.01
Leukemia	RPMI-8226	--	< 0.01	< 0.01	< 0.01
Leukemia	SR	2.2	0.16	< 0.01	< 0.01
Melanoma	LOX IMVI	3.4	0.19	< 0.01	< 0.01
Melanoma	M14	2.2	< 0.01	< 0.01	< 0.01
Melanoma	MALME-3M	3.0	0.13	< 0.01	< 0.01
Melanoma	SK-MEL-2	61.7	0.48	0.02	0.112
Melanoma	SK-MEL-28	2.3	< 0.01	< 0.01	< 0.01
Melanoma	SK-MEL-5	2.1	0.17	0.01	0.013
Melanoma	UACC-257	4.8	0.04	< 0.01	< 0.01
Melanoma	UACC-62	3.3	0.10	0.01	0.018
Non-Small Cell Lung	A549/ATCC	4.1	< 0.01	< 0.01	< 0.01
Non-Small Cell Lung	EKVX	2.8	--	--	--
Non-Small Cell Lung	HOP-62	3.3	0.03	< 0.01	< 0.01
Non-Small Cell Lung	HOP-92	2.6	0.46	< 0.01	0.017
Non-Small Cell Lung	NCI-H226	--	--	--	--
Non-Small Cell Lung	NCI-H23	4.3	0.07	< 0.01	< 0.01
Non-Small Cell Lung	NCI-H322M	3.5	0.03	< 0.01	< 0.01
Non-Small Cell Lung	NCI-H460	3.2	0.25	< 0.01	0.047
Non-Small Cell Lung	NCI-H522	--	< 0.01	< 0.01	< 0.01
Ovarian	IGROV1	3.4	0.23	< 0.01	< 0.01
Ovarian	OVCAR-3	9.3	0.17	< 0.01	< 0.01
Ovarian	OVCAR-4	8.9	0.20	< 0.01	< 0.01
Ovarian	OVCAR-5	3.6	0.16	< 0.01	< 0.01
Ovarian	OVCAR-8	3.9	0.10	< 0.01	< 0.01
Ovarian	SK-OV-3	72.4	1.38	0.03	0.051
Prostate	DU-145	2.6	0.55	< 0.01	0.043
Prostate	PC-3	38.9	0.23	< 0.01	< 0.01
Renal	786-0	3.1	0.25	< 0.01	< 0.01
Renal	A498	3.0	0.39	0.01	< 0.01
Renal	ACHN	3.1	0.25	0.02	0.025
Renal	CAKI-1	3.0	--	--	--
Renal	RXF 393	1.9	< 0.01	< 0.01	< 0.01
Renal	SN12C	--	0.03	< 0.01	< 0.01
Renal	TK-10	3.2	0.37	< 0.01	0.013
Renal	UO-31	2.8	< 0.01	0.03	< 0.01

Table 6: In Vitro Growth Inhibition (GI₅₀) of NCI Human Transformed Cell Lines of Several Compounds of the Current Invention.

Cancer Type	Cell Line	79 GI ₅₀ (μM)	87 GI ₅₀ (μM)	12 GI ₅₀ (μM)
Breast	BT-549	< 0.01	0.02	0.041
Breast	HS 578T	< 0.01	< 0.01	< 0.005
Breast	MCF7	< 0.01	0.04	< 0.005
Breast	MDA-MB-231/ATCC	< 0.01	< 0.01	< 0.005
Breast	MDA-MB-435	< 0.01	< 0.01	< 0.005
Breast	MDA-N	< 0.01	0.014	< 0.005
Breast	NCI/ADR-RES	0.86	0.28	1.26
Breast	T-47D	< 0.01	0.048	0.0088
CNS	SF-268	< 0.01	< 0.01	< 0.005
CNS	SF-295	< 0.01	0.047	0.018
CNS	SF-539	< 0.01	0.081	0.022
CNS	SNB-19	< 0.01	0.038	0.016
CNS	SNB-75	< 0.01	0.012	< 0.005
CNS	U251	< 0.01	0.028	0.0078
Colon	COLO 205	< 0.01	< 0.01	< 0.005
Colon	HCC-2998	< 0.01	< 0.01	< 0.005
Colon	HCT-116	< 0.01	0.037	0.0089
Colon	HCT-15	< 0.01	0.066	0.17
Colon	HT29	< 0.01	< 0.01	< 0.005
Colon	KM12	< 0.01	< 0.01	< 0.005
Colon	SW-620	< 0.01	< 0.01	< 0.005
Leukemia	CCRF-CEM	< 0.01	< 0.01	< 0.005
Leukemia	HL-60(TB)	< 0.01	< 0.01	< 0.005
Leukemia	K-562	< 0.01	0.024	< 0.005
Leukemia	MOLT-4	< 0.01	0.02	< 0.005
Leukemia	RPMI-8226	< 0.01	< 0.01	< 0.005
Leukemia	SR	< 0.01	0.032	< 0.005
Melanoma	LOX IMVI	< 0.01	0.027	< 0.005
Melanoma	M14	< 0.01	< 0.01	< 0.005
Melanoma	MALME-3M	< 0.01	0.024	0.010
Melanoma	SK-MEL-2	< 0.01	0.056	0.0096
Melanoma	SK-MEL-28	< 0.01	< 0.01	0.01
Melanoma	SK-MEL-5	< 0.01	0.028	0.014
Melanoma	UACC-257	< 0.01	0.017	0.008
Melanoma	UACC-62	< 0.01	0.045	0.027
Non-Small Cell Lung	A549/ATCC	< 0.01	< 0.01	< 0.005
Non-Small Cell Lung	EKVX	< 0.01	0.081	0.023
Non-Small Cell Lung	HOP-62	< 0.01	0.01	< 0.005
Non-Small Cell Lung	HOP-92	< 0.01	0.088	0.011
Non-Small Cell Lung	NCI-H226	< 0.01	0.0052	0.021
Non-Small Cell Lung	NCI-H23	< 0.01	0.022	< 0.005
Non-Small Cell Lung	NCI-H322M	< 0.01	0.021	< 0.005
Non-Small Cell Lung	NCI-H460	< 0.01	0.22	0.015
Non-Small Cell Lung	NCI-H522	< 0.01	< 0.01	< 0.005
Ovarian	IGROV1	< 0.01	0.052	0.013

Ovarian	OVCAR-3	< 0.01	0.05	0.012
Ovarian	OVCAR-4	< 0.01	0.048	< 0.005
Ovarian	OVCAR-5	< 0.01	0.051	0.017
Ovarian	OVCAR-8	< 0.01	0.033	0.0076
Ovarian	SK-OV-3	< 0.01	0.35	0.018
Prostate	DU-145	< 0.01	0.22	0.017
Prostate	PC-3	< 0.01	0.018	< 0.005
Renal	786-0	< 0.01	0.047	0.0065
Renal	A498	< 0.01	0.10	0.016
Renal	ACHN	< 0.01	0.19	0.039
Renal	CAKI-1	< 0.01	0.064	0.038
Renal	RXF 393	< 0.01	0.011	< 0.005
Renal	SN12C	< 0.01	< 0.01	< 0.005
Renal	TK-10	< 0.01	0.029	0.01
Renal	UO-31	< 0.01	0.016	0.063

Table 7: In Vitro Growth Inhibition (GI₅₀) of NCI Human Transformed Cell Lines of Several Compounds of the Current Invention and Olomoucine.

Cancer Type	Cell Line	74 GI ₅₀ (μM)	78 GI ₅₀ (μM)	77 GI ₅₀ (μM)	Olomoucine GI ₅₀ (μM)
Breast	BT-549	0.16	0.04	< 0.01	79
Breast	HS 578T	< 0.01	--	< 0.01	63
Breast	MCF7	< 0.01	< 0.01	0.03	50
Breast	MDA-MB-231/ATCC	< 0.01	< 0.01	0.04	100
Breast	MDA-MB-435	--	--	--	63
Breast	MDA-N	< 0.01	< 0.01	0.01	79
Breast	NCI/ADR-RES	0.24	14.45	0.03	100
Breast	T-47D	< 0.01	0.03	0.01	63
CNS	SF-268	< 0.01	--	< 0.01	50
CNS	SF-295	< 0.01	0.21	0.04	79
CNS	SF-539	0.07	--	0.22	32
CNS	SNB-19	< 0.01	< 0.01	0.03	63
CNS	SNB-75	< 0.01	< 0.01	< 0.01	25
CNS	U251	< 0.01	0.02	0.09	50
Colon	COLO 205	< 0.01	< 0.01	0.02	32
Colon	HCC-2998	--	< 0.01	--	63
Colon	HCT-116	< 0.01	0.03	0.05	40
Colon	HCT-15	< 0.01	1.48	< 0.01	40
Colon	HT29	< 0.01	< 0.01	< 0.01	63
Colon	KM12	< 0.01	< 0.01	< 0.01	40
Colon	SW-620	< 0.01	< 0.01	< 0.01	40
Leukemia	CCRF-CEM	< 0.01	--	< 0.01	40
Leukemia	HL-60(TB)	--	< 0.01	--	40
Leukemia	K-562	< 0.01	0.02	0.02	100
Leukemia	MOLT-4	< 0.01	< 0.01	0.01	63
Leukemia	RPMI-8226	< 0.01	< 0.01	< 0.01	50
Leukemia	SR	< 0.01	--	0.02	25
Melanoma	LOX IMVI	< 0.01	--	0.04	32

- 233 -

Melanoma	M14	< 0.01	< 0.01	< 0.01	100
Melanoma	MALME-3M	0.01	0.01	0.05	100
Melanoma	SK-MEL-2	0.06	0.02	0.51	100
Melanoma	SK-MEL-28	< 0.01	0.01	< 0.01	50
Melanoma	SK-MEL-5	0.06	0.10	0.08	40
Melanoma	UACC-257	< 0.01	0.02	0.02	79
Melanoma	UACC-62	0.04	0.03	0.12	32
Non-Small Cell Lung	A549/ATCC	< 0.01	< 0.01	< 0.01	50
Non-Small Cell Lung	EKVX	--	0.05	--	100
Non-Small Cell Lung	HOP-62	< 0.01	0.02	< 0.01	32
Non-Small Cell Lung	HOP-92	0.03	--	0.13	50
Non-Small Cell Lung	NCI-H226	--	0.02	--	50
Non-Small Cell Lung	NCI-H23	< 0.01	0.01	0.01	79
Non-Small Cell Lung	NCI-H322M	< 0.01	< 0.01	< 0.01	63
Non-Small Cell Lung	NCI-H460	< 0.01	0.05	0.22	63
Non-Small Cell Lung	NCI-H522	< 0.01	< 0.01	< 0.01	40
Ovarian	IGROV1	< 0.01	< 0.01	0.09	40
Ovarian	OVCAR-3	< 0.01	0.03	0.02	79
Ovarian	OVCAR-4	< 0.01	0.02	< 0.01	100
Ovarian	OVCAR-5	0.03	< 0.01	0.04	40
Ovarian	OVCAR-8	< 0.01	0.02	0.02	63
Ovarian	SK-OV-3	0.22	0.06	0.19	100
Prostate	DU-145	0.02	0.06	0.13	40
Prostate	PC-3	< 0.01	< 0.01	0.02	100
Renal	786-0	< 0.01	0.04	0.03	63
Renal	A498	0.03	0.03	0.03	32
Renal	ACHN	0.03	0.32	0.11	25
Renal	CAKI-1	--	0.79	--	32
Renal	RXF 393	< 0.01	< 0.01	< 0.01	20
Renal	SN12C	< 0.01	< 0.01	< 0.01	100
Renal	TK-10	< 0.01	0.07	0.05	63
Renal	UO-31	0.01	0.17	< 0.01	32

The following data in Table 8 summarize the *in vivo* properties of several compounds of the current invention. These data were cooperatively obtained at the National Cancer Institute in their Hollow Fiber Assay according to published procedures (Hollingshead, M.G., et al "In Vivo Cultivation of Tumor Cells in Hollow Fibers," *Life Sciences*, 1995, 57(2), 131-141 which is hereby incorporated by reference).

Table 8: In Vivo Evaluation of Several Compounds of the Current Invention.

Compound	MTD (mg/kg)	IP Score	SC score	Cell Kill	Cell Types Killed
73	100	2	0	N	--
17	100	8	0	Y	H522
38	100	0	4	N	--
78	6.3	34	0	Y	H23, H522, OVCAR3, SF29
79	6.3	26	6	N	--
86	6.3	38	0	Y	OVCAR 3, OVCAR 5, H522
87	25	30	2	Y	H522
12	3.1	26	4	Y	H522, MDA-MB-435
93	25	22	8	Y	OVCAR-3
94	50	22	2	Y	COLO205, OVCAR-3, H522
103	6.3	38	6	Y	OVCAR 3, OVCAR 5, H522, MDA-MB-435, SF295
109	400	18	10	Y	H522
112	50	18	2	Y	OVCAR-3
113	200	18	4	Y	H23, H522, OVCAR-3
110	50	14	0	Y	OVCAR-3
124	25	28	0	Y	H23, H522, OVCAR-3, MDA-MB-435

5

The following data in Table 9 summarize the *in vivo* properties of several compounds of the current invention. The protocol for the experiment is as follows. The dose-range finding study consists of four groups of three athymic mice each (four dose levels). The compound is administered on the basis of individual animal body weight. The route is intraperitoneal (IP) and the treatment schedule is daily for 14 days (qd x 14) or once every 4 days for 12 days (q4d x 3). The mice were observed for survival, and body weights recorded weekly.

The efficacy study consists of three compound-treated groups (six mice/group), a positive control-treated group (six mice), and a vehicle-treated control group of 12 mice. Test compounds were administered IP under the treatment schedules listed above (qd x 14 or q4d x 3), whereas the positive control agent (Taxol) was administered intravenously (IV) at a dosage level of 15 mg/kg/dose for five consecutive days (qd x 5). All agents were administered on the basis of individual animal body weight. Treatment began when the implanted tumors were approximately 100 mg in size (range of 65 to 200 mg). The mice were observed daily for survival. Each tumor was measured by caliper in two dimensions and converted to tumor mass using the formula for a prolate ellipsoid ($a \times b^2/2$) and assuming unit density. Tumor measurements and animal body weights were recorded twice weekly. Antitumor activity was assessed by the delay in tumor growth of the treated

- 235 -

groups in comparison to the vehicle-treated control group, partial and complete regressions, and tumor-free survivors.

Table 9: In Vivo Evaluation of Several Compounds of the Current Invention.


Compound	Dose (mg/kg)	Route	Schedule	Tumor free/total	T-C (days)
78	0.5	IP	qd x 14	0/6	0.8
78	0.33	IP	qd x 14	0/6	1.8
78	0.22	IP	qd x 14	0/6	-0.8
78	1.5	IP	q4d x 3	0/6	2.5
78	1.0	IP	q4d x 3	0/6	1.1
78	0.67	IP	q4d x 3	0/6	2.0
12	0.6	IP	q4d x 3	0/6	0.3
12	0.4	IP	q4d x 3	0/6	0.2
12	0.27	IP	q4d x 3	0/6	1.2
87	15	IP	q4d x 3	0/6	2.5
87	10	IP	q4d x 3	0/6	2.9
87	6.7	IP	q4d x 3	0/6	1.4

Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is solely for that purpose, and variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

- 236 -

WHAT IS CLAIMED:

1. A compound of the following formula:

**Formula I**

wherein:

R_1 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_6 -straight chain alkyl;
 C_2 - C_6 -straight alkenyl chain;
 C_3 - C_6 -branched alkyl chain;
 C_3 - C_6 -branched alkenyl chain;
 C_3 - C_7 -cycloalkyl;
 CH_2 -(C_3 - C_7 -cycloalkyl);
 CH_2CF_3 ;
 $CH_2CH_2CF_3$; and
 $CH(CF_3)_2$;

the combination of X, D, and Q are either:

$D=Q=N$ and $X=CH$; or
 $D=X=N$ and $Q=CH$; or
 $Q=X=N$ and $D=CH$; or
 $Q=N$ and $D=X=CH$;

V= NH;
 O;
 S; or
 CH_2 ;

R_2 = phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O)_2R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F, Cl, Br, CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;
 1-naphthyl;

- 237 -

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;

3-pyridyl;

4-pyridyl;

2-pyrimidyl;

4-pyrimidyl;

5-pyrimidyl;

thiophene-2-yl;

thiophene-3-yl;

2-furanyl;

3-furanyl;

oxazol-2-yl;

oxazol-4-yl;

oxazol-5-yl;

thiazol-2-yl;

thiazol-4-yl;

thiazol-5-yl;

imidazol-2-yl;

imidazol-4-yl;

pyrazol-3-yl;

pyrazol-4-yl;

isoxazol-3-yl;

isoxazol-4-yl;

isoxazol-5-yl;

isothiazol-3-yl;

isothiazol-4-yl;

isothiazol-5-yl;

1,3,4-thiadiazol-2-yl;

benzo[b]furan-2-yl;

benzo[b]thiophene-2-yl;

2-pyrrolyl;

3-pyrrolyl;

1,3,5-triazin-2-yl;

pyrazin-2-yl;

pyridazin-3-yl;

pyridazin-4-yl;

2-quinolinyl;

3-quinolinyl;

4-quinolinyl;

1-isoquinolinyl;

3-isoquinolinyl; and

4-isoquinolinyl; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

- 238 -

H;
C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh; and
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
above in R₂;

R₄= H;
C₁-C₄-straight chain alkyl; or
C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a
5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

A= CH₂;
(CH₂)₂;
(CH₂)₃;
OCH₂CH₂; or
CHCH₃;

Y= H;
OR₁;
N(R₁)₂;
N(R₁)C(O)R₃;
N(R₁)C(O)R₅;
N(R₁)C(O)CH(R₆)NH₂;
N(R₁)SO₂R₃;
N(R₁)C(O)NHR₃; or
N(R₁)C(O)OR₆;

R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh; or
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
above in R₂;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein

- 239 -

R_3 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_4 -straight chain alkyl; and
 C_3 - C_4 -branched chain alkyl;

Y= H;
 OR_1 ;
 $N(R_1)_2$;
 $N(R_1)C(O)R_3$;
 $N(R_1)SO_2R_3$; or
 $N(R_1)C(O)NHR_3$.

3. A compound according to claim 1, wherein
two of X, D, and Q = N;


R_3 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_4 -straight chain alkyl; and
 C_3 - C_4 -branched chain alkyl;

Y= H;
 OR_1 ;
 $N(R_1)_2$;
 $N(R_1)C(O)R_3$;
 $N(R_1)SO_2R_3$; or
 $N(R_1)C(O)NHR_3$;

or a pharmaceutically acceptable salt thereof.

4. A compound of the following formula:



Formula III

wherein:

- 240 -

R_1 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_6 -straight chain alkyl;
 C_2 - C_6 -straight alkenyl chain;
 C_3 - C_6 -branched alkyl chain;
 C_3 - C_6 -branched alkenyl chain;
 C_3 - C_7 -cycloalkyl;
 CH_2 -(C_3 - C_7 -cycloalkyl);
 CH_2CF_3 ;
 $CH_2CH_2CF_3$; and
 $CH(CF_3)_2$;

the combination of X, D, and Q are either:

$D=Q=N$ and $X=CH$; or
 $D=X=N$ and $Q=CH$; or
 $Q=X=N$ and $D=CH$; or
 $Q=N$ and $D=X=CH$;

V= NH;
 O;
 S; or
 CH_2 ;

R_2 = phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F, Cl, Br, CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;
 1-naphthyl;
 2-naphthyl;
 heterocycles selected from the group consisting of:
 2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;

- 241 -

imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

n= 0-3;

A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;

R₃ are the same or different and independently selected from the group consisting of:
 H;

- 242 -

C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh; and
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

R₅= C₃-C₇-cycloalkyl;

R₆ = C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl; or
C₂-C₄-alkenyl chain;

or a pharmaceutically acceptable salt thereof.

5. A compound according to claim 4, wherein two of X, D, and Q = N.
6. A method of treating a mammal with a disorder mediated by elevated levels of cell proliferation compared to a healthy mammal comprising:
administering a therapeutically effective amount of the compound of claim 1 to the mammal under conditions effective to treat the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal.
7. The method of claim 6, wherein the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal is cancer, wherein the cancer is selected from the group consisting of breast cancer, colon cancer, central nervous system cancer, leukemia, melanoma, lung cancer, ovarian cancer, prostate cancer, and renal cancer.
8. The method of claim 7, wherein the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal is restenosis.
9. The method of claim 7, wherein the mammal is human.
10. A pharmaceutical composition of matter comprising the compound of claim 1 and one or more pharmaceutical excipients.

- 243 -

11. A method of treating a mammal with a disorder mediated by elevated levels of cell proliferation compared to a healthy mammal comprising:

administering a therapeutically effective amount of the compound of claim 4 to the mammal under conditions effective to treat the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal.


12. The method of claim 11, wherein the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal is cancer, wherein the cancer is selected from the group consisting of breast cancer, colon cancer, central nervous system cancer, leukemia, melanoma, lung cancer, ovarian cancer, prostate cancer, and renal cancer.

13. The method of claim 11, wherein the disorder mediated by elevated levels of cell proliferation compared to a healthy mammal is restenosis.

14. The method of claim 11, wherein the mammal is human.

15. A pharmaceutical composition of matter comprising the compound of claim 4 and one or more pharmaceutical excipients.

16. A process for preparation of a purine derivative compound of the formula:



Formula I

wherein:

R₁ are the same or different and independently selected from the group consisting of:
H;

- 244 -

C₁-C₆-straight chain alkyl;
C₂-C₆-straight alkenyl chain;
C₃-C₆-branched alkyl chain;
C₃-C₆-branched alkenyl chain;
C₃-C₇-cycloalkyl;
CH₂-(C₃-C₇-cycloalkyl);
CH₂CF₃;
CH₂CH₂CF₃; and
CH(CF₃)₂;

the combination of X, D, and Q are either:

D=Q=N, and X=CH; or
D=X=N, and Q=CH; or
Q=X=N, and D=CH; or
Q=N, and D=X=CH;

V= NH;
O;
S; or
CH₂;

R₂= phenyl;
substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;
3-pyridyl;
4-pyridyl;
2-pyrimidyl;
4-pyrimidyl;
5-pyrimidyl;
thiophene-2-yl;
thiophene-3-yl;
2-furanyl;
3-furanyl;
oxazol-2-yl;
oxazol-4-yl;
oxazol-5-yl;
thiazol-2-yl;
thiazol-4-yl;
thiazol-5-yl;
imidazol-2-yl;
imidazol-4-yl;
pyrazol-3-yl;

- 245 -

pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinoliny;
 3-quinoliny;
 4-quinoliny;
 1-isoquinoliny;
 3-isoquinoliny; and
 4-isoquinoliny; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined

above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

A= CH₂;
 (CH₂)₂;

- 246 -

(CH₂)₃;
OCH₂CH₂; or
CHCH₃;

Y= H;
OR₁;
N(R₁)₂;
N(R₁)C(O)R₃;
N(R₁)C(O)R₅;
N(R₁)C(O)CH(R₆)NH₂;
N(R₁)SO₂R₃;
N(R₁)C(O)NHR₃; or
N(R₁)C(O)OR₆;


R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh; or
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
above in R₂;

or a pharmaceutically acceptable salt thereof;

said process comprising:

reacting a first intermediate compound of the formula:



Formula IX

where Z= Br or I

with a compound of the formula: R₂-B(OH)₂, R₂-Sn(n-Bu)₃, or R₂-Sn(Me)₃, or
mixtures thereof, under conditions effective to form the purine derivative compound.


- 247 -

17. A process according to claim 16, wherein if Y is NHR_1 , said process further comprises:

reacting the purine derivative compound with $\text{R}_3\text{C}(\text{O})\text{Cl}$ or $\text{R}_5\text{C}(\text{O})\text{Cl}$ or $\text{R}_3\text{SO}_2\text{Cl}$ or R_3NCO or $\text{R}_6\text{OC}(\text{O})\text{Cl}$ under conditions effective to form a final product having the same formula as the purine derivative compound except that Y is $\text{NR}_1\text{C}(\text{O})\text{R}_3$ or $\text{NR}_1\text{C}(\text{O})\text{R}_5$ or $\text{NR}_1\text{SO}_2\text{R}_3$ or $\text{NR}_1\text{C}(\text{O})\text{NHR}_3$ or $\text{NR}_1\text{C}(\text{O})\text{OR}_6$.


18. A process according to claim 16 further comprising:

reacting a second intermediate compound of the formula:



Formula VII

with a second compound of the formula:



Formula VIII


wherein: $\text{V}_1 = \text{NH}_2$;
 OH ;
 SH ;

under conditions effective to form the first intermediate compound.

19. A process according to claim 18 further comprising:


reacting a third intermediate compound of the formula:

- 248 -


**Formula VI**

with a compound of the formula R_1 -Z under conditions effective to form the second intermediate compound.

20. A process according to claim 16, wherein the purine derivative compound has the formula:

**Formula III**

21. A process for preparation of a purine derivative compound of the formula:

**Formula I**

wherein:

- 249 -

R_1 are the same or different and independently selected from the group consisting of:

H;
 C_1 - C_6 -straight chain alkyl;
 C_2 - C_6 -straight alkenyl chain;
 C_3 - C_6 -branched alkyl chain;
 C_3 - C_6 -branched alkenyl chain;
 C_3 - C_7 -cycloalkyl;
 CH_2 -(C_3 - C_7 -cycloalkyl);
 CH_2CF_3 ;
 $CH_2CH_2CF_3$; and
 $CH(CF_3)_2$;

the combination of X, D, and Q are either:

$D=Q=N$, and $X=CH$; or
 $D=X=N$, and $Q=CH$; or
 $Q=X=N$, and $D=CH$; or
 $Q=N$, and $D=X=CH$;

V= NH;
 O;
 S; or
 CH_2 ;

R_2 = phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R_1 , OR_1 , SR_1 , $S(O)R_1$, $S(O_2)R_1$, NHR_1 , NO_2 , $OC(O)CH_3$, $NHC(O)CH_3$, F, Cl, Br, CF_3 , $C(O)R_1$, $C(O)NHR_1$, phenyl, and $C(O)NHCHR_1CH_2OH$;

1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;

- 250 -

imidazol-4-yl;
 pyrazol-3-yl;
 pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined

above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

- 251 -

A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂;
 CHCH₃;


Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;

R₃= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂;

or a pharmaceutically acceptable salt thereof, said process comprising:

reacting a first intermediate compound of the formula:




Formula XV


under reductive or hydrogenation conditions effective to form the purine derivative compound.

- 252 -

22. A process according to claim 21 further comprising:
reacting a second intermediate compound of the formula:


**Formula XIV**

with a second compound of the formula:

**Formula XII**

under conditions effective to form the first intermediate compound.


23. A process according to claim 22 further comprising:
reacting a third intermediate compound of the formula:

**Formula VII**

with a compound of the formula: $R_2-B(OH)_2$, $R_2-Sn(n-Bu)_3$, or $R_2-Sn(Me)_3$, or mixtures thereof, under conditions effective to form the second intermediate compound.


24. A process according to claim 21 further comprising:
reacting a fourth intermediate compound of the formula:

- 253 -


**Formula XIII**

with a compound of the formula: $R_2-B(OH)_2$, $R_2-Sn(n-Bu)_3$, or $R_2-Sn(Me)_3$, or mixtures thereof, under conditions effective to form the first intermediate compound.

25. A process according to claim 24 further comprising:
reacting a fifth intermediate compound of the formula:

**Formula VII**


with a compound of the formula:

**Formula XII**


under conditions effective to form the fourth intermediate compound.

26. A process according to claim 21, wherein the purine derivative compound has the formula:

- 254 -

**Formula III**

27. A process for preparation of a purine derivative compound of the formula:

**Formula I**

wherein:

R₁ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 CH₂CH₂CF₃; and
 CH(CF₃)₂;

the combination of X, D, and Q are either:

D=Q=N, and X=CH; or
 D=X=N, and Q=CH; or
 Q=X=N, and D=CH; or
 Q=N, and D=X=CH;

V= NH;
 O;

- 255 -

S; or
CH₂;

R₂= phenyl;
substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;

3-pyridyl;

4-pyridyl;

2-pyrimidyl;

4-pyrimidyl;

5-pyrimidyl;

thiophene-2-yl;

thiophene-3-yl;

2-furanyl;

3-furanyl;

oxazol-2-yl;

oxazol-4-yl;

oxazol-5-yl;

thiazol-2-yl;

thiazol-4-yl;

thiazol-5-yl;

imidazol-2-yl;

imidazol-4-yl;

pyrazol-3-yl;

pyrazol-4-yl;

isoxazol-3-yl;

isoxazol-4-yl;

isoxazol-5-yl;

isothiazol-3-yl;

isothiazol-4-yl;

isothiazol-5-yl;

1,3,4-thiadiazol-2-yl;

benzo[b]furan-2-yl;

benzo[b]thiophene-2-yl;

2-pyrrolyl;

3-pyrrolyl;

1,3,5-triazin-2-yl;

pyrazin-2-yl;

pyridazin-3-yl;

pyridazin-4-yl;

2-quinoliny;

3-quinoliny;

- 256 -

4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or
 substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

Y= H;
 OR₁;
 N(R₁)₂;
 N(R₁)C(O)R₃;
 N(R₁)C(O)R₅;
 N(R₁)C(O)CH(R₆)NH₂;
 N(R₁)SO₂R₃;
 N(R₁)C(O)NHR₃; or
 N(R₁)C(O)OR₆;


R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;

- 257 -


C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
 above in R₂; or a pharmaceutically acceptable salt thereof; said process
 comprising:

reacting a first intermediate compound of the formula:



Formula XIV

with a compound of the formula:



Formula VIII


where V₁= NH₂;
 OH; or
 SH;

under conditions effective to form the purine derivative compound.

28. A process according to claim 27, wherein if Y is NHR₁, said process further comprises:

reacting the purine derivative compound with R₃C(O)Cl or R₅C(O)Cl or
 R₃SO₂Cl or R₃NCO or R₆OC(O)Cl under conditions effective to form a final product
 having the same formula as the purine derivative compound except that Y is
 NR₁C(O)R₃ or NR₁C(O)R₅ or NR₁SO₂R₃ or NR₁C(O)NHR₃ or NR₁C(O)OR₆.


29. A process according to claim 27 further comprising:
reacting a second intermediate compound of the formula:



Formula XVIII

with a compound of the formula R₁-Z under conditions effective to form the first intermediate compound.


30. A process according to claim 29 further comprising:
reacting a third intermediate compound of the formula:



Formula XVII

with a compound of the formula 2,6-dichloropurine (**Formula IV**) under conditions effective to form the second intermediate compound.

31. A process according to claim 30 further comprising:
reacting a fourth intermediate compound of the formula:




Formula V

- 259 -


with a compound of the formula: $R_2-B(OH)_2$, $R_2-Sn(n-Bu)_3$, or $R_2-Sn(Me)_3$, or mixtures thereof, under conditions effective to form the third intermediate compound.

32. A process according to claim 28, wherein the purine derivative compound has the formula:



Formula III

33. A process for preparation of a purine derivative compound of the formula:



Formula XX

wherein:

R_1 are the same or different and independently selected from the group consisting of:

- H;
- C_1 - C_6 -straight chain alkyl;
- C_2 - C_6 -straight alkenyl chain;
- C_3 - C_6 -branched alkyl chain;
- C_3 - C_6 -branched alkenyl chain;
- C_3 - C_7 -cycloalkyl;
- CH_2 -(C_3 - C_7 -cycloalkyl);
- CH_2CF_3 ;
- $CH_2CH_2CF_3$; and
- $CH(CF_3)_2$;

- 260 -

the combination of X, D, and Q are either:

D=Q=N and X=CH; or

D=X=N and Q=CH; or

Q=X=N and D=CH; or

Q=N and D=X=CH;

V= NH;

O;

S; or

CH₂;

R₂= phenyl;

substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;

3-pyridyl;

4-pyridyl;

2-pyrimidyl;

4-pyrimidyl;

5-pyrimidyl;

thiophene-2-yl;

thiophene-3-yl;

2-furanyl;

3-furanyl;

oxazol-2-yl;

oxazol-4-yl;

oxazol-5-yl;

thiazol-2-yl;

thiazol-4-yl;

thiazol-5-yl;

imidazol-2-yl;

imidazol-4-yl;

pyrazol-3-yl;

pyrazol-4-yl;

isoxazol-3-yl;

isoxazol-4-yl;

isoxazol-5-yl;

isothiazol-3-yl;

isothiazol-4-yl;

isothiazol-5-yl;

1,3,4-thiadiazol-2-yl;

benzo[b]furan-2-yl;

- 261 -

benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined

above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

A= CH₂;
 (CH₂)₂;
 (CH₂)₃;
 OCH₂CH₂; or
 CHCH₃;

Y= NR₁C(O)R₃;
 NR₁SO₂R₃;
 NR₁C(O)NHR₃; or
 NR₁C(O)OR₆

- 262 -

$R_5 =$ C₃-C₇-cycloalkyl;

$R_6 =$ C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; or
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined above in R₂;

or a pharmaceutically acceptable salt thereof;


said process comprising:

reacting a first intermediate compound having the same formula as the purine derivative compound except that Y=NHR₁, with R₃COCl or R₅COCl or R₃SO₂Cl or R₃NCO or R₆OC(O)Cl under conditions effective to form the purine derivative compound.

34. A process according to claim 33 further comprising:

reacting a second intermediate compound having the same formula as the first intermediate compound except that Y is NH₂, with R₈CH₂Z or R₈CHO under conditions effective to form the first intermediate compound where Y=NHR₁ or N(R₁)₂, wherein Z is Br or I and R₈=C₁-C₅-straight chain alkyl; C₂-C₅-straight alkenyl chain; C₃-C₅-branched alkyl chain; C₃-C₅-branched alkenyl chain; C₃-C₇-cycloalkyl; CF₃; CH₂CF₃.

35. A process for preparation of a purine derivative compound of the formula:



Formula XIX

wherein:

R₁ are the same or different and independently selected from the group consisting of:
 H;

- 263 -

C₁-C₆-straight chain alkyl;
 C₂-C₆-straight alkenyl chain;
 C₃-C₆-branched alkyl chain;
 C₃-C₆-branched alkenyl chain;
 C₃-C₇-cycloalkyl;
 CH₂-(C₃-C₇-cycloalkyl);
 CH₂CF₃;
 CH₂CH₂CF₃; and
 CH(CF₃)₂;

the combination of X, D, and Q are either:

D=Q=N and X=CH; or
 D=X=N and Q=CH; or
 Q=X=N and D=CH; or
 Q=N and D=X=CH;

V= NH;
 O;
 S; or
 CH₂;

R₂= phenyl;
 substituted phenyl, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of R₁, OR₁, SR₁, S(O)R₁, S(O₂)R₁, NHR₁, NO₂, OC(O)CH₃, NHC(O)CH₃, F, Cl, Br, CF₃, C(O)R₁, C(O)NHR₁, phenyl, and C(O)NHCHR₁CH₂OH;

1-naphthyl;

2-naphthyl;

heterocycles selected from the group consisting of:

2-pyridyl;
 3-pyridyl;
 4-pyridyl;
 2-pyrimidyl;
 4-pyrimidyl;
 5-pyrimidyl;
 thiophene-2-yl;
 thiophene-3-yl;
 2-furanyl;
 3-furanyl;
 oxazol-2-yl;
 oxazol-4-yl;
 oxazol-5-yl;
 thiazol-2-yl;
 thiazol-4-yl;
 thiazol-5-yl;
 imidazol-2-yl;
 imidazol-4-yl;
 pyrazol-3-yl;

- 264 -

pyrazol-4-yl;
 isoxazol-3-yl;
 isoxazol-4-yl;
 isoxazol-5-yl;
 isothiazol-3-yl;
 isothiazol-4-yl;
 isothiazol-5-yl;
 1,3,4-thiadiazol-2-yl;
 benzo[b]furan-2-yl;
 benzo[b]thiophene-2-yl;
 2-pyrrolyl;
 3-pyrrolyl;
 1,3,5-triazin-2-yl;
 pyrazin-2-yl;
 pyridazin-3-yl;
 pyridazin-4-yl;
 2-quinolinyl;
 3-quinolinyl;
 4-quinolinyl;
 1-isoquinolinyl;
 3-isoquinolinyl; and
 4-isoquinolinyl; or

substituted heterocycle, wherein the substituents (1-2 in number) are in any position and are independently selected from the group consisting of Br, Cl, F, R₁, and C(O)CH₃;

R₃ are the same or different and independently selected from the group consisting of:

H;
 C₁-C₄-straight chain alkyl;
 C₃-C₄-branched chain alkyl;
 C₂-C₄-alkenyl chain;
 (CH₂)_nPh; and
 (CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined

above in R₂;

R₄= H;
 C₁-C₄-straight chain alkyl; or
 C₃-C₄-branched chain alkyl;

R₃ and R₄ can be linked together by a carbon chain to form with intervening atoms a 5-8-membered saturated or unsaturated ring;

n₁= 0-3;

n= 0-3;

A= CH₂;
 (CH₂)₂;

- 265 -

(CH₂)₃;
OCH₂CH₂; or
CHCH₃;

Y= NHC(O)CH(R₆)NH₂;

R₅= C₃-C₇-cycloalkyl;

R₆= C₁-C₄-straight chain alkyl;
C₃-C₄-branched chain alkyl;
C₂-C₄-alkenyl chain;
(CH₂)_nPh; or
(CH₂)_n-substituted phenyl, wherein the phenyl substituents are as defined
above in R₂;

or a pharmaceutically acceptable salt thereof;

said process comprising:

reacting a first intermediate compound having the same formula as the purine
derivative compound except that Y is NH₂, with a compound of the formula:
PNHCH(R₆)CO₂H under conditions effective to form the purine derivative compound
after a suitable deprotection strategy,

wherein

P= C(O)OtBu;
C(O)OCH₂Ph;
Fmoc
Benzyl; or
Alloc.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 March 2003 (20.03.2003)

PCT

(10) International Publication Number
WO 03/022219 A3

(51) International Patent Classification⁷: C07D 473/22,
473/18, A61K 31/52, 31/522, A61P 35/00, 35/02

(21) International Application Number: PCT/US02/28731

(22) International Filing Date:
9 September 2002 (09.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/950,543 11 September 2001 (11.09.2001) US

(71) Applicant: ALBANY MOLECULAR RESEARCH,
INC. [US/US]; 21 Corporate Circle, Albany, NY
12212-5098 (US).

(72) Inventor: TROVA, Michael, Peter; 3002 Morgan Court,
Schenectady, NY 12306 (US).


(74) Agents: GOLDMAN, Michael, L. et al.; Nixon Peabody
LLP, P.O. Box 31051, Clinton Square, Rochester, NY
14603-1051 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,
VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: BIARYLAMINO PURINE ANTIPROLIFERATIVE AGENTS



(57) Abstract: The compounds of the present invention are 2,6,9-trisubstituted purine derivatives which are inhibitors of cyclin/cdk complexes. The compounds of the current invention also are potent inhibitors of human cellular proliferation. As such, the compounds of the present invention constitute pharmaceutical compositions with a pharmaceutically acceptable carrier. Such compounds are useful in treating a disorder mediated by elevated levels of cell proliferation in a mammal compared to a healthy mammal by administering to such mammal an effective amount of the compound. Examples of the compounds of the present invention are represented by the following chemical structures; with X, Y, D, Q, V, A, R₁, R₂, R₃, R₄, and n₁ defined herein.





Published:

— *with international search report*

(88) Date of publication of the international search report:

13 November 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/28731

A. CLASSIFICATION OF SUBJECT MATTER		
IPC(7) : C07D 473/22, 473/18; A61K 31/52, 31/522 A61P 35/00, 35/02		
US CL : 544/276, 277; 514/263.2, 263.21		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
U.S. : 544/276, 277; 514/263.2, 263.21		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,E	US 6479487 B1 (DUMONT ET AL) 12 November 2002, see Formula I.	1-35
A,P	US 6413974 B1 (DUMONT ET AL) 02 July 2002, see Formula II.	1-35
A	FR 2793794 (Hoechst MARION ROUSSEL) 24 November 2000	1-35
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"B" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
06 January 2003 (06.01.2003)	15 JUL 2003	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703)305-3230	Authorized officer Mark L. Berch Telephone No. 703-308-1235	

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

PCT/US02/28731

Continuation of Item 4 of the first sheet:

Title too long. New Title:

Biaryl amino Purine
antiproliferative agents